

UNIVERSIDAD COMPLUTENSE DE MADRID

FACULTAD DE CIENCIAS GEOLÓGICAS
DEPARTAMENTO DE PALEONTOLOGÍA



TESIS DOCTORAL

**Cricetodontini (Rodentia, Mammalia) del Mioceno medio y superior del área de
Daroca (Aragón, España):
sistemática y filogenia**

MEMORIA PARA OPTAR AL GRADO DE DOCTORA

PRESENTADA POR

Paloma López Guerrero

Directores

María Ángeles Álvarez Sierra
Israel García Paredes

Madrid, 2014

CRICETODONTINI (RODENTIA, MAMMALIA) DEL MIOCENO MEDIO Y SUPERIOR DEL ÁREA DE DAROCA (ARAGÓN, ESPAÑA): SISTEMÁTICA Y FILOGENIA

Cricetodontini (Rodentia, Mammalia) from the middle and upper Miocene in the Daroca area (Aragón, Spain): Systematics & Phylogeny

PALOMA LÓPEZ GUERRERO

DIRECTORES DE TESIS

DRA. MARÍA ÁNGELES ÁLVAREZ SIERRA
DR. ISRAEL GARCÍA PAREDES

TESIS DOCTORAL

**MADRID
2014**



DEPTO. DE PALEONTOLOGÍA
FACULTAD DE GEOLOGÍA
UNIVERSIDAD COMPLUTENSE DE MADRID




CSIC

CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS

**DEPTO. DE GEOLOGÍA SEDIMENTARIA Y CAMBIO
MEDIOAMBIENTAL**
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS
INSTITUTO DE GEOCIENCIAS-IGEO

**CRICETODONTINI (RODENTIA, MAMMALIA) DEL MIOCENO MEDIO
Y SUPERIOR DEL ÁREA DE DAROCA (ARAGÓN, ESPAÑA):
SISTEMÁTICA Y FILOGENIA**



MEMORIA DE TESIS DOCTORAL PRESENTADA POR
PALOMA LÓPEZ GUERRERO
BAJO LA DIRECCIÓN DE LOS DOCTORES
MARÍA ÁNGELES ÁLVAREZ SIERRA E ISRAEL GARCÍA PAREDES



UNIVERSIDAD COMPLUTENSE DE MADRID
FACULTAD DE GEOLOGÍA
DEPARTAMENTO DE PALEONTOLOGÍA



CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS
UNIVERSIDAD COMPLUTENSE DE MADRID
INSTITUTO DE GEOCIENCIAS-IGEO
DEPARTAMENTO DE GEOLOGÍA SEDIMENTARIA Y CAMBIO
MEDIOAMBIENTAL



Vº Bº DE LA DIRECTORA DE TESIS
MARÍA ÁNGELES ÁLVAREZ SIERRA

Vº Bº DEL DIRECTOR DE TESIS
ISRAEL GARCÍA PAREDES

FDO.
PALOMA LÓPEZ GUERRERO

A mi familia y amigos
A Toño

“The past history of our globe must be explained by what can be seen to be happening now. No powers are to be employed that are not natural to the globe, no action to be admitted except those of which we know the principle.”

James Hutton
Theory of the Earth (1785)

“The most consequential change in man's view of the world, of living nature and of himself came with the introduction, over a period of some 100 years beginning only in the 18th century, of the idea of change itself, of change over periods of time: in a word, of evolution.”

Ernst Mayr
Evolution, Scientific American (1978)

“La ciencia se construye intuitivamente, tenemos lagunas de lógica, pasamos años deambulando en la nada, nos frustramos, nos tiramos de los pelos. Éste es realmente el poder de la genialidad: La fuerza de voluntad para cometer los errores necesarios que te conducen a la respuesta correcta.”

Michio Kaku

–Quien no haya experimentado la irresistible atracción de la ciencia no podrá comprender su tiranía.–

Mary Shelley
Frankenstein (1818)

Esta tesis ha sido financiada gracias a una beca predocotoral del programa de Personal Investigador en Formación de la Universidad Complutense de Madrid y por los proyectos CGL2004-02094/BTE, CGL2007-65208, CGL2008-04200/BTE, y CGL 2011-28877.

Índice



AGRADECIMIENTOS	i
PRESENTACIÓN	vii
ABSTRACT	ix
RESUMEN	xiv

I. INTRODUCCIÓN 1

Capítulo 1. Introducción	3
INTRODUCCIÓN A LOS ROEDORES	5
CRICÉTIDOS	6
TRIBU CRICETODONTINI	7
Capítulo 2. Objetivos	13
Capítulo 3. Contexto Geológico	17
INTRODUCCIÓN	19
GEOLOGÍA	20
CONTEXTO CRONOLÓGICO Y BIOESTRATIGRÁFICO	28

II. MATERIAL Y MÉTODOS 33

Capítulo 4. Material y Métodos	35
MATERIAL	37
MÉTODOS	40
Métodos de Campo	40
Métodos de laboratorio	42
Estudio del material	43
Método de estudio filogenético	54
Bibliografía (capítulos 1-4)	57

III. PALEONTOLOGÍA SISTEMÁTICA 75

Chapter 5. Revision of <i>Cricetodon soriae</i> (Rodentia, Mammalia), new data from the middle Aragonian (middle Miocene) of the Calatayud-Daroca basin (Zaragoza, Spain)	76
INTRODUCTION	81
MATERIAL & METHODS	84
SYSTEMATIC PALEONTOLOGY	88
Genus <i>Cricetodon</i> Lartet, 1851	88
<i>Cricetodon soriae</i> López Martínez et al., 2006.	89
<i>Cricetodon soriae</i> from the Calatayud-Daroca Basin	91
<i>Cricetodon soriae</i> from Madrid Basin	94
Comparison with other species of <i>Cricetodon</i>	99
FINAL REMARKS & CONCLUSIONS	104
Conclusions	105
LITERATURE CITED	107
Chapter 6. <i>Cricetodon</i> from the middle and late Aragonian of the Calatayud-Daroca Basin: a taxonomical description	115

INTRODUCTION	119
MATERIAL & METHODS	121
SYSTEMATIC PALEONTOLOGY	123
<i>Cricetodon sansaniensis</i> Lartet, 1851	123
<i>Cricetodon sansaniensis</i> from the Calatayud-Daroca Basin	124
<i>Cricetodon jotae</i> Mein & Freudenthal, 1971	130
<i>Cricetodon jotae</i> from the Calatayud-Daroca Basin	131
FINAL REMARKS AND CONCLUSIONS	138
LITERATURE CITED	141

Chapter 7. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution

145

INTRODUCTION	149
MATERIAL & METHODS	151
SYSTEMATIC PALEONTOLOGY	154
<i>Cricetodon nievei</i> sp. nov.	154
<i>Cricetodon nievei</i> from the Calatayud-Daroca Basin	156
Comparisons and discussion	160
FINAL REMARKS AND CONCLUSIONS	169
LITERATURE CITED	175

Chapter 8. Species of *Hispanomys* of the late Aragonian and early Vallesian (middle-late Miocene) of the Calatayud-Daroca Basin, Zaragoza, Spain

185

INTRODUCTION	189
MATERIAL & METHODS	191
SYSTEMATIC PALEONTOLOGY	194
Genus <i>Hispanomys</i> Mein & Freudenthal, 1971	194
<i>Hispanomys aguirrei</i> (Sesé Benito, 1977 in López Martínez et al., 1977)	195
<i>Hispanomys aguirrei</i> from the Calatayud-Daroca Basin	195
<i>Hispanomys lavocati</i> (Freudenthal, 1966)	206
<i>Hispanomys lavocati</i> from the Calatayud-Daroca Basin	206
<i>Hispanomys nombrevillae</i> (Freudenthal, 1966)	214
<i>Hispanomys nombrevillae</i> from the Calatayud-Daroca Basin	215
<i>Hispanomys aragonensis</i> (Freudenthal, 1966)	223
<i>Hispanomys aragonensis</i> from the Calatayud-Daroca Basin	224
FINAL REMARKS & CONCLUSIONS	239
LITERATURE CITED	240
APPENDIX 8.1	245
APPENDIX 8.2	263
APPENDIX 8.3	281

IV. BIOESTRATIGRAFÍA

299

Chapter 9. Cricetodontini from the Calatayud-Daroca Basin (Spain): an up-to-date of their stratigraphical distributions

301

INTRODUCTION	305
RECORDED SPECIES OF CRICETODONTINI	307
<i>Cricetodon soriae</i> López Martínez et al., 2006 in Hernández Fernández et al., 2006	307
<i>Cricetodon sansaniensis</i> Lartet, 1851	307
<i>Cricetodon jotae</i> Mein & Freudenthal, 1971	308
<i>Cricetodon nievei</i> sp. nov.	308
<i>Hispanomys aguirrei</i> (Sesé Benito, 1977 in López Martínez et al., 1977)	309
<i>Hispanomys lavocati</i> (Freudenthal, 1966)	309
<i>Hispanomys nombrevillae</i> (Freudenthal, 1966)	309
<i>Hispanomys aragonensis</i> (Freudenthal, 1966)	310

CONCLUSIONS	315
LITERATURE	316
CITED V.	321
FILOGENIA	
Chapter 10. First phylogenetic approach to Cricetodontini (Rodentia, Mammalia) from the Miocene and Pliocene of Europe and Asia Minor	323
INTRODUCTION	327
MATERIAL & METHODS	329
RESULTS & DISCUSSION	335
Results	335
Discussion	336
Cricetodontini from the Calatayud-Daroca Basin	338
Comments on systematics	340
Paleobiogeography and Evolutionary History of the Cricetodontini	342
FINAL REMARKS & CONCLUSIONS	348
LITERATURE CITED	350
APPENDIX 10.1	361
APPENDIX 10.2	365
VI. RESUMEN Y CONCLUSIONES	379
Chapter 11. Resumen y conclusiones	381
Conclusiones taxonómicas	384
Conclusiones bioestratigráficas	388
Conclusiones filogenéticas	389

..... AGRADECIMIENTOS

Al igual que un artículo científico, esta tesis tiene varios autores. Por orden, las autorías de mayor importancia dentro de un artículo son por un lado, los primeros firmantes, autores intelectuales, aportadores de ideas y soluciones así como ejecutores responsables de que el artículo salga adelante y por otro, los últimos, ideólogos, canalizadores y apoyo sustancial.

En este caso, los primeros puestos deben reservarse a mis directores, la Dra. María de los Ángeles Álvarez Sierra y el Dr. Israel García Paredes. Me han enseñado paciencia, dignidad, honestidad y humildad para formarme como investigadora. Tengo que agradecer la oportunidad que me brindó Marián que nunca ha demostrado flaqueza en este proyecto, por mucho que le llevase el caos o el barroquismo hecho Word a su despacho. La Dedicación que tiene a la ciencia y a la docencia por partes iguales, su Sacrificio y el Sentido de la Justicia que posee, han despertado en mí lo mejor que puede inspirar un director de tesis, admiración. Además, siempre he contado con el apoyo de Israel, gran científico forjado en la excelencia y magnífico trabajador. Su compromiso con este mundo va más allá de estar 12 horas mirando un ordenador o a través de la lupa, su compromiso con la honestidad y la “NoVentaDeMotos” me ha enseñado la grandeza de lo sencillo.

Y los últimos firmantes deben ser mis padres, Francisco e Isabel, ellos me han dado las armas y herramientas, los valores, me han hecho ser lo que soy y han sido el gran soporte de esta tesis. Ellos han sido los responsables de todas las cosas buenas que tiene el pasar por este viaje.

Las enseñanzas de unos y otros se entremezclan y no es difícil reconocer que tanto en la vida como en la ciencia hay que ser pacientes, honestos, humildes y personas dignas.

Ahora entra el turno de los demás firmantes: colegas y amigos que se han comprometido tanto con esta tesis que no se puede discernir qué es suyo y qué es nuestro. Entre ellos, estoy especialmente agradecida al Dr. Pablo Peláez-Campomanes, sabio consejero, que no dudaba en acudir al rescate cada vez que le robaba un día en el museo y que se apasionaba tanto por este tema que nos contagiaba a los demás; incluso cuando no podíamos aguantar el mirar un solo diente más. *Also, I would like to express my most sincere gratitude to Dr. Lars van den Hoek Ostende, my Dutch*

family, your heart is as big as yourself. You had made me feel like I belong to Leiden and that I have a friend there. Not only for making me laugh (practically every minute) but also for your tremendous support and friendship. Yours is the phrase: "Friends are more important than fossils" and I am always trying to follow it. También agradezco profundamente el apoyo de mis compañeras. Las futuras doctoras: Adriana Oliver y Verónica Hernández Ballarín. Adriana y yo hemos ido juntas paso a paso y me siento privilegiada por haber contado con su apoyo en todos los momentos, no sólo los difíciles. Además, he podido aprender de ella la rigurosidad y profesionalidad con la que hace tu trabajo. También tengo que agradecer el tiempo que he compartido con Verónica, la sorpresa. La tranquilidad y serenidad que aporta han sido un bálsamo para las desesperaciones.

Afortunadamente, he podido compartir trabajo, despacho o conversaciones con muchos más autores nacionales e internacionales que son referencia en su campo y que siempre han tenido una palabra de amabilidad hacia esta tesis: Dr. Jérôme Prieto, Dr. Isaac Casanovas-Vilar, Dr. Jan van Dam, Dr. Kees Hordijk, Dra. Wilma Wessels, Dr. Hans de Bruijn, Dr. Marc Furió, Anne Blair Gould, Dra. Carmen Sesé, Dr. János Hír, Dr. Olivier Maridet, Dra. Gudrun Daxner-Höck, Dra. Anneke Bosma, Dra. Madelaine Böhme. Debo expresar mi gratitud hacia aquellas personas que me acogieron en las instituciones que visité durante estos años, Dr. Gertrud Rössner, Dr. Michael Rummel, Dr. Loïc Costeur, Dr. Reinhard Ziegler, Dra. Ursula Göhlich, así como agradecer los comentarios y ayuda prestada a los Dres. Paco Ruiz y Mathíjs Freudenthal. Especial cariño le tengo a los momentos que me dedicó el gran Dr. Albert van der Meulen. Su cercanía, su trato cariñoso, el entusiasmo que desprendía en aquellas conversaciones en Utrecht en las que enseñaba las maravillas que manaban de su mente y que hacía "por diversión" son impagables. No se puede explicar la sensación de estar aprendiendo y viviendo un momento crucial en tu vida, mientras te enseña gráficos y PCAs hechas en un momento de lucidez visionaria.

Durante los primeros años de esta investigación tuve la gran suerte de compartir grandes momentos con la Dra. Nieves López Martínez. Desafortunadamente, falleció en 2010 y esta tesis se adolece de ello, no habiendo podido comentar con ella los aspectos más interesantes. Nieves fue una gran impulsora de la paleontología a todos los niveles y su legado nos sigue llegando a través de su linaje.

Hay otra persona que, muy desgraciadamente, no pude conocer. El Dr. Remmert Daams es, para los que empezamos después de su marcha, un mito. Es alguien que sale en todas las conversaciones que tuve a lo largo de estos años. Bien porque el material que estudié lo encontró él, o bien porque sus notas manuscritas aparecen en artículos prestados, en etiquetas en salas de colecciones y en cuadernos de campo. Es una especie de guía omnipresente detrás de cada uno de los sacos que trié, cada uno de los dientes que miré y cada uno de los artículos que leí. Nunca se agradecerá lo suficiente el esfuerzo de tantos años sobre la lupa.

Y después de los autores, lo siguiente que se ve en una publicación es la afiliación. La primera

que debo poner es el Museo Nacional de Ciencias Naturales-CSIC, en concreto “la cochambre” y “eldes pachode Pablo”. Fueron los primeros sitios en los que disfruté de contratos en los que me inicié en la investigación y que aunque han pasado unos años de desvinculación contractual, no he sentido el desarraigo. Debo agradecer el apoyo científico y moral del personal que trabaja allí: sobre todo, agradezco al Dr. Jorge Morales su tiempo dedicado a disertar algunos temas de esta tesis, el apoyo que siempre tiene hacia los investigadores en formación y que siempre tenga tiempo para hablar de cualquier tema (siempre que haya sucedido en 1974). Agradezco a la Dra. Carmén Sesé la disponibilidad que siempre ha tenido, tanto para prestar el material o bibliografía como para charlar un rato acerca de estos fósiles. Asimismo, agradezco a la Dra. Maite Alberdi que accediera a formar parte de la propuesta del tribunal. Agradezco enormemente la ayuda prestada en el capítulo 10 por el Dr. Juan López-Cantalapiedra, que posee la Curiosidad (con mayúsculas) que se le supone a cualquier científico. Igualmente, el Dr. Israel Sánchez siempre estuvo dispuesto a echar una mano ya fuera en la filogenia o en cómo enviar un manuscrito, por todo, muchas gracias. También tengo que agradecer el interés demostrado por (el futuro Doctor) Oscar Sansisidro al que acudí para solucionar dudas y que adquirió tal compromiso que se involucró en la tarea al máximo nivel. Gracias, Oscar, por contribuir también a que las figuras del Capítulo 10 lucieran muchísimo más bonitas. Agradezco también los comentarios de la Dra. Raquel López-Antoñanzas y el tiempo que me prestó constantemente. Además de contar con la calidad profesional, el museo cuenta con un gran equipo de compañeros que ha sido un inmenso soporte todos estos años. Muchas gracias Adri, Vero, Gema, Blanca, Juan, Oscar, Enrique, Patricia, Susana, Israel, Vicky, María, Alberto, Dani y Cayetana.

La segunda afiliación que he tenido ha sido en el Departamento de Paleontología de la Universidad Complutense de Madrid. No fue un cambio traumático al ser el centro en el que estudié la licenciatura y que considero mi casa. Allí encontré excelentes científicos y mejores personas que me ayudaron a participar activamente en el departamento, a intentar aportar mi granito. Agradezco mucho la ayuda prestada y los consejos del Dr. Manuel Hernández Fernández, que siempre está dispuesto a escuchar y lo que es mejor, a aportar soluciones. También tengo que dar las gracias a la Dra. Laura Domingo que siempre dispuesta a oír cualquier tipo de idea y a dar su sincera opinión. Por esto (y muchas cosas más que pasan cuando trabajas “espalda con espalda”) Laura, es algo de lo que nunca estaré lo suficientemente agradecida. Gracias también a la Dra. Graciela Sarmiento, la Dra. Marisol Ureta y la Dra. Alejandra García-Frank de las que aprendí el significado del concepto “dedicación docente”, por tener esa extremada sensibilidad hacia los más jóvenes y dotar de sentido común a todas sus palabras. Agradezco al Dr. Sixto Fernández López sus palabras de amabilidad, espero haber cumplido sus expectativas. Igualmente tengo que agradecer al Dr. Diego García-Bellido y al Dr. Javier Élez el interés y la amabilidad siempre mostrada hacia este proyecto. Recuerdo con especial cariño sus consejos y palabras de ánimo durante el campamento de Geología Regional en Aguilar de Campoo, ya en los últimos años de

la licenciatura, que sirvieron para “encauzarme” hacia esta tesis doctoral. Agradezco también la ayuda presatada por D^a Charo Palomo en todos los temas administrativos. Muchas gracias a mis amigos de la UCM con los que podido compartir muchos momentos además de un despacho. Por todos los desvaríos que han soportado, risas y discusiones en los cafés autofinanciados, muchas gracias Ana Rosa, Javi, Laura, Sole, Isra, Alicia, Belén, Manuel, Xabi, Omid, Jonathan, Ángela, Daniel, Ismael, Ester, Juan Alberto, Fernando, Angélica y Ricardo. Muchas gracias a mis amigos geólogos, cada uno en una parte del mundo, pero siempre unidos: Elena, muchas gracias por ser una constante en mi vida, aun a 6000km y diez años después sigo riéndome con tus mismos chistes malos. Aunque lo que realmente agradezco de Elena es que me enseñara a hacer tablas dinámicas en Excel. Muchas gracias Kiko por mostrarte siempre tan entusiasta con esta tesis y alegrarte tanto como yo de que por fin esté sobre la mesa. Muchas gracias a Virginia “larubia”, Virginia “lamorena”, Rocío, Álvaro, Raquel y Mario. Pocas veces se sabe el momento exacto en el que te haces amigo de alguien y pero para mí fue en un coche camino a Oliva escuchando (a nuestro pesar) a Los Planetas, montando una tienda de campaña, tomando el sol y viendo el amanecer sobre el mar.

He tenido la suerte de disfrutar de tres estancias en el extranjero, dos en Naturalis Biodiversity Center en Leiden (Países Bajos) y una en el Bayerische Staatssammlung für Paläontologie und Geologie de Munich (Alemania), además, he disfrutado de una beca del proyecto Synthesys que me permitió visitar el Naturhistorisches Museum de Viena (Austria). Allí encuentran mi más profundo agradecimiento los profesionales que me hicieron un hueco y no dejaron que cayera en la melancolía de estar lejos de casa. Las estancias me sirvieron para centrar, recoger datos y fuerzas para seguir. Muchas gracias Lars, Anne, Helena, Stefan, Jérôme, Gertrud, Oliver y Ursula por ser mi familia en el exilio.

Al final de todo artículo, una vez leídos los autores, las afiliaciones y todo el contenido, hay un apartado de Agradecimientos, en los que se reconoce la labor de apoyo a todas aquellas personas que han contribuido a la mejora del artículo. Vayan estos meta-agradecimientos por delante.

A los revisores y consejeros: Irene, (dentro de poco haremos 30 años de amistad y no puedo entender la vida sin ella), Marie, Pilar, Sara y Chema, han sido grandes apoyos a la hora de ignorar un poco la Paleontología y mirar hacia el lado Perdido de la vida. Mirian, con su valor, fortaleza, inteligencia y sabiduría ha iluminado cada rincón oscuro de estos años. Además debo agradecer sus constantes consejos editoriales y lingüísticos que han contribuido a que la introducción sea más legible y han evitado el desastre de que alguien tan profano en Latín como yo pusiera nombre a una especie. Muchas gracias Benks y Miri por responder siempre que sí a la pregunta: ¿En serio quieres que te lo cuente? (el Miocénico es mío y el Micénico es tuyo). Sois la pieza única que me hace falta, vuestra amistad es una solución sólida en la que no se puede hacer una frontera clara entre lo que das y lo que recibes.

A los conservadores: mi Familia. Padres, Hermanas Marisa y Maribel, Hermanos Mario y José Manuel, y sobrinos Álvaro y Diana, sois los conservadores de mi cordura. Y a mi familia Castellana: Antonio, Angelines, Julián, Ester, Jorge, Marga, Diego y Manolo. Agradezco especialmente a Marga que haya dedicado un rato de su tiempo a adecentar el capítulo 11 con sus comentarios y revisiones.

Tanto los revisores como los conservadores han sido el corazón de esta tesis, cada vez que me recordaban que la paleontología no es lo importante y que todo lo grande qué es, no tiene sentido entre ellos. Que no les importa cuántos SCI tienes, les importa que no llegues tarde a comer, que pases más tiempo con tus hermanos, que les acompañes a Dinópolis o que te lo pases bien el día de su cumpleaños. Todavía no encuentro palabras para agradecerles que me hayan apoyado en todo esto, que me hicieran sentir que podía hacerlo, ni siquiera dudasen un segundo de mi capacidad, incluso que fingieran que les interesaba muchísimo lo que pasaba con los ratones de Mioceno, haciendo preguntas de corta-pega o dando cabezadas cada vez que les hablaba. Padres, no os podéis imaginar lo increíblemente agradecida que os estoy por darme una educación (tanto dentro como fuera de la academia), enseñarme a valorarla y aprovechar la vida con humildad y buen humor. Gracias por enseñarme que el camino a la felicidad no está dentro de la universidad, está junto a vosotros. Os quiero mucho a todos.

En los trabajos del equipo de micromamíferos al que pertenezco, siempre se hace una mención especial en los agradecimientos a todas las personas que han trabajado en las campañas de campo y triado del material, en resumen, al núcleo duro del trabajo. Ese núcleo duro que aguanta carros, carretas y trípodes por los campos de Daroca y Nombrevilla es Antonio Martín Monge. Toño quiso participar en esta tesis realizando las fotos de los yacimientos que aparecen en la introducción, sin darse cuenta de que años atrás ya se había “ganado los agradecimientos”, escuchando, empatizando, apoyando, dialogando y consolando. Por todas las noches de lloros, meses separados, encuentros y reencuentros en aeropuertos, conversaciones por Skype en el suelo del baño muniqué, se ha ganado no sólo unos agradecimientos, el cielo entero. Muchas gracias Toño.

A continuación en una publicación se ofrecería una lista de las referencias bibliográficas que se han usado, pero yo sólo tengo una:

López-Guerrero, P. Cricetodontini (Rodentia, Mammalia) del Mioceno medio y superior del área de Daroca (Aragón, España): sistemática y filogenia. 420 pp. Thesis Dissertation. Universidad Complutense de Madrid. Madrid.

.....PRESENTACIÓN

Las páginas que siguen a esta presentación tienen un tema central muy claro, el estudio de dos géneros extintos de cricétidos. A lo largo de este trabajo he intentado explicar de la mejor manera posible, qué pasó con *Cricetodon* e *Hispanomys* durante el Mioceno medio y superior. Si bien se centra principalmente en el estudio sistemático, también se ha acompañado esta parte con estudios en otros campos como son la bioestratigrafía, las relaciones filogenéticas y la paleobiogeografía. Una tarea que parecía fácil al principio, se empezó a volver complicada con los años y de nuevo, cuando echo un vistazo a las casi 400 páginas que siguen a este prefacio, parece simple.

Este volumen sigue la Normativa de Desarrollo del régimen relativo a la elaboración, tribunal, defensa y evaluación de la tesis doctoral del Real Decreto 1393/2007. Se compone de una introducción, un apartado de paleontología sistemática, otro de síntesis bioestratigráfica, un estudio filogenético y unas conclusiones generales. Se ha considerado necesario un capítulo individual para discutir endetalle tanto la primera presencia de la tribu en la cuenca, como la descripción de una especie nueva. Otros dos capítulos tratan al resto de especies de *Cricetodon* y de *Hispanomys*. Cada capítulo se ha estructurado a modo de artículo científico, de manera que cuenta con una introducción propia en la que se trata en concreto el tema planteado y unas conclusiones directamente relacionadas con el contenido del capítulo. Este tipo de estructura nos ha permitido avanzar en la publicación de los capítulos en revistas científicas y parte del contenido de esa tesis se encuentra en diferentes estadios de publicación. Las referencias bilbiográficas de cada uno de los artículos publicados se presentan al inicio de cada uno; aún así, a continuación se presenta una lista de los capítulos y artículos escritos:

El capítulo 5: *Revision of Cricetodon soriae (Rodentia, Mammalia), new data from the middle Aragonian (middle Miocene) of the Calatayud-Daroca basin (Zaragoza, Spain)*, ha sido publicado en la revista Journal of Vertebrate Paleontology. Aquí se hablará de la primera especie de *Cricetodontini* registrada en el área de estudio.

El capítulo 6: *Cricetodon from the middle and late Aragonian of the Calatayud-Daroca Basin (Spain): a taxonomical description*, se ocupa de otras dos especies de *Cricetodon* y ha sido publicado

junto con el capítulo 9: *Cricetodontini from the Calatayud-Daroca Basin (Spain): an up-to-date of their stratigraphical distribution*, en el que se analiza el valor bioestratigráfico de las especies estudiadas. Ambos capítulos forman parte de un mismo artículo sintético publicado en la revista Comptes Rendus Palevol.

El capítulo 7: *New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution* describe una nueva especie registrada en el Aragoniense superior y está publicado en la revista Bulletin of Geosciences.

El capítulo 8: *Species of Hispanomys of the late Aragonian and early Vallesian (middle-late Miocene) of the Calatayud-Daroca Basin, Zaragoza, Spain* se ocupa de todas las especies de *Hispanomys* del área estudiada. La presencia de dos de ellas en uno de los yacimientos estudiados se ha publicado en el volumen del VI Encuentro de Jóvenes Investigadores en Paleontología, en la revista Palaeontologica Nova, Publicaciones del Seminario de Paleontología de Zaragoza (SEPAZ). Ese estudio, realizado en el año 2008, ha sido reescrito en su mayor parte para la presente tesis doctoral. Las descripciones y las discusiones han sido ampliadas, y se han descrito fósiles de nuevos yacimientos, por lo que el capítulo 8 no es una transcripción de esa publicación. Aún así hay que reconocer que fue el germen del mismo.

El capítulo 10: *First phylogenetic approach to Cricetodontini (Rodentia, Mammalia) from the Miocene and Pliocene of Europe and Asia Minor* es un capítulo que se encarga de evaluar las relaciones filogenéticas y está proyectado para ser enviado en los próximos meses.

Cricetodontini (Rodentia, Mammalia) from the middle and upper Miocene in the Daroca area (Aragón, Spain): Systematics & Phylogeny.

Introduction

This dissertation presents the systematic study of two fossil genera of the family Cricetidae (Rodentia, Mammalia). The order Rodentia is one of the most diverse and abundant extant groups of mammals and its origin is dated 57 Ma. Currently, it counts with more than 2,200 species worldwide distributed. The studied fossils belong to the tribe Cricetodontini which was very common in Europe during the Miocene. Its representatives stand out among other Miocene cricetids for their large size. The taxonomy, as well as the other micromammal groups, is mainly based on the dental characters.

The first occurrence of the tribe is located in the early Miocene of Asia Minor. Its members remained in Anatolia during most part of the early Miocene and they occurred in Europe during the latest early Miocene. Then, the tribe was widely extended in Europe being the middle Miocene (middle and late Aragonian) a very interesting interval. At this time, the diversity of the tribe increases, including species with more complex morphology and higher intraspecific variability than the older representatives. Their decline starts in the late Miocene (early Vallesian). The number of species decrease, until their extinction at the beginning of the Pliocene. On the light of this evolutionary history, the most interesting interval to study the Cricetodontini is the middle to late Miocene (middle and late Aragonian, and early Vallesian).

The taxonomy of the Cricetodontini is controversial and some aspects remain unresolved. For example, the generic ascription of some species were debated and the inclusion of some genera in the tribe is still dubious. Because of this, the species of Cricetodontini have not been recommended for the proposal of detailed biostratigraphical scales. Only the genera of Criceto-

dontini were used in biostratigraphical studies and they defined large time intervals and major continental units. However, recent studies in Cricetodontini from Central Europe and Anatolia demonstrate that their species have served as a source to establish biostratigraphical subdivisions. Thus, a better knowledge of the taxonomy of the Cricetodontini from the Iberian Peninsula could reveal them as good biostratigraphical indicators as well.

Apart from that, the establishment of the phylogenetic relationships of Cricetodontini are also complicated due to the mosaic morphology displayed by some of their representatives. Moreover, the phylogenetic hypotheses proposed before are outdated and therefore, it is necessary to make a phylogenetic study of the Cricetodontini.

The Calatayud-Daroca Basin (Zaragoza, Spain) offers the best context for studying the Cricetodontini since many fossil sites are recovered within this time range. In addition, most of the fossiliferous levels found in the Calatayud-Daroca Basin are stratigraphically superposed which allow to assess a numerical dating by means of paleomagnetostratigraphy. Fifty-four localities from the Calatayud-Daroca Basin provide fossils of Cricetodontini from the middle and late Aragonian, and early Vallesian. Their age estimation ranges from 14.06 to 9.98 Ma. Due to the high number of localities and their suitability for being numerically dated, the Calatayud-Daroca Basin is one of the most studied basins in the Iberian Peninsula.

Synthesis

- Aims

The main aims of this study are: to characterize, morphologically and metrically, the representatives of Cricetodontini in the Calatayud-Daroca Basin in order to resolve their taxonomy. Another aim is to specify their stratigraphical distributions during the middle and late Miocene (middle and late Aragonian, and early Vallesian) and evaluate their biostratigraphical use. And finally, the last aim is to assess the phylogenetic relationships among Cricetodontini from Europe and Asia by means of a cladistic analysis.

- Results in taxonomy and phylogeny

The whole sample includes 3,181 teeth that have been morphometrically studied. These fossils are distributed among eight species. Four of them belong to *Cricetodon* and four to *Hispanomys*. A new species, *Cricetodon nievei*, was described due to the unique combination of characters that the specimens from sites Toril 2, 3A, 3B and Las Planas 5H present.

Cricetodon soriae López Martínez et al., in Hernández Fernández et al., 2006 is earliest occurrence of the tribe in the basin and also, the oldest representative of the tribe on the Iberian Peninsula. Its stratigraphical distribution covers the local biozone E (MN5; 14.06-13.80 Ma)

during the middle Aragonian. The results of the cladistic analysis indicate that *Cricetodon soriae* is closely related to other species from southwestern Europe such as *C. sansaniensis* Lartet, 1851, *C. jotae* Mein and Freudenthal, 1971a and *C. meini* Freudenthal, 1963.

Cricetodon sansaniensis is present in five localities covering the local biozone F (MN6 *pro parte*; 13.76–13.68 Ma) for the late Aragonian. *Cricetodon sansaniensis* were previously related with *C. aureus* Mein and Freudenthal, 1971b and *C. albanensis* Mein and Freudenthal, 1971a. But, in present study, *C. sansaniensis* is related to *C. soriae*, *C. jotae* and *C. meini*.

Cricetodon jotae is present in 13 sites belonging to local biozones, G1 and G2, (MN6 pp.) and G3 (MN7/8 pp.) in the late Aragonian. The first occurrence of *C. jotae* is in Las Planas 5B (local biozone G1; 13.56 Ma). The youngest occurrence of *C. jotae* is in Las Planas 5K, (local biozone G2; 13.08 Ma). However, *C. jotae* is present in younger localities from the local biozone G3 with no numerical age. *Cricetodon jotae* was related to *C. meini*. Both are sister species in the results of the phylogenetic analysis supporting this relationship.

The new species, *Cricetodon nievei*, is found in four sites, all of them belong to the G3 biozone (MN7/8 pp.; 12.60–12.65 Ma) from the late Aragonian. *Cricetodon nievei* shares more features with the species from the Central Europe basins than with those from the Calatayud-Daroca Basin. The results of the cladistic analysis agrees with this; *Cricetodon nievei* is grouped with *C. bolligeri* Rummel, 1995 which is endemic from Central Europe.

Hispanomys aguirrei (Sesé Benito, 1977 in López Martínez et al., 1977) is registered in seven localities from the local biozone G3 (MN7/8 pp.; 12.39–11.33 Ma) from the late Aragonian. The results of the cladistic analysis report that *Hispanomys aguirrei* is situated among the other species of *Hispanomys* indicating that its assignation to *Hispanomys* is correct. It is sister species to *H. lavocati* (Freudenthal, 1966).

Hispanomys lavocati is recovered from three sites placed in the local biozone G3 (MN7/8 pp.; 11.87–11.88 Ma) for the late Aragonian. *Hispanomys lavocati* is situated in the strict consensus as a member of *Hispanomys* closely related with the other southwestern Europe *Hispanomys* from the late Aragonian.

Hispanomys nombrevillae (Freudenthal, 1966) is registered in seven localities from the local biozone H (MN7/8 pp. and MN9 pp.; 11.20 Ma–10.77 Ma) from the late Aragonian and the early Vallesian. *Hispanomys nombrevillae* was related with *H. bijugatus* (Mein and Freudenthal, 1971a) and also considered as the descendant of *H. aguirrei*. In the cladistic analysis, *H. nombrevillae* is sister species to the derived *Hispanomys* revealing that *H. nombrevillae* is more closely related to *H. aragonensis* (Freudenthal, 1966) than to *H. aguirrei*/*H. bijugatus*.

Hispanomys aragonensis is present in seven localities from the local biozone I (MN9 pp.;

10.72–9.98 Ma) for the early Vallesian. It was related to *H. decedens* (Schaub, 1925). In the cladogram, these two species are situated in separate clades, precluding the relationship between them. Apart from this, *H. aragonensis* is the most basal taxon to the clade that includes *H. peralensis* Van de Weerd, 1976 supporting this relationship previously stated.

Conclusions

In general, the morphological patterns observed in the Cricetodontini from the Calatayud-Daroca Basin fit the following model: basal morphology during the older record in the middle, late Aragonian (*C. soriae*, *C. sansaniensis*); mosaic morphology in most of the species from the late Aragonian (*C. jotae*, *C. nievei*, *H. aguirrei* and *H. lavocati*), and finally, derived forms as *H. nombrevillae* and *H. aragonensis* in the early Vallesian. This evolutionary history is also recognized in other European faunas Cricetodontini but not synchronously.

The tribe Cricetodontini is present in local biozones from E to I, its stratigraphical distribution spans an age range from 14.06 to 9.98 Ma. The stratigraphical distributions of the studied species, generally, fit well with the local biozonation; those of *Cricetodon soriae*, *C. sansaniensis*, *H. nombrevillae* and *H. aragonensis* correlate with local biozones E, F, H and I, respectively. The distribution of *C. jotae* is long and it ranges about 1 my covering several local biozones (G1, G2 and partly G3). Four species of Cricetodontini (*C. jotae*, *C. nievei*, *H. aguirrei* and *H. lavocati*) are present in local biozone G3 (late Aragonian). On the light of the distribution of this species and the unusually long duration of the G3 (1 my) the redefinition and subdivision of this biozone is recommended. Nevertheless, we suggest a study of the complete rodent fauna before proposing a new biozonation.

Three main clades can be observed in our cladistic analysis. The cladogram reveals that *Deperetomys* and *Byzantinia* are monophyletic, *Cricetodon* is paraphyletic and the species of *Hispanomys* together with *Ruscinomys* Freudenthal, 1966 form a clade.

The centre of origin of the studied Cricetodontini is Anatolia, where the tribe was present since the early Miocene. *Deperetomys* extends its range in Asia Minor and Central Europe, *Cricetodon* is wide spread across Europe and Asia Minor, whereas *Byzantinia* is strongly linked with Asia Minor. *Hispanomys*+*Ruscinomys* are linked with south-western Europe.

Cricetodon soriae, *C. sansaniensis* and *C. jotae* are grouped together in a clade with *C. meini*. *Cricetodon nievei* is clustered with the species from Central Europe, clearly separated from the other Cricetodontini from the Calatayud-Daroca Basin. The species of *Hispanomys* are split in two clades: *H. aguirrei* and *H. lavocati* are basal, and *H. nombrevillae* and *H. aragonensis* are derived.

Present study of the tribe Cricetodontini in Calatayud-Daroca Basin contributes to evidence

their importance in mammalian paleontology. We have recognized their suitability in biostratigraphical studies and resolved some important systematic and phylogenetic questions. However, in order to confirm some of the phylogenetic hypotheses made and also to better understand the evolutionary history of the tribe, it is recommended to widen the study to other rodent taxa and add supplementary morphological data.

Thus, future research should focus on the study of other genera of rodents from the late Aragonian and early Vallesian from Calatayud-Daroca Basin to redefine the local biozonation. In addition it would be interesting to extend the phylogenetic study to other genera from Asia Minor and Southwestern Europe in order to resolve some phylogenetic issues, especially those concerning *Deperetomys*, *Cricetodon* and *Hispanomys*.

Cricetodontini (Rodentia, Mammalia) del Mioceno medio y superior del área de Daroca
(Aragón, España): Sistemática y filogenia

Introducción

Esta tesis doctoral presenta el estudio sistemático y bioestratigráfico de dos géneros extintos de la familia Cricetidae (Rodentia, Mammalia). El orden Rodentia tiene sus orígenes hace 57 Ma y sus representantes llegan hasta nuestros días. Es el grupo de mamíferos más diverso que existe en la actualidad y sus especies están distribuidas por toda la superficie terrestre, excepto en el continente antártico. Los géneros estudiados en la presente tesis doctoral se clasifican en la subfamilia Cricetodontinae y en la tribu Cricetodontini. Esta tribu está presente desde principios del Mioceno hasta el Plioceno de Europa. Forman parte de las comunidades de roedores de forma continua durante el Mioceno medio y superior. El primer registro se encuentra en el Mioceno inferior de Asia Menor. Los representantes de la tribu se mantienen dentro de Anatolia durante la mayor parte del Mioceno inferior y se registran en Europa a finales del Mioceno inferior. A partir de ese momento las especies de Cricetodontini se distribuyen ampliamente por toda Europa y Asia. En la segunda mitad del Mioceno medio la diversidad de la tribu aumenta, incluyendo especies con morfología más compleja y con una mayor variabilidad intraespecífica. Su declive en Europa y Asia Menor comienza a principios del Mioceno superior; su distribución geográfica empieza a restringirse al suroeste de Europa, encontrando el último representante de la tribu en Francia. A la luz de su historia evolutiva, parece que el intervalo más interesante para el estudio de los Cricetodontini es el Mioceno medio y superior (Aragoniense medio y superior, y Vallesiense inferior)

Este grupo tiene una alta variabilidad morfológica que dificulta la asignación genérica de algunas especies. Así mismo, la inclusión de algún género en la tribu (p.e. *Deperetomys*) ha sido

igualmente discutida. Por otro lado, algunas especies presentan una combinación de caracteres basales y derivados que se ajusta a un modelo evolutivo en mosaico. Esto ha llevado a recomendar que se evite el uso de las especies de Cricetodontini para realizar escalas bioestratigráficas locales y su uso bioestratigráfico para el Mioceno ibérico se limitó a caracterizar solamente grandes intervalos de tiempo. Sin embargo, en otras cuencas europeas y de Asia Menor, se está poniendo de manifiesto la utilidad bioestratigráfica de las especies de Cricetodontini encontradas allí.

Existe una gran dificultad para estimar las relaciones filogenéticas entre las especies de Cricetodontini debido al patrón morfológico en mosaico descrito anteriormente. Además, las hipótesis filogenéticas propuestas hasta la fecha se han quedado desactualizadas al haberse descrito muchas especies posteriormente. Por eso es necesario acompañar al estudio taxonómico de un estudio filogenético de los representantes de Cricetodontini.

La Cuenca de Calatayud-Daroca (Zaragoza, España) ofrece las condiciones óptimas ya que presenta uno de los registros de micromamíferos fósiles mejor documentados y estudiados del Mioceno. Además, muchos de los yacimientos están en superposición estratigráfica lo que ha permitido datar numéricamente a la gran mayoría. Existen 54 yacimientos en el que se tiene registro de Cricetodontini durante el Aragoniense medio y superior y Vallesiense inferior. Su edad estimada se sitúa entre los 14,06 y los 9,98 Ma. Debido al gran número de yacimientos y su idoneidad para ser datados, la Cuenca de Calatayud-Daroca es una de las más estudiadas en la Península Ibérica.

Síntesis

- Objetivos

Los principales objetivos de esta tesis doctoral son: caracterizar, morfológica y métricamente, las especies de Cricetodontini presentes en la Cuenca de Calatayud-Daroca y así resolver su taxonomía. Con esta información taxonómica se estudiarán sus distribuciones estratigráficas y se evaluará la utilidad bioestratigráfica de los Cricetodontini dentro de la cuenca. Finalmente, se discutirán las relaciones filogenéticas de cada una de las especies encontradas y las del resto de Europa y Asia Menor a través de la metodología cladística.

- Resultados de la taxonomía y la filogenia

El estudio taxonómico resuelve que los 3.181 fósiles se reparten entre ocho especies pertenecientes a dos géneros de la tribu. Cuatro son especies del género *Cricetodon* y cuatro de *Hispanomys*. Se ha descrito una nueva especie, *Cricetodon nievei*, debido a la combinación novedosa y única de las características morfológicas y métricas que posee.

Cricetodon soriae es la especie más antigua de la tribu Cricetodontini en la cuenca y, además,

es el miembro más antiguo de la tribu en la Península Ibérica. Está presente en la biozona E, entre los 14,06 Ma y los 13,80 Ma durante el Aragoniense medio. *Cricetodon soriae* se relaciona estrechamente con otras especies de Europa occidental, como son *C. sansaniensis*, *C. jotae* y *C. meini*.

Cricetodon sansaniensis está presente en cinco localidades que cubren la biozona local F y abarcan un intervalo temporal de 13,76 a 13,68 Ma. *Cricetodon sansaniensis* no parece tener relación con ninguna de las especies con las que se había relacionado anteriormente y se rechaza su relación directa con *C. aureus*. En cambio, sí parece tener una relación estrecha con *C. soriae*.

Cricetodon jotae se ha encontrado en 13 localidades situadas en las biozonas locales, G1 y G2 (MN6 *pro parte*), así como en la parte inferior de la G3 (MN7/8 pp.) en el Aragoniense superior. La primera presencia de *C. jotae* se produce en Las Planas 5B (biozona local G1), datada en 13,56 Ma. La última localidad datada numéricamente con *C. jotae* es Las Planas 5K (13,08 Ma; biozona local G2). Sin embargo, *Cricetodon jotae* está presente en las localidades más recientes pertenecientes a biozona local G3, de las cuales no se dispone de edad numérica. *Cricetodon jotae* es especie hermana de *C. meini*, lo que confirma la relación propuesta por los autores.

La nueva especie *Cricetodon nievei* se ha encontrado en cuatro yacimientos de la biozona local G3 (12,65-12,60 Ma). Se caracteriza por presentar una combinación de caracteres primitivos y derivados. *Cricetodon nievei* es más parecido a las especies de *Cricetodon* e *Hispanomys* de la Cuenca de Vallès-Penedès, Francia y Europa central que a las especies registradas en Calatayud-Daroca. En el análisis filogenético, *Cricetodon nievei* se agrupa con las especies procedentes de Europa central, claramente separado de los otros Cricetodontini de la Cuenca de Calatayud-Daroca.

Hispanomys aguirrei se registra en siete localidades de la biozona local G3 (MN7/8 pp.; 12,39-11,33 Ma) del Aragoniense superior. *Hispanomys aguirrei* se asigna a *Hispanomys*, ya que se encuentra entre las otras especies de este género. Parece confirmarse que *H. aguirrei* se relaciona con *H. lavocati*, su especie hermana.

Hispanomys lavocati se ha incluido en el género *Hispanomys* a pesar de las propuestas de otros autores, que lo incluyen en *Cricetodon*. *Hispanomys lavocati* es una de las especies más grandes de *Hispanomys* descritas y está presente en tres yacimientos de la biozona local G3 (MN7/8 pp.; 11,87-11,88 Ma) del Aragoniense superior. *Hispanomys lavocati* parece tener una relación directa con las otras especies de *Hispanomys* del suroeste de Europa.

Hispanomys nombrevillae está presente en siete localidades de la biozona local H (MN7/8 pp. y MN9 pp.; 11,20 Ma-10,77 Ma) del Aragoniense superior y Vallesiense inferior. *Hispanomys nombrevillae* aparece como una especie basal al clado que contiene todas las especies derivadas de *Hispanomys*, estado más estrechamente relacionada con *H. aragonensis* que con *H. aguirrei* o *H.*

bijugatus.

Hispanomys aragonensis se encuentra en siete localidades de la biozona local I (MN9 pp.; 10,72-9,98 Ma) del Vallesiense inferior. *Hispanomys aragonensis* se encuentra entre las especies de *Hispanomys* más derivadas, el análisis rechaza su relación directa con *H. decedens*. Se relaciona directamente con *H. peralensis* tal y como se había discutido anteriormente.

Conclusiones

La tribu Cricetodontini está presente en las biozonas locales desde la E hasta la I, con un rango de edad estimada de 14,06 a 9,98 Ma. Las distribuciones estratigráficas de las especies estudiadas, en general, se correlacionan con la biozonación local, por lo que se puede indicar que se trata de buenos indicadores bioestratigráficos. Las distribuciones de *C. soriae*, *C. sansaniensis*, *H. nombrevillae* e *H. aragonensis* coinciden con las biozonas locales E, F, I y H, respectivamente. Sin embargo, la distribución de *Cricetodon jotae* es muy amplia y oscila alrededor de un 1 Ma cubriendo varias biozonas locales (G1, G2 y G3). Por su parte, *C. nievei* se encuentra únicamente en la parte media de la G3. A la luz de la distribución de las especies y de lo inusualmente larga que es la biozona local G3 (1 Ma) se recomienda que ésta se revise y se redefine o subdivida. Sin embargo, también se sugiere que se estudie todo el conjunto de las especies de roedores antes de proponer una nueva biozonación.

Las especies estudiadas en el análisis filogenético se dividen en tres clados principales que corresponden a *Deperetomys*, *Byzantinia* e *Hispanomys*+*Ruscinomys*. El cladograma indica que *Deperetomys* y *Byzantinia* son monofiléticos, *Cricetodon* es parafilético y que las especies de *Hispanomys* forman un clado junto con las de *Ruscinomys*.

El centro de origen de Cricetodontini es claramente Asia Menor. También que *Byzantinia* e *Hispanomys* están estrechamente vinculados con un área geográfica específica: Asia Menor y el suroeste de Europa respectivamente. Por otra parte, *Deperetomys* se extiende por Asia Menor y Europa Central, y *Cricetodon* está ampliamente distribuido por toda Europa y Asia Menor. También se han detectado tres momentos migratorios (hace 18 Ma, 15 Ma y 14 Ma respectivamente) que tienen su origen en Anatolia y en los que participan diferentes especies que dan lugar a las encontradas en la Cuenca de Calatayud-Daroca.

Cricetodon soriae, *C. sansaniensis* y *C. jotae* están agrupados en un clado junto con *C. meini*. *Cricetodon nievei* se sitúa con las especies de Europa central, claramente separadas de las especies de la Península Ibérica. Por su parte, las especies de *Hispanomys* se separan en dos clados: *H. aguirrei* e *H. lavocati* en uno más basal que el que contiene a *H. nombrevillae* e *H. aragonensis*.

Este estudio de la tribu Cricetodontini de Calatayud-Daroca contribuye a evidenciar su

importancia en la paleontología de mamíferos. Hemos reconocido su aptitud para la bioestratigrafía y se han resuelto algunos de sus problemas sistemáticos y filogenéticos. Sin embargo, para confirmar algunas de las hipótesis aquí planteadas, es recomendable ampliar el estudio a otros grupos de roedores así como añadir más datos morfológicos a la matriz filogenética.

Así, las siguientes investigaciones encargadas de redefinir la biozonación local deberían enfocarse en el estudio de otros roedores de la Cuenca de Calatayud-Daroca del Aragoniense superior y del Vallesiense inferior. Además, sería muy interesante extender el estudio filogenético a otros géneros de Asia Menor y del suroeste de Europa.

I. Introducción

Capítulo 1

Introducción



INTRODUCCIÓN

“—¿Pero qué es lo que se supone que voy a ver?!
 (...) —No lo sé. En cierto sentido, depende de ti.”
 Stanislaw Lem, *Solaris*.

INTRODUCCIÓN A LOS ROEDORES

1

El origen del orden Rodentia (Mammalia) se ha datado en 57 Ma (Hartenberger, 1998; Montgelard et al., 2002) y sus representantes llegan hasta nuestros días. Es el grupo de mamíferos más diverso que existe en la actualidad. Cuenta con más de 2.200 especies (Wilson y Reeder, 2005) distribuidas por toda la superficie terrestre, excepto en el continente antártico (Figura 1.1). En la presente tesis doctoral se estudian dos géneros de cricétidos extintos característicos del Mioceno. Los cricétidos, cuyos representantes son conocidos con el nombre común de hámsters, pertenecen a la superfamilia Muroidea. Existen diferentes opiniones respecto a su clasificación dentro de Muroidea y su relación con otro grupo de roedores, los múridos. Algunos autores (e.g., Hershkovitz, 1962; McKenna y Bell, 1997; Michaux et al., 2001; Musser y Carleton, 2005) consideran que los cricétidos y los múridos pertenecen a la misma familia: Muridae; otros (e.g., Lavocat, 1978; Reig, 1984; Flynn et al., 1985; Hartenberger, 1998; Freudenthal y Martín Suárez, 1999; Rummel, 1999; Maridet y Ni, 2013; Flynn et al., 2014), separan los miembros en dos familias diferentes: Cricetidae y Muridae. Aunque sigue sin haber consenso respecto a esta cuestión (e.g., Gomes Rodrigues et al., 2009), en esta tesis doctoral se va a seguir la segunda opción, en la que los cricétidos se incluyen dentro de la familia Cricetidae. Finalmente, siguiendo la clasificación de Mein y Freudenthal (1971a) que incluye representantes fósiles, los géneros estudiados en la presente tesis doctoral se clasifican en la subfamilia Cricetodontinae y en la tribu Cricetodontini.

CRICÉTIDOS

Características principales de la familia Cricetidae

En la actualidad, Cricetidae es una de las familias de mamíferos más numerosas, con 681 especies repartidas en 130 géneros y 6 subfamilias (Musser y Carleton, 2005). Los cricétidos ocupan un amplio espectro de hábitats de climas tanto secos y húmedos, como fríos y calientes (Nowak, 1999). La morfología externa de los cricétidos es muy particular: tienen cuerpos pequeños, alargados, con colas cortas, ojos pequeños y orejas que están casi completamente ocultas en la piel (Figura 1.2). Su comportamiento es muy diverso, la mayoría son nocturnos y entran en letargo durante los periodos fríos para reducir su consumo energético. Aunque son bastante sedentarios, algunas especies pasan por ciclos de grandes poblaciones que pueden dispersarse a grandes distancias en periodos de abundancia (Nowak, 1999). En lo que respecta al modo de alimentación, los cricétidos son igualmente diversos; pueden ser carnívoros, omnívoros o herbívoros (Nowak, 1999). La fórmula dental es, generalmente, $1/1, 0/0, 0/0, 3/3=16$.




FIGURA 1.1. Distribución de las familias de roedores actuales y recientemente extintas (basado en Wilson y Reeder, 2005).

TRIBU CRICETODONTINI

La tribu Cricetodontini Simpson, 1945, forma un distintivo grupo de cricétidos de gran tamaño que se encuentran presentes desde principios del Mioceno hasta el Plioceno de Europa, Asia y África. Esta tribu incluye los géneros *Cricetodon*, *Hispanomys*, *Byzantinia*, *Ruscinomys*, *Deperetomys*, *Zramys*, *Eumyarion* y *Metamys* (Mein y Freudenthal, 1971a; Rummel, 1999). Las características más importantes de la tribu son, aparte de su tamaño, el grosor del esmalte, que puede presentar rugosidades; los surcos de los incisivos superiores y la tendencia de los molares a aumentar la hipsodoncia, a simplificar la morfología y a reducir el tamaño del M3 superior, así como a presentar el anterocono dividido (Rummel, 1999). La presencia de ectolofos muy variables en morfología y longitud dentro de cada una de las especies es también característica. En el capítulo 4 (Material y Métodos) se explicará con detalle la morfología de los ectolofos. Como se discute a lo largo del apartado III (Paleontología Sistemática), tienen una gran variabilidad interespecífica y por eso son caracteres diagnósticos en varias especies de Cricetodontini (Mein y Freudenthal,



FIGURA 1.2. Ejemplos de algunos roedores cricétidos actuales. (a-c). *Mesocricetus auratus* (d). *Onychomys* sp.

1971b; Agustí, 1982; Rummel, 1999). En general, se ha observado que los ectolofos siguen una tendencia al aumento de longitud y número de crestas que los componen (Agustí, 1982).

Durante el Mioceno medio y superior, los Cricetodontini forman parte de las comunidades de roedores de forma continua (De Bruijn y Ünay, 1996; Rummel, 1999; Kálin y Kempf, 2009; Daams et al., 1999c; López-Guerrero et al., en prensa). El primer registro de la tribu procede de la localidad de Inonak (Mioceno inferior, 22,564-22,268 Ma; De Bruijn et al., 1993) en Asia Menor. Los representantes de la tribu se mantienen dentro de Anatolia durante la mayor parte del Mioceno inferior y, tras un periodo sin registro, se encuentran en Europa en la isla de Aliveri (Grecia) a finales del Mioceno inferior (17,20-16,40 Ma; Hilgen et al., 2012). A partir de ese momento las especies de Cricetodontini se distribuyen ampliamente por toda Europa y Asia (De Bruijn et al., 1993; De Bruijn y Ünay, 1996; Rummel, 1999; Wu et al., 2009; Qiu, 2010) y su presencia es muy común durante la primera mitad Mioceno medio. En la segunda mitad del Mioceno medio se registra un gran número de especies nuevas de Cricetodontini en toda Europa (De Bruijn et al., 1993; Rummel, 1999; Kálin y Kempf, 2009; Prieto et al., 2010; Álvarez-Sierra et al., 2013), la mayoría de ellas son endémicas de las cuencas sedimentarias en las que se encuentran. Su declive en Europa y Asia Menor comienza a principios del Mioceno superior; su distribución geográfica empieza a restringirse al suroeste de Europa, encontrando el último representante de la tribu en Francia (De Bruijn et al., 1993; De Bruijn and Ünay, 1996; Rummel, 1999; López-Guerrero et al., 2011a; Álvarez-Sierra et al., 2013).

En la Península Ibérica, las especies de Cricetodontini siguen la misma historia evolutiva señalada anteriormente. En el Aragoniense medio (MN5) se registra por primera vez este grupo, se trata de una especie endémica de la Península Ibérica (De Bruijn et al., 1993; Hernández Fernández et al., 2006; López-Guerrero et al., 2007, 2013a; Hernández-Ballarín et al., 2010). Al comienzo del Aragoniense superior (MN6) se encuentran especies que se distribuyen ampliamente por toda Europa (Bolliger, 1994; Rummel, 2000, 2001; Rummel y Kálin, 2003; López-Guerrero et al., en prensa). La segunda mitad del Aragoniense superior (MN7/8) es un periodo muy interesante para este grupo. Se produce un aumento de la diversidad de Cricetodontini, hay un mayor número de especies que presentan alta variabilidad morfológica. Además, las especies que aparecen son endémicas (Álvarez-Sierra et al., 2003; 2013; López-Guerrero et al., 2014). Este aumento en el número de taxones de Cricetodontini durante el Aragoniense superior se refleja en el incremento del número de especies de *Cricetodon*, así como por la primera presencia de varias especies de *Hispanomys* (López-Guerrero et al., 2008, 2014, en prensa). Estas especies de *Hispanomys* poseen una amplia variabilidad morfológica que, en ocasiones, ha dificultado o puesto en duda su asignación genérica, como se verá en el capítulo 8. El Vallesiense inferior se caracteriza porque las especies de Cricetodontini presentan menor variabilidad morfológica y hay también un descenso en el número de especies (Álvarez-Sierra, 1983; Van de Weerd, 1976; Van Dam, 1997; Agustí, 1980; López-Guerrero et al., en prensa). Esta pérdida de biodiversidad se acusa durante el

Vallesiense superior que sólo cuenta con 3 especies de *Hispanomys* en todo el registro ibérico (Van Dam et al., 2001; Hartenberger, 1965; Casanovas-Vilar, 2007; López-Antoñanzas et al., 2010).

La historia evolutiva de los Cricetodontini tuvo lugar en un contexto de cambios climáticos importantes. A comienzos del Mioceno medio se ha detectado un intervalo temporal caracterizado por altas temperaturas a nivel global, con desarrollo de climas húmedos tropicales y subtropicales (Zachos et al., 2001; Domingo et al., 2012; Hérán et al., 2010). Este evento es conocido como el Óptimo Climático del Mioceno (cuyas siglas en inglés son MCO) y ha sido datado en ~17 a 14 Ma (Zachos et al., 2001). El MCO ha sido detectado tanto en sedimentos marinos (Abels, 2005; Mourik, 2011) como continentales (Domingo et al., 2012; Hérán et al., 2010). El MCO fue seguido por una caída relativamente rápida de las temperaturas, el Enfriamiento del Mioceno Medio o MMC (según sus siglas en inglés) que inicia el régimen climático frío del Neógeno (Lear et al., 2000; Zachos et al., 2001; Shevenell et al., 2004). Como se verá en los capítulos 5 y 9, el MMC es casi simultáneo a la primera presencia de *Cricetodon* en la Península Ibérica. Shevenell et al. (2004) describieron la Transición Climática del Mioceno Medio (MMCT en inglés) que tuvo lugar durante el Mioceno medio (14,20 a 13,80 Ma), relacionando el patrón de las temperaturas calculadas con fluctuaciones en la capa de hielo antártico. Finalmente, durante el Mioceno superior, se produce un enfriamiento adicional y una expansión de la capa de hielo de la Antártida (Kenneth y Barker, 1990) y del Ártico (Thiede y Vorren, 1994) que coinciden con el declive de la tribu Cricetodontin hasta su extinción en el Plioceno inferior.

Dada la historia evolutiva de los Cricetodontini y el contexto paleoclimático en la que ésta se desarrolla, parece que uno de los intervalos más interesantes para un estudio en profundidad de este grupo es la primera presencia en el Aragoniense medio (Mioceno medio). Igualmente interesante es estudiar esta tribu durante su amplia distribución por toda Europa y el rápido aumento de biodiversidad del Aragoniense superior (Mioceno medio). Finalizando con el estudio del inicio de su declive en el Vallesiense inferior (Mioceno superior).

La taxonomía de los Cricetodontini es complicada. Este grupo tiene una alta variabilidad morfológica por lo que, en ocasiones, que la asignación genérica de alguna especie es muy difícil (Prieto et al., 2010; López-Guerrero et al., 2014) y que la inclusión de algún género en la tribu (p.e. *Deperetomys*) fuera discutida (Maridet y Ni, 2013). Algunas especies presentan una combinación de caracteres basales y derivados que se ajusta a un modelo evolutivo en mosaico. Este tipo de modelo evolutivo ha llevado a algunos autores (Klein Hofmeijer y De Bruijn, 1988; De Bruijn et al., 1993) a recomendar que se evite el uso de las especies de Cricetodontini para realizar escalas bioestratigráficas, afirmando que su utilidad está sobrestimada. Ésta pudo ser una de las causas por la que el uso bioestratigráfico de los miembros de Cricetodontini para el Mioceno ibérico, se limitó a caracterizar grandes intervalos de tiempo (Daams et al., 1999c). Por ejemplo, se indicó que *Cricetodon* es un género que caracteriza el Aragoniense e *Hispanomys* el Vallesiense (Daams et al.,

1999c). Sin embargo, en otras cuencas europeas y de Asia Menor, se está poniendo de manifiesto la utilidad bioestratigráfica de las especies de Cricetodontini encontradas allí (Ünay et al., 2003; Kálin y Kempf, 2009; Böhme et al., 2012; De Bruijn et al., 2013), lo que hace muy interesante un estudio taxonómico en profundidad de las especies de Cricetodontini de la Península Ibérica que permita hacer una evaluación de la utilidad bioestratigráfica de las mismas similar a la realizada en centroeuropa.

Muchos de los estudios recientes sobre Cricetodontini señalan la dificultad de inferir las relaciones filogenéticas entre las especies debido al patrón morfológico en mosaico descrito anteriormente (Prieto et al., 2010, 2014; López-Guerrero et al., 2014, en prensa). Además, las hipótesis filogenéticas propuestas hasta la fecha (Mein y Freudenthal, 1971a; Agustí, 1981; Van de Weerd, 1976) se han quedado desactualizadas al haberse descrito muchas especies posteriormente. Por eso, es necesario acompañar al estudio taxonómico, de un estudio filogenético de los representantes de Cricetodontini.

Existen muy pocas cuencas ibéricas en las estén bien documentados los intervalos temporales más interesantes para el estudio de la tribu Cricetodontini (Aragoniense medio y superior, y Vallesiense inferior). Los mejores afloramientos se dan en la Cuenca del Duero (e. g., Álvarez-Sierra y García Moreno, 1986; López Martínez et al., 1986; García-Moreno, 1987; Álvarez-Sierra, 1987; Álvarez-Sierra et al., 1987; Álvarez Sierra et al., 1990; Krijgsman et al., 1996), en el Vallés-Penedés (e. g., Agustí, 1977, 1980; Agustí y Gilbert, 1982; Agustí et al., 2005; Alba et al., 2006; Casanovas-Vilar, 2007) y en la Cuenca de Madrid (e. g., Peláez-Campomanes et al., 2003; Morales et al., 2004; Hernández-Ballarín et al., 2010, 2011; López-Antoñanzas et al., 2010). Sin embargo, en la Cuenca de Calatayud-Daroca se dan las condiciones óptimas ya que presenta uno de los registros de micromamíferos fósiles mejor documentados y estudiados del Mioceno (e.g., Freudenthal, 1963; 1966; De Bruijn y Mein, 1968; Daams, 1974, 1981, 1990, 1999; Álvarez-Sierra, 1987; Cuenca Bescós, 1988; Daams y Freudenthal, 1988; De Jong, 1988; Freudenthal y Daams, 1988; Daams et al., 1999a, 1999b, 1999c; Van den Hoek Ostende, 2003, 2012; Van der Meulen et al., 2003; 2012; Hordijk y Van der Meulen, 2007; López-Guerrero et al., 2008, 2009, 2011b, 2013a; García-Paredes et al., 2009, 2010; Hordijk, 2010; Oliver y Peláez-Campomanes, 2013).

El excepcional registro de las faunas de micromamíferos, así como la continuidad estratigráfica de las secciones estudiadas, ha permitido establecer en la Cuenca de Calatayud-Daroca la referencia de un piso continental para el Cenozoico, el Aragoniense (Daams et al., 1977; Daams y Freudenthal, 1981), establecido en el área de Villafeliche, en Zaragoza. Está definido y subdividido en función de las faunas de roedores. En la actualidad se cuenta con estudios de paleomagnetoes-tratigrafía (Krijgsman et al., 1994, 1996; Garcés, et al. 2003) que correlacionan las secciones datadas con yacimientos de micromamíferos con la Escala Temporal de Polaridad Geomagnética (cuyas

siglas en inglés son GPTS). La suma de todos estos factores de superposición estratigráfica, de ordenación bioestratigráfica y correlación con la GPTS ha permitido datar numéricamente la gran mayoría de los yacimientos estudiados (Daams et al., 1999c; Van Dam et al., 2006; Van Dam et al., en prensa). Por todo lo anteriormente discutido, se ha considerado que la Cuenca de Calatayud-Daroca es la más idónea para estudiar las faunas de Cricetodontini. En ella se encuentran un gran número de yacimientos datados en el intervalo seleccionado para este estudio (Aragoniense medio-Vallesiense inferior).

Los trabajos asociados a la presente tesis doctoral que se han ido publicando (López-Guerrero et al., 2008, 2009, 2013a, 2013b, 2014; en prensa) ponen al día la taxonomía de los géneros *Cricetodon* e *Hispanomys* en la Cuenca de Calatayud-Daroca. Sin embargo, parte del material de Cricetodontini de la Cuenca de Calatayud-Daroca ya se estudió con anterioridad, tanto en trabajos taxonómicos (e.g., Freudenthal, 1963, 1966; Mein y Freudenthal, 1971a) como bioestratigráficos (e.g., Daams y Freudenthal, 1988; Daams et al., 1999b). También se citó la presencia de Cricetodontini en varios yacimientos, sin describir el material en detalle (Daams et al., 1999c; Álvarez-Sierra et al., 2003; López-Guerrero et al., 2011b). Por otro lado, también están estudiados los Cricetodontini de la Cuenca de Teruel-Alfambra, próxima a Calatayud-Daroca (Van de Weerd, 1976; Van Dam, 1997), así como los del resto de cuencas de la Península Ibérica y Europa (e.g., De Bruijn et al., 1993; Aguilar et al., 1994, 2007; Rummel, 1995, 2000, 2001; De Bruijn y Ünay, 1996; Rummel y Kálin, 2003; Agustí et al., 2005; Hernández Fernández et al., 2006; López-Antoñanzas y Mein, 2009, 2011; López-Antoñanzas et al., 2010). Por lo que se cuenta con muchos trabajos que sirven de apoyo y consulta para esta investigación.

Teniendo en cuenta todo lo anterior, en el presente trabajo se realizará una revisión taxonómica y sistemática de las especies de Cricetodontini de la Cuenca de Calatayud-Daroca. Con esta información se evaluará la utilidad bioestratigráfica de los Cricetodontini dentro de la cuenca y se discutirán las relaciones filogenéticas de cada una de las especies encontradas. A continuación se detallarán los mismos.

Capítulo 2

Objetivos



OBJETIVOS

"If you watch animals objectively for any length of time, you're driven to the conclusion that their main aim in life is to pass on their genes to the next generation."

David Attenborough

2

Como ya se ha comentado anteriormente, la tribu Cricetodontini tiene un gran interés, especialmente durante el Aragoniense medio y superior y del Vallesiense inferior (Mioceno medio y superior), y por ello se ha planteado su estudio en la Cuenca de Calatayud-Daroca. El principal objetivo de esta tesis es abordar el estudio sistemático y bioestratigráfico del grupo. Para ello hay que determinar taxonómicamente las especies presentes en cada uno de los yacimientos, evaluar su utilidad bioestratigráfica y hacer un análisis cladístico que nos ayude a entender las relaciones filogenéticas entre dichas especies.

De esta forma, los objetivos que, de forma efectiva, se pretenden cumplir en esta tesis doctoral son los siguientes:

1. Revisar la taxonomía de la tribu en el intervalo de tiempo estudiado. Para este primer objetivo, hay que llevar a cabo los aspectos particulares propios de un estudio sistemático:
 - Conocer la variabilidad morfológica y métrica del material de cada yacimiento para comprobar si los fósiles pertenecen a una sola especie o a varias.
 - Agrupar los especímenes que presentan la misma morfología y talla, para caracterizar su variabilidad.
 - Comparar cada uno de estos grupos morfológicos con todas de las especies de Cricetodontini descritas hasta la fecha para identificar a qué especie pertenecen o si se trata de una especie nueva.

- Discutir todos los aspectos taxonómicos de las especies encontradas en la cuenca, enmendando su diagnosis, si procediera.

A partir del estudio de la variabilidad en morfología y talla de cada uno de los molares, se realizará una asignación taxonómica de los fósiles de *Cricetodontini* del Aragoniense medio y superior, y del Vallesiense inferior.

2. Como segundo objetivo se ha fijado evaluar la utilidad bioestratigráfica de las especies. Debido a su gran complejidad y a la dificultad de establecer los límites morfológicos entre especies, los miembros de *Cricetodontini* se han usado tradicionalmente en el área estudiada únicamente a nivel de género y para caracterizar sólo grandes intervalos temporales. Con la determinación taxonómica de las especies en la Cuenca de Calatayud-Daroca y el estudio de sus distribuciones estratigráficas, se podrá evaluar su utilidad bioestratigráfica.
3. El tercer objetivo consiste en llevar a cabo un análisis filogenético de las especies de *Cricetodontini* mediante la metodología cladística. Para ello, se ha tenido que completar el primer objetivo y conocer la variabilidad morfológica y métrica de los miembros de *Cricetodontini*. Al usar el método cladístico, se han tenido que decidir aspectos esenciales como:
 - Establecer qué especies van a formar parte del análisis, entre el resto de especies de *Cricetodontini* de Europa, Asia Menor y China.
 - Decidir qué caracteres son los más idóneos para describir tanto la variabilidad genérica como la específica.

A partir de los resultados del estudio filogenético, se obtendrá el cladograma más parsimonioso que servirá de base para la comprobación de las hipótesis filogenéticas previas a este trabajo y durante el transcurso del mismo. Igualmente se propondrán nuevas hipótesis de parentesco de los *Cricetodontini* del Mioceno medio y superior de Calatayud-Daroca.

Capítulo 3

Contexto Geológico



CONTEXTO GEOLÓGICO

“Geology is the science which investigates the successive changes that have taken place in the organic and inorganic kingdoms of nature; it enquires into the causes of these changes, and the influence which they have exerted in modifying the surface and external structure of our planet.”

Sir Charles Lyell

INTRODUCCIÓN

3

El área estudiada en la presente tesis doctoral se encuentra en el sector NE de la Península Ibérica, en la provincia de Zaragoza, entre las localidades de Calatayud y Nombrevilla (Figura 3.1). Geológicamente, forma parte del complejo de grábenes tectónicos sedimentarios situados entre Calatayud y Teruel. Este gran complejo está formado por dos grandes subfosas que presentan direcciones estructurales diferentes, aunque actualmente estén conectadas (Sanz Rubio, 1999). Estas grandes subfosas se conocen como las cuencas de Alfambra-Teruel-Ademuz y de Calatayud-Montalbán (Ferreiro y Ruiz, 1991). La Cuenca de Calatayud-Montalbán está formada por dos subcuencas: la de Calatayud, también llamada de Calatayud-Daroca (Colomer y Santanach, 1988; Anadón y Moissenet, 1996) y la de Montalbán, separadas durante gran parte del Neógeno por el umbral de Daroca (Sanz Rubio, 1999; Sanz-Rubio et al., 2003). Como consecuencia de esta separación, los sedimentos que las rellenan son de diferentes edades: la Cuenca de Montalbán, en el sector meridional, contiene depósitos generalmente más antiguos y la Cuenca de Calatayud-Daroca, en el sector septentrional, cuenta con sedimentos más modernos (Sanz Rubio, 1999; Sanz-Rubio et al., 2003). Esta última contiene todos los yacimientos estudiados en esta tesis doctoral.

La Cuenca de Calatayud-Daroca cuenta con una extensión de unos 60 Km de largo, es relativamente estrecha, entre 10 y 26 Km de ancho, y tiene una orientación NO-SE (Alcalá et al., 2000). Está limitada por los macizos paleozoicos de Ateca-Daroca en el borde occidental y de Calatayud-Montalbán en su límite oriental (Ferreiro y Ruiz, 1991; Álvarez-Sierra et al., 2003).

GEOLOGÍA



FIGURA 3.1. Situación geográfica de las cuencas de Calatayud-Montalbán y Alfambra-Teruel-Ademuz. En detalle las subfosas de Calatayud-Daroca y Montalbán (Modificado de Gómez Cano et al., 2011).

La Cuenca de Calatayud-Daroca se formó durante el Neógeno (Cenozoico) en un régimen distensivo como resultado de los cambios de la zona de convergencia entre África y Eurasia desde los Pirineos hasta la Cordillera Bética (Alcalá et al., 2000). Esta cuenca pertenece a un grupo importante de depocentros que forman parte de un sistema extensional de larga amplitud y todavía activo que une los grábenes de Valencia con los de Francia y Centroeuropa (Alonso-Zarza et al., 2002). Las rocas sedimentarias del Cenozoico en España fueron depositadas durante y después de la compresión alpina en el área ibérica (Alonso-Zarza et al., 2002).

La estructura de la cuenca es compleja y presenta dos grandes variaciones (Julivert, 1954; Ferreiro y Ruiz, 1991; Alcalá et al., 2000). En el margen norte (área de Calatayud) la estructura tiene

forma de gráben, mientras que el margen suroeste presenta fallas *strike-slip*, como la falla del Jiloca, de dirección NO-SE (Julivert, 1954; Ferreiro y Ruiz, 1991; Anadón y Moissenet, 1996; Alcalá et al., 2000) que están acompañadas de cabalgamientos de bajo ángulo del basamento sobre depósitos aluviales sintectónicos (Colomer y Santanach, 1988; Casas et al., 2003).

Los depósitos del Cenozoico continental de las cuencas de la Península Ibérica han sido objeto de muchos estudios estratigráficos, sedimentarios y paleontológicos (e.g., López Martínez, 1987; Calvo et al., 1993; Alonso-Zarza et al., 2002). Igualmente, los trabajos centrados en los depósitos terciarios de la Cuenca de Calatayud-Daroca han sido numerosos (e.g., Julivert, 1954; Colomer y Santanach, 1988; Daams et al., 1999b, 1999c; Sanz Rubio, 1999; Alonso Zarza y

Calvo, 2000; Alcalá et al., 2000; Casas et al., 2003; Sanz-Rubio et al., 2003; Van Dam et al., en prensa). El relleno sedimentario de la Cuenca de Calatayud-Daroca comprende materiales detríticos de carácter aluvial (conglomerados, areniscas, limos y arcillas) en los sectores es de la cuenca, donde se acumulan potencias de hasta 1.200 m (Sanz-Rubio et al., 2003). El Neógeno de la Cuenca de Calatayud-Daroca ha sido dividido en tres unidades litoestratigráficas (Inferior, Intermedia y Superior) separadas por rupturas sedimentarias. No se han encontrado en la cuenca afloramientos datados de materiales previos al Ramblense (Mioceno inferior) (Sanz-Rubio et al., 2003). Sin embargo, se ha considerado que el inicio de la sedimentación tuvo lugar durante el Paleógeno, probablemente durante el Oligoceno (Sanz Rubio, 1999; Sanz-Rubio et al., 2003). Los yacimientos estudiados en la presente tesis doctoral se encuentran situados en la citada Unidad Intermedia que está formada por un gran espesor de yesos laminados y depósitos dolomíticos en el centro de la cuenca, mientras que en las áreas de borde de cuenca, hay localizados sistemas lacustres extensos y aislados (Sanz Rubio, 1999). La edad asignada a la Unidad Intermedia es Aragoniense medio-Vallesiense (Sanz-Rubio et al., 2003).

Los depósitos de la sección tipo del Aragoniense se encuentran situados al este de Villafeliche. Se sitúan encima de formaciones Paleozoicas, generalmente relacionados a través de una discordancia, aunque localmente hay contactos por fallas. La estructura del área tipo es, en general, relativamente simple. Se trata de una antiforma cuyo flanco sur está afectado tectónicamente, plegándose mientras que el flanco norte sufre un fallado normal. De esta manera, cerca del margen Paleozoico, la base del Mioceno presenta un buzamiento de aproximadamente 50° NE. Este buzamiento disminuye rápidamente hacia el área de la Rambla de Vargas (Figura 3.2). En el margen norte, los depósitos Aragonienses buzando progresivamente hacia el NE y se vuelven casi verticales como resultado de una falla (Daams et al., 1999c; Figura 3.2). En las áreas es de la Cuenca de Calatayud-Daroca, los materiales de edad Aragoniense corresponden a depósitos de evaporitas, fundamentalmente yesos laminados y margas que pasan a unidades de carácter carbonatado en las partes más altas (Sanz Rubio, 1999; Alcalá et al., 2000). Finalmente, en el lado noroeste de la Cuenca de Calatayud-Daroca, los depósitos de edad Aragoniense son materiales siliciclásticos constituidos por limos y arenas rojizas que culminan en un tramo de calizas (Alcalá et al., 2000)

Se han descrito tres unidades litoestratigráficas no formales dentro del Aragoniense: Valdemoros-Vargas, Las Umbrías y Las Planas. Parte de los yacimientos estudiados en la presente tesis doctoral se han encontrado en las unidades de Las Umbrías y Las Planas. A continuación se detalla su litoestratigrafía.

Unidad de Las Umbrías—Esta unidad contiene calizas tobáceas, margas verdosas y grises, y carbonatos nodulares. El espesor puede alcanzar unos 22 m. La parte superior consiste en una capa de carbonatos nodulosos finos que se encuentran depositados sobre una capa de lutitas




FIGURA 3.2. Mapa geológico y columna estratigráfica sintética del área tipo del Aragoniense en la Cuenca de Calatayud-Daroca (Zaragoza). En el mapa se señalan los principales yacimientos fosilíferos (Modificado de Daams et al., 1999c).

rojas que se puede seguir lateralmente en casi toda el área y representa un buen nivel guía. Estos depósitos carbonatados y margosos se formaron en un ambiente palustre en el que pequeños lagos someros alternaban con períodos de exposición subaérea (Platt y Wrigth, 1992; Alonso Zarza et al., 1992). El carácter somero de los lagos se inferido a partir de la presencia común de estructuras relacionadas con suelos, como bioturbación por raíces, estructuras hidromórficas y nodulares (Daams et al., 1999c).

Unidad de Las Planas—Esta unidad está compuesta por lutitas rojas con intercalación de margas y calizas. Las lutitas rojas afloran en forma de grandes capas, de hasta 10 m de espesor. Las margas verdes y grises aparecen asociadas a los carbonatos, generalmente por encima o debajo de éstos. Los carbonatos son dolomicritas nodulares típicas que presentan huellas de desecación o de bioturbación por raíces; pueden tener de 0,5 a 1,80 m. El espesor total puede llegar a los 110 m. Esta asociación de facies sugiere que los depósitos se acumularon en zonas de bajo relieve en partes distales de abanicos aluviales. Los carbonatos y margas se depositaron en lagos o charcas muy someros que experimentaron frecuentes episodios de exposición subaérea (Daams et al., 1999c).

En cuanto a la litoestratigrafía y sedimentología de los depósitos de edad Vallesiense inferior en la Cuenca de Calatayud-Daroca, se encuentran lutitas rojas, con intercalaciones locales de gravas y arenas, que se alternan con niveles carbonatados en los que predominan calizas nodulosas y acumulaciones de oncolitos y tobas (Alcalá et al., 2000). Esta asociación de facies correspondería a un ambiente de abanicos aluviales que se intercalan con lagos carbonatados muy someros y con periodos largos de exposición subaérea (Alcalá et al., 2000).

Secciones estratigráficas estudiadas

Las prospecciones realizadas la Cuenca de Calatayud-Daroca han hecho posible localizar un gran número de yacimientos de edad Aragoniense medio, superior y Vallesiense inferior. De éstos, 54 contienen representantes de la tribu Cricetodontini, la mayoría de ellos se agrupan en varias secciones estratigráficas. Las tres secciones más importantes en las que se han encontrado los yacimientos de este estudio son: Parte de la sección tipo del Aragoniense (en concreto, la sección de Valdemoros-Las Umbrías-Las Planas), Toril-Nombrevilla y Cañada (Daams et al., 1999c; Álvarez-Sierra et al., 2003; López-Guerrero et al., 2011b).

Sección tipo del Aragoniense—(Figuras 3.2; 3.3) esta sección está compuesta por otras tres parcialmente solapadas. Todas ellas situadas en la Rambla de Vargas, próxima al municipio de Villafeliche. La sección de Valdemoros es la parte inferior de la sección tipo y se encuentra al sur de la Rambla de Vargas; la sección de Las Umbrías continúa la sección de Valdemoros al norte de la rambla, y, finalmente, la sección de Las Planas es la parte superior de la sección tipo (Daams et




FIGURA 3.3. (a) Tramo medio y superior de la sección tipo del Aragoniense en las áreas de Las Umbrías y Las Planas. Las columnas muestran detalles de las estructuras sedimentarias, los yacimientos están señalados mediante el símbolo de un hueso. Modificado de Daams et al., 1999c (Fotografía (c) de A. Martín Monge). (b) Vista general de los yacimientos Las Umbrías. (c) Yacimiento Las Planas 5H.




FIGURA 3.4. (a). Columnas estratigráficas de Los Andurriales, Pedregueras, Toril y Nombrevilla, correlacionadas con la escala cronoestratigráfica (Modificado de Van Dam et al., en prensa). (b). Vista general de la sección de Nombrevilla. (Fotografía de A. Martín Monge).




FIGURA 3.5. Vista general de los yacimientos Toril 3A (a); Toril 3B (b); Nombrevilla 2 (c); Nombrevilla 3 y Nombrevilla 4 (d).
(Fotografías de A. Martín Monge)

al., 1999c; Van der Meulen et al., 2012). En total, se han encontrado 37 yacimientos de los cuales 14, de edad Aragoniense medio y superior, se han estudiado en la presente tesis doctoral (Daams et al., 1999c; Van der Meulen et al., 2012).

Sección Toril-Nombrevilla—(Figuras 3.4; 3.5) esta sección está compuesta a su vez por otras tres. La sección de Toril, que incluye las localidades de Toril y Pedregueras, está situada al NO próxima a la ciudad de Daroca. Se completa con otras dos secciones situadas más al SE, en los alrededores de la localidad de Nombrevilla (Alcalá et al., 2000; Álvarez Sierra et al., 2003). Se han estudiado los 17 yacimientos de edad Aragoniense superior y el Vallesiense inferior (Álvarez-Sierra et al., 2003).

Sección Cañada—está situada al SE de la sección tipo del Aragoniense, entre ésta y la sección de Toril-Nombrevilla. Consta de nueve yacimientos, seis de ellos de edad Vallesiense y dos de edad Turolense que no se han incluido en este estudio (López-Guerrero et al., 2011b).

Otras secciones también importantes, pero de menor extensión son: Los Andurriales, Valalto, Armantes, Alcocer y Paje. Finalmente, se incluyen 5 localidades aisladas: Manchones, Borjas, Arroyo del Val 6, Villafeliche 9 y Solera. (Alcalá et al., 2000; Álvarez Sierra et al., 2003; Garcés et al., 2003; López-Guerrero et al., 2008; Van der Meulen et al., 2012; Van Dam et al., en




FIGURA 3.6. Mapa y ortofoto con la situación geográfica de las secciones Cañadas, Andurriales, Pedregueras, Toril y Nombrevilla (Modificado de Van Dam et al., en prensa).

CONTEXTO CRONOLÓGICO Y BIOESTRATIGRÁFICO

prensa; Figura 3.6).

Como se ha explicado en la introducción, muchos de los yacimientos estudiados están estratigráficamente superpuestos. Este hecho no es muy frecuente en el caso de las localidades de vertebrados continentales y ha permitido realizar la propuesta de una biozonación local. También se han podido hacer estudios de paleomagnetoe-stratigrafía (Krijnsman et al., 1994, 1996; Garcés et al., 2003; Van Dam et al., en prensa) y establecer la correlación con la escala magnetoestratigráfica. Por tanto, la gran mayoría de los yacimientos estudiados tienen dataciones numéricas. En esta tesis doctoral se han usado las edades propuestas por Daams et al. (1999b, 1999c), Van Dam et al. (2006) y Van Dam et al. (en prensa) recalibradas de acuerdo con la Astronomically Tuned Neogene Time Scale (ATNTS2004; Lourens et al., 2004; ATNTS2012; Hilgen et al., 2012). En total, el intervalo de tiempo que abarcan estos yacimientos es desde unos 15 Ma hasta unos 9 Ma aproximadamente (Daams et al., 1999b; Garcés et al., 2003; Van Dam et al., 2006; Van der Meulen et al., 2012; Van Dam et al., en prensa). Este intervalo de tiempo corresponde, en parte, a los pisos continentales Aragoniense y Vallesiense.

El piso continental Aragoniense fue propuesto y definido por Daams et al. (1977) en el área de la Rambla de Vargas (Figura 3.2) situada en las inmediaciones de la localidad de Villafeliche. Daams y Freudenthal (1981) y posteriormente Daams et al. (1987) plantearon una biozonación local y subdividieron el Aragoniense en seis zonas: B, C, D, E, F y G. Se incluyeron algunos cambios en los siguientes trabajos (Daams y Freudenthal, 1988) como fue la propuesta de subdivisión de las biozonas locales D (D1, D2 y D3) y G (G1, G2 y G3). En la última revisión integral del Aragoniense (Daams et al., 1999b) se estudiaron 108 yacimientos, se describieron detalladamente las unidades litoestratigráficas y la sección tipo del Aragoniense. También se proporcionaron las edades numéricas de las localidades siguiendo las correlaciones magnetoestratigráficas con la GPTS propuestas por Krijnsman et al. (1994, 1996) y dataron el Aragoniense entre 17 y 11 Ma. En ese trabajo se cambió la subdivisión de la biozona D, que pasó a ser Da, Db, Dc y Dd. Recientemente, Van der Meulen et al. (2012) hicieron una nueva revisión bioestratigráfica del Aragoniense inferior y medio. Pusieron al día el contenido faunístico de los yacimientos del área tipo del Aragoniense, con el que redefinieron las biozonas desde la B hasta la F, dividiendo la biozona C (Ca y Cb). Utilizando las nuevas edades numéricas y datos magnetoestratigráficos (Van Dam et al., 2006) dataron numéricamente los límites de las biozonas locales y los principales eventos bióticos, así como se definieron formalmente cada biozona.

El término *Vallesiense* fue definido por Crusafont en 1950 en la cuenca catalana del Vallès-Penedès para denominar a las primeras faunas europeas que contienen el équido *Hipparion*. El primer registro de *Hipparion* se ha considerado como el límite de este piso continental. Sin embargo, estudios magnetoestratigráficos en secciones de la Cuenca de Calatayud-Daroca (Garcés et al., 2003) y en la Cuenca del Vallès-Penedès (Garcés et al., 1996) sugirieron que la llegada de este équido a la Península Ibérica fue diacrónica, datando antes la primera presencia de *Hipparion* en la Cuenca del Vallès-Penedès que en la Cuenca de Calatayud-Daroca (Garcés et al., 2003).

El Vallesiense tiene dos divisiones: el Vallesiense inferior; dentro del cual se definieron las biozonas locales H e I (Daams y Freudenthal, 1981) para la Cuenca de Calatayud-Daroca, y el Vallesiense superior en el que Van de Weerd (1976) describió una única biozona en la Cuenca de Alfambra-Teruel-Ademuz, *Progonomys hispanicus*, que más tarde se subdividió en tres (J1, J2 y J3) (Van Dam, 1997; Van Dam et al., 2001). El final del Vallesiense inferior se ha considerado como un periodo de crisis (Agustí y Moyá Solá, 1990) en grupos como por ejemplo, los eomíidos, cricétidos y glíridos. Sin embargo, recientes trabajos (Casanovas-Vilar et al., 2005, 2013, 2014; García-Paredes y Casanovas-Vilar, 2013; Maddern et al., 2013) relativizan las dimensiones de la crisis del Vallesiense y justifican la conclusión sacada por Agustí y Moyá Solá (1990) en la escasez de yacimientos encontrados en el Vallesiense superior en contraposición con el Vallesiense inferior, en ese momento.

En la tabla 3.1 se presentan los yacimientos estudiados junto con su datación numérica calculada por Daams et al. (1999c), Van Dam et al. (2006) y Van Dam et al. (en prensa).

En la figura 3.7 se indica el marco cronológico y bioestratigráfico usado en esta tesis doctoral. Los yacimientos de este estudio están situados entre las biozonas locales E e I. A continuación se detallan cada una de estas biozonas señalando las especies que las definen.

La biozona E—definida por Daams y Freudenthal (1981), ha sido estudiada y redefinida recientemente por Van der Meulen et al. (2012). Está caracterizada por la presencia de dos taxones: la primera presencia común de *Heteroxerus grivensis* y de *Megacricetodon rafaeli*. También presenta como eventos característicos: la primera presencia de *Cricetodon* y una especie nueva de *Heteroxerus* no descrita. *Pseudodryomys ibericus*, *Simplomys simplicidens*, *Pseudodryomys rex*, *Megacricetodon collongensis* y *Democricetodon lacombai* tienen sus últimas presencias en esta biozona. El yacimiento más antiguo de la biozona es Las Umbrias 11 (14,06 Ma), y el más moderno, Las Umbrias 20 (13,80 Ma).

La biozona F—según Daams y Freudenthal (1981) se caracteriza por la presencia de dos especies de *Megacricetodon* que Daams et al. (1999c) determinaron como *M. gersii* y *M. rafaeli*. También se consideran taxones característicos: *Democricetodon darocensis* y *Cricetodon* (aunque son poco abundantes). La fauna de glíridos es pobre. El yacimiento más antiguo de la biozona es Las

Umbrías 22 (13,76 Ma), y el más moderno, Valalto 2C (13,68 Ma según Daams et al., 1999c).

La biozona G1—descrita por Daams y Freudenthal (1988) y redefinida por Daams et al (1999c), se caracteriza por la presencia de dos especies de *Megacricetodon*: *M. gersii* y *M. minor*. El yacimiento más antiguo de la biozona es Las Planas 5B (13,56 Ma) y el más moderno es Valalto 1 (13,30 Ma de acuerdo con Daams, 1999c).

La biozona G2—definida por Daams y Freudenthal (1988) y posteriormente redefinida por Daams et al. (1999c), se caracteriza por la presencia de faunas de cricétidos similares a la G1, pero *M. gersii* es sustituido por *M. crusafonti*. Los glíridos son poco abundantes, aunque muy diversos; puede encontrarse *Myomimus dehmi*, *Muscardinus thaleri*, *Myoglis meini* y dos

especies de *Microdyromys*. También se ha comentado la presencia del eomúido *Emyops* (Daams et al., 1999c). El yacimiento más antiguo es Manchones (13,25 Ma de acuerdo con Daams et al., 1999c), y el más moderno es Las Planas 5K (13,08 Ma).

La biozona G3—se definió en 1988 por Daams y Freudenthal y se redefinió por Daams et al. (1999c). Está caracterizada por la presencia de *Megacricetodon minor* y una asociación transicional entre *M. crusafonti* y *M. ibericus*. *Cricetodon* es relativamente frecuente y *Democricetodon darocensis* se reemplaza por *D. crusafonti* en la parte superior. Los glíridos son comparables a los de la G2, y los castores pueden estar presentes. El yacimiento más antiguo es Villafeliche 9 (12,85 Ma según Daams et al., 1999c) y el más moderno es Nombrevilla 2 (11,87 Ma).

La Biozona H—se describió en 1981 por Daams y Freudenthal. Pertenece en parte al Aragoniense superior y al Vallesiense inferior, se redefinió por Daams et al. (1999c) por la presencia de dos especies del género *Megacricetodon* diferentes a las presentes en la biozona local G3: *Megacricetodon ibericus* y *M. minor* o *M. debruijni*, así como por la presencia y mayor abundancia de *Hispanomys* frente a *Cricetodon*. El glírido *Myomimus dehmi* es frecuente pero *Democricetodon crusafonti*, *Muscardinus hispanicus* y los Castoridae no lo




FIGURA 3.7. Esquema gráfico del marco cronológico en el que están situados los yacimientos estudiados. Se presentan las correlaciones propuestas entre las épocas del Mioceno, los pisos continentales, las zonas de mamíferos neógenos y la biozonación local (Daams et al., 1999; Garcés et al., 2003; Van der Meulen et al., 2012; Van Dam et al., 2006; Van Dam et al., en prensa).

son. Hay dos yacimientos que, siendo de edad Aragoniense superior, pertenecen a la biozona H: Nombrevilla 9 y Nombrevilla 10. El resto (Tabla 3.1) son vallesienses (Álvarez-Sierra et al., 2003; Garcés et al., 2003; López-Guerrero et al., en prensa).

La Biozona I—se describió en 1981 por Daams y Freudenthal. Posteriormente se redefinió por Daams et al. (1999c) por la presencia de *Megacricetodon debruijni*. *Cricetulodon* e *Hispanomys* son comunes y *Democricetodon sulcatus* y los Castoridae son poco frecuentes. El glirido *Ramys multicrestatus* se registra por primera vez, *Muscardinus hispanicus* y *Myomimus dehmi* son relativamente comunes. *Hipparion* está presente. El yacimiento más antiguo de esta biozona es Nombrevilla 13 (10'72 Ma según Van Dam et al., en prensa) y el más moderno es Nombrevilla 19 (9,98 Ma según Van Dam et al., en prensa).

Localidad	Ab.	Biozona	Edad	Localidad	Ab.	Biozona	Edad
Nombrevilla 19	NOM19	I	9,98	Alcocer 3	AC3	G3	—
Pedregueras 2C	PED2C	I	10,38	Alcocer 2	AC2	G3	—
Pedregueras 2A	PED2A	I	10,62	Alcocer 1	AC1	G3	—
Pedregueras 1A	PDA	I	—	Villafeliche 9	VL9	G3	—
Nombrevilla 14	NOM14	I	10,68	Arroyo del Val VI	ARV6	G2	—
Nombrevilla 13	NOM13	I	10,72	Las Planas 5K	LP5K	G2	13,08
Nombrevilla 22	NOM22	I	—	Las Planas 5L	LP5L	G2	13,16
Cañada 11	CAÑ11	H	—	Borjas	BOR	G2	—
Cañada 10	CAÑ10	H	—	Manchones	MAN	G2	—
Cañada 9	CAÑ9	H	—	Valtalto 1	VT1	G1	13,30
Cañada 8	CAÑ8	H	—	Las Planas 5C	LP5C	G1	13,55
Nombrevilla 1	NOM1	H	10,77	Las Planas 5B	LP5B	G1	13,56
Nombrevilla 10	NOM10	H	11,18	Valtalto 2C	VT2C	F	13,68
Nombrevilla 9	NOM9	H	11,20	Valtalto 2B	VT2B	F	13,70
Carrilanga 1	CARR1	G3	11,33	Armantes 7	ARM7	F	—
Nombrevilla 4	NOM4	G3	11,80	Las Umbrías 21	LUM21	F	13,75
Nombrevilla 3	NOM3	G3	11,84	Las Umbrías 22	LUM22	F	13,76
Nombrevilla 2	NOM2	G3	11,87	Las Umbrías 20	LUM20	E	13,80
Solera	SOL	G3	12,01	Las Planas 4C	LP4C	E	13,88
Paje 2	PJE2	G3	12,26	Las Umbrías 19	LUM19	E	13,95
Paje 1	PJE1	G3	12,39	Las Planas 4B	LP4B	E	13,96
Las Planas 5H	LP5H	G3	12,60	Las Planas 4BA	LP4BA	E	—
Toril 3B	TOR3B	G3	12,65	Las Planas 4A	LP4A	E	13,98
Toril 3A	TOR3A	G3	12,65	Las Umbrías 16	LUM16	E	14,04
Toril 2	TOR2	G3	—	Las Umbrías 11	LUM11	E	14,06
Toril 1	TOR1	G3	—				

TABLA 3.1. Listado de los yacimientos estudiados, sus abreviaturas (ab.), la biozona a la que pertenecen y la edad calculada está basada en algunos casos en la correlación directa con la escala geomagnética. En otros casos, la edad está obtenida a partir de los yacimientos datados directamente y las tasas de sedimentación calculadas para estos depósitos. La metodología de cálculo de las edades se recoge en Krijgman et al, 1994; Daams et al., 1999b, 1999c.; Garcés et al., 2003 y Van Dam et al., en prensa. Las edades están recogidas de Daams et al. (1999c); Van Dam et al. (2006) y Van Dam et al., en prensa.

II. Material y Métodos

Capítulo 4

Material y Métodos



MATERIAL Y MÉTODOS

“Scientific method includes, in short, all the processes by which the observing and amassing of data are regulated with a view to facilitating the formation of explanatory conceptions and theories.”

John Dewey

MATERIAL

4

En el presente trabajo se han estudiado 3.181 molares de roedores cricétidos provenientes de 54 yacimientos. La tabla 4.1 muestra la información acerca del número de especímenes estudiados, el yacimiento al que pertenecen y la institución en la que están conservados. Para comparar el material de este estudio con las especies de Cricetodontini, se han estudiado directamente 925 fósiles de 12 localidades tipo, todas ellas situadas fuera de la Cuenca de Calatayud-Daroca (Tabla 4.2). En cada capítulo del apartado (Paleontología sistemática) se especifica qué material se ha estudiado y a qué institución pertenece. Además, se han estudiado 344 molares de las colecciones alemanas de la Cuenca de Antepaís Nordalpina, así como 90 fósiles de la Molasa de la baja Austria (Tabla 4.2). Algunos de estos fósiles centroeuropeos ya habían sido publicados (Boom, 1991; Rummel, 2001; Heißig, 2006; Daxner-Höck, 2003) y otros son inéditos, pero todos ellos han sido descritos y medidos para este trabajo. Estos fósiles han aportado información acerca de la variabilidad intraespecífica de determinadas especies de amplia distribución geográfica. En total, se han medido y descrito casi 4500 fósiles.




TABLA 4.1. Marco cronológico y bioestratigráfico de los yacimientos estudiados de la Cuenca de Calatayud-Daroca. Se incluye la información con la institución en la que están conservados y el número de especímenes estudiados separados por elemento dental. Abreviaturas: Yac., yacimiento; Ab., Abreviatura; Inst., institución; IAUU, Instituut voor Aardwetenschappen, Utrecht University (Países Bajos); MNCN, Museo Nacional de Ciencias Naturales-CSIC, Madrid (España); RGM, Naturalis Biodiversity Center, antiguo Rijksmuseum van Geologie en Mineralogie, Leiden (Países Bajos); N, Número de elementos. Las barras azules representan gráficamente, con escalado logarítmico, el número de fósiles de cada yacimiento.

Especie	Localidad	M1	M2	M3	m1	m2	m3	N	Institución	(a)
<i>C. meini</i>	Burtg-Balzhausen	11	12	1	10	—	10	44	NA	
<i>C. aff. meini</i>	Ebershausen	—	3	2	—	2	—	7	BSGP	
<i>Cricetodon</i> sp.	Gallenbach2b	27	18	8	18	16	9	96	BSGP	
<i>C. aureus</i>	Göttschlag	5	3	—	3	—	—	11	BSGP	
<i>Cricetodon</i> sp.	Hahnenberg	—	—	—	1	—	2	3	BSGP	
<i>C. cf. meini</i>	Höhenraunau	1	—	—	—	—	1	2	NA	
<i>Cricetodon</i> sp.	Höll	1	1	1	—	1	1	5	BSGP	
<i>C. cf. sansaniensis/C. cf. engesseri</i>	Kirrberg-Tongrube	4	5	2	6	5	2	24	NA	
<i>Cricetodon</i> sp.	Kleinensenbach	3	2	2	5	3	3	18	BSGP	
<i>C. aureus</i>	Laimering2	20	15	12	13	—	—	60	BSGP	
<i>C. aureus</i>	Laimering3	6	10	—	8	6	9	39	BSGP	
<i>C. aureus</i>	Laimering5	—	1	—	4	1	3	9	BSGP	
<i>Cricetodon</i> sp.	Mörge	2	1	2	—	1	2	8	NA	
<i>Cricetodon</i> sp.	Petersbuch 14	1	—	—	—	—	—	1	BSGP	
<i>C. aureus</i>	Unterneul 1a	3	1	1	—	—	2	7	BSGP	
Total Alemania	Total Alemania	84	72	31	68	35	44	334	Total Alemania	
<i>C. meini</i>	Mühlbach 1	2	5	5	3	7	1	23	NHW	
<i>C. meini</i>	Mühlbach 2	10	8	7	9	4	7	45	NHW	
<i>C. meini</i>	Goldberg, Nördlinger Ries/BRD	1	2	3	3	4	2	15	NHW	
<i>C. meini</i>	Grund	1	—	—	—	—	1	2	NHW	
<i>Cricetodon</i> sp.	Bullendorf	—	—	2	1	1	1	5	NHW	
Total Austria	Total Austria	14	15	17	16	16	12	90	Total Austria	
Especie	Localidad tipo	M1	M2	M3	m1	m2	m3	N	Institución	(b)
<i>'C'. fandli</i>	Gratkorn	2	1	3	1	1	3	11	BSGP	
<i>C. bolligeri</i>	Petersbuch 10	2	1	1	2	1	1	8	BSGP	
<i>H. lavocati</i>	Hostalets de Pierola	10	5	4	2	2	—	23	ICP	
<i>H. aguirrei</i>	Escobosa de Calatañazor	79	41	35	71	43	28	297	MNCN	
<i>'C'. klariankae</i>	Felsőtárkány-Felnémet	9	11	11	13	16	9	69	PM	
<i>H. bijugatus</i>	La Grive-Saint Alban, L3	10	5	5	6	5	5	36	RGM	
<i>H. decedens</i>	La Grive-Saint Alban, L5'	10	10	10	10	10	10	60	RGM	
<i>C. albanensis</i>	La Grive-Saint Alban, M	27	26	23	25	15	6	122	RGM	
<i>C. sansaniensis</i>	Sansan	9	6	8	15	—	—	38	MHNB	
<i>C. soriae</i>	SomosaguasN	—	1	1	1	2	3	8	UCM	
<i>C. meini</i>	Vieux-Collonges	25	33	66	36	6	6	172	RGM	
<i>C. aureus</i>	Vieux-Collonges	13	8	25	12	17	7	82	RGM	
Total	Total localidades tipo	196	148	192	194	118	78	926		

TABLA 4.2. Número de ejemplares estudiados para realizar la comparación con el material de Calatayud-Daroca. (a). Localidades del Mioceno medio de Europa Central. Los fósiles que están actualmente en supervisión se han citado bajo nomenclatura abierta como *Cricetodon* sp. Asignaciones taxonómicas obtenidas de Heißig, 1997, 2006; Rummel, 2000, 1995; Daxner-Höck, 2003; Prieto, 2007 y Seehuber, 2008. (b). Localidades tipo de algunas especies de *Cricetodontini*. La edad de cada una de estas localidades se especifica en la sección *Material and Methods* de cada uno de los capítulos del apartado III (Paleontología Sistemática). Abreviaturas: BSPG, Bayerische Staatssammlung für Paläontologie und Geologie, Munich (Alemania); MNCN, Museo Nacional de Ciencias Naturales-CSIC, Madrid (España); UCM, Universidad Complutense de Madrid, (España); RGM, Naturalis Biodiversity Center, antiguo Rijksmuseum van Geologie en Mineralogie, Leiden (Países Bajos); MHNB, Muséum d'Histoire Naturelle de Bâle (Suiza); PM, Páztó Múzeum, Páztó (Hungría); MHW, Naturhistorisches Museum Wien, Viena (Austria); NA, Naturmuseum Augsburg, Augsburg (Alemania).

MÉTODOS

MÉTODOS DE CAMPO

Los fósiles de la Cuenca de Calatayud-Daroca fueron recogidos en numerosas campañas de campo. La prospección de los yacimientos de esta tesis doctoral, así como la recolección del material estudiado, se llevó a cabo durante las campañas dirigidas por el Dr. Remmert Daams desde la década de 1980 y que continuó el equipo de paleontólogos de micromamíferos que trabajan conjuntamente desde la UCM y del MNCN-CSIC hasta los años 2000. A continuación, se resumen los aspectos más importantes en las labores de campo en paleontología de micromamíferos.

La gran mayoría de los yacimientos de este estudio son de origen fluviolacustre (Alcalá et al., 2000; Álvarez-Sierra et al., 2003; Azanza et al., 2004; López-Guerrero et al., 2011b). Debido a la baja densidad de fósiles que presentan este tipo de yacimientos, es necesario recoger muestras muy grandes de sedimento para su procesado. Se recogen pruebas y se procesa el sedimento de manera que si el resultado es positivo, se explota el yacimiento. El número de fósiles que se deben encontrar para considerar una muestra positiva varía según el interés de los mismos. El tamaño de la muestra final que se recoge varía igualmente en función del interés de los fósiles encontrados. A continuación la muestra pasa por una serie de procesos en los que el sedimento se seca (Figura 4.1) y se rehidrata para pasarlo después por la mesa de lavado-tamizado similar a la descrita por Daams y Freudenthal (1988) (Figura 4.2a). Todas estas técnicas de tratamiento de fósiles de microvertebrados se pueden encontrar descritas en detalle en numerosos trabajos previos (Daams y Freudenthal, 1988; Álvarez-Sierra, 1987; López-Martínez, 1992; Peláez-Campomanes,



FIGURA 4.1. Proceso de secado del sedimento previo al lavado-tamizado.

1993; García Paredes, 2006; Casanovas-Vilar, 2007). Una vez terminado el proceso de lavado, el concentrado mayor de 0,7 mm se deja secar en tamices (Figura 4.2b) y se guarda en sacos debidamente siglados para su almacenaje y posterior procesamiento en el laboratorio.



FIGURA 4.2. (a). Vista general y detalles de la mesa de lavado ideada por M. Freudenthal. El agua a presión es conducida por las mangueras y un sistema de tolvas hace que el sedimento pase de un tamiz a otro. (b). Proceso de secado después del lavado-tamizado.

MÉTODOS DE LABORATORIO

Habitualmente, en el laboratorio se realizan nuevos lavados de la muestra utilizando distintos reactivos. En el caso de las muestras de la Cuenca de Calatayud-Daroca, normalmente, se utiliza el ácido acético diluido al 10%. El ácido reacciona con el carbonato, de manera que se reduce sustancialmente el tamaño de la fracción carbonatada y, por tanto, también del concentrado final. Posteriormente, se seca la muestra mediante un horno y se realiza el fraccionamiento de la misma mediante tamices, usando una tamizadora o a mano (Figura 4.3). Así se han separado las distintas fracciones (>2 mm, 2-1,5 mm, 1,5-1 mm, 1-0,7 mm) para cada muestra.

A continuación se realiza el triado. En este proceso se separan los fósiles de los granos de arena o restos vegetales. Para ello, se necesita la ayuda de una lupa de visera o una lupa binocular para los tamaños menores de 2 mm, mientras que la fracción más grande se puede separar a simple vista.



FIGURA 4.3. Métodos de relavado en el laboratorio. Tratamiento con ácido acético, tamizado con cedazos a mano o con tamizadora (Fotografías de A. R. Gómez Cano).

Cuando ya se tienen todos los restos fósiles aislados, se montan y siglan. Como sistemas de almacenaje en unas ocasiones, se usan cápsulas de plástico en las que se mete individualmente cada resto. En otras se colocan los pequeños restos de los micromamíferos sobre masas plásticas flexibles que se montan sobre un soporte rígido como puede ser un portaobjetos o una ficha de *Lego*, por ejemplo. La gran ventaja del uso de la masa flexible es que no se toca el fósil para su estudio, y además, permite ocupar menos espacio para almacenar las piezas, ya que se pueden siglar hasta 50 ejemplares en un mismo portaobjetos. La desventaja de este tipo de montaje radica en el tipo de masa que se use, ya que se ha comprobado que algunas masas flexibles (por ejemplo, la parafina) son inadecuadas porque reaccionan con la dentina, destruyéndola por completo en algunos casos. En los últimos años se ha usado un tipo de masa limpiatipos *Pelikan* que es bastante inerte, no es grasa y es muy moldeable. Sin embargo, este limpiatipos ya no se fabrica y se está sustituyendo por goma de borrar carboncillo, fabricada en caucho sintético.

ESTUDIO DEL MATERIAL

Aspectos generales

En primer lugar, se ha descrito la morfología dental de los fósiles, especialmente los caracteres de la la superficie oclusal, tal y como es habitual en el estudio de los roedores fósiles (Sesé Benito, 1977; Agustí, 1981; Álvarez-Sierra, 1987; García-Moreno, 1987; Martínez-Salanova, 1987; Cuenca Bescós, 1988; Lacombe, 1988; Martín-Suárez, 1988; Sevilla, 1988; Herráez, 1993; Peláez-Campomanes, 1993; García-Alix, 2006; Minwer-Barakat, 2006; García Paredes, 2006;



FIGURA 4.4. Lupas binoculares utilizadas para el triado y estudio de los fósiles. (a). Olympus wild 308700 (50x) del Museo Nacional de Ciencias Naturales. (b). Leica 50x del *Naturalis Biodiversity Center*, (Leiden).

Casanovas-Vilar, 2007; Oliver y Peláez-Campomanes, 2013). Las observaciones se han hecho a través de varias lupas binoculares (wild 308700 (50x) y Leica 50x) (Figura 4.4).

También se ha estudiado la talla de todas las piezas dentales, midiendo su longitud y anchura máxima. Con las medidas, se han realizado análisis estadísticos siempre que el número mínimo de molares lo haya hecho posible. Se ha usado el programa SPSS (Statistical Package for the social Sciences) en la versión 11.5.1 (SPSS, 2002).

Las fotografías de los molares se han llevado a cabo en el *Laboratorio de microscopía electrónica* del MNCN-CSIC. Se utilizó un microscopio electrónico de barrido ambiental FEI Quanta 200 (Figura 4.5). Se han construido varias láminas de cada una de las especies estudiadas que incluyen ejemplares de uno o varios yacimientos, según el caso.



FIGURA 4.5. Equipo de microscopía electrónica utilizado para hacer las fotos usadas en las láminas de los capítulos del apartado III Paleontología sistemática (Fotografía de J. García Guinea).

Nomenclatura

Los dientes de los cricétidos presentan una corona dental baja (dentición braquidonta), con un patrón de cúspides separadas por valles (dentición de tipo bunodonta). Se ha seguido la nomenclatura propuesta por Mein y Freudenthal (1971b) y modificada por López-Guerrero et al. (2013a) (Figura 4.6). En los cricétidos, el patrón dental está compuesto por cinco cúspides principales en los primeros molares, tanto superiores como inferiores, que pasan a ser cuatro en los segundos y terceros molares. Puede aparecer un número variable de crestas o lofos conectando las cúspides. Así mismo, pueden aparecer pequeñas cúspides accesorias en los bordes del diente, cuyo desarrollo es mucho menor que el de las cúspides principales.




FIGURA 4.6. Terminología de las partes de los dientes yugales de Cricetodontini propuesta por Mein y Freudenthal (1971b) y modificada por López-Guerrero et al. (2013a). Sólo el M1, M3 y m1 se han dibujado, pero esta nomenclatura se puede aplicar al resto de los molares inferiores y superiores.

Caracteres morfológicos

A continuación, se presenta una descripción detallada de algunos de los caracteres utilizados en los estudios morfológicos presentados en el apartado (Paleontología sistemática). El resto de caracteres y sus estados de carácter se ilustran a lo largo de los siguientes capítulos tanto en las tablas de morfología de los apéndices del capítulo 8, como en las figuras y tablas del capítulo 10.

Se han estudiado un total de 82 caracteres morfológicos, que presentan entre todos 282 estados de carácter. Éstos se reparten de la siguiente manera:

M1: 54 estados de carácter agrupados en 16 caracteres.

M2: 55 estados de carácter, repartidos entre 16 caracteres.

M3: 40 estados de carácter de 12 caracteres.

m1: 56 estados de carácter, recogidos de 16 caracteres.

m2: 41 estados de carácter, agrupados en 11 caracteres.

m3: 36 estados de carácter, repartidos entre 11 caracteres.

Para los 4.500 molares estudiados, esto supone un total de más de 150.000 observaciones realizadas durante el presente trabajo.

La mayoría de los caracteres estudiados son comunes a todos los cricétidos, aunque algunos de ellos son característicos de los representantes de la tribu Cricetodontini, como los ectolofos anteriores, los grandes estilos en el protoseno de los molares superiores o el metalofúlido posterior en el primer molar inferior.

Molares superiores—Los molares superiores tienen gran valor informativo y acaparan un tercio de los caracteres usados en la diagnosis de los géneros estudiados. Llama la atención las diferencias que hay en el contorno de diferentes especies, que puede ser recto o sinuoso (=trilobulado) (Figura 4.7); esta diferencia está relacionada con dos aspectos: la presencia o ausencia de cíngulos en el borde labial de los dientes y la de ectolofos completos, tanto el anterior como el posterior. De esta manera, el contorno sinuoso o trilobulado estaría presente en especímenes que no poseen cíngulos en los valles labiales y cuyos ectolofos están completos, creando una cresta continua.




FIGURA 4.7. Tipos de contorno en el M1

M1

Anterocono/Anterocone: esta cúspide se encuentra en el borde anterior del diente y está situada en su eje longitudinal. Está dividido en dos lóbulos por un surco. En algunas especies, este surco no llega hasta la base de la corona, por lo que en especímenes muy desgastados puede parecer que no está dividido, sino que es alargado. Sin embargo, el anterocono nunca es una cúspide simple del tipo que presentan otros cricétidos como *Democricetodon* (Figura 4.8).

Anterolofos lingual o labial/Lingual or labial anteroloph: en algunas ocasiones, el anterocono presenta una cresta de esmalte que, partiendo desde el ápice de la cúspide, recorre el protoseno o el




FIGURA 4.8. Tipos de división del anterocono

anteroseno. A estas crestas se les da el nombre de anterolofos lingual y labial, respectivamente. Es muy importante diferenciar éstas de los cíngulos que aparecen en los bordes del protoseno y del anteroseno. El cíngulo es un recrecimiento del esmalte en el borde del diente que une dos cúspides, pero que no está preferentemente asociado a ninguna de ellas. Por su parte, un anterolofos está claramente asociado al anterocono, parte de éste y recorre el valle. El anterolofos lingual puede estar ausente (Figura 4.9a), ser “corto” (Figura 4.9b) o “largo” (Figura 4.9c), mientras que el anterolofos labial suele ser siempre corto.

Espina lingual o labial del anterolofulo/Lingual or labial spur of the anterolophule: el antero-






FIGURA 4.9. Desarrollo del anterolofos lingual


Espina labial



Espina lingual

FIGURA 4.10. Espina labial y lingual del anterolóbulo

lóbulo es la cresta de esmalte que une el protocono con el anterocono. Esta cresta puede desarrollar dos espinas que se nombran según su dirección lingual o labial. La espina lingual del anterolóbulo suele ser muy larga y llegar hasta el borde anterolingual del protoseno, mientras que la labial puede presentar longitudes variadas. Si es corta (no recorre más de la mitad de la longitud del anteroseno) no recibe ningún nombre en especial. Esta cresta labial nunca toca el borde labial del diente. Esto ocurre por dos razones: porque nunca llega a ser tan larga y porque cuando se desarrolla más, se curva posteriormente y se une al protolóbulo (Figura 4.10). Cuando ocurre esto último, puede hablarse de un “protolóbulo anterior”. Ambas estructuras, “protolóbulo anterior” y “espina labial del anterolóbulo conectada con el protolóbulo”, son muy difíciles de distinguir. En los capítulos 5-8 se dice que un molar tiene “espina labial de anterolóbulo” cuando es corta y no está conectada al protolóbulo mientras que cuando está conectada, se define como “protolóbulo anterior”.



Espina del Protocono

FIGURA 4.11. Espina del protocono

Espina del protocono/*Protocone spur*: el protocono puede presentar una espina en su cara posterior. Esto hace, que su forma no sea simétrica. La presencia de esta espina determina que el recorrido del seno adquiera una forma curvada anteriormente, mientras que en su ausencia es recto (Figura 4.11).

Protoseno/*Protosinus*: este valle, situado entre el anterocono y el protocono, puede presentar varias estructuras en el borde. Se han descrito éstas como “estructuras que cierran el protoseno” y pueden ser: un cíngulo, en el sentido de lo anteriormente descrito; un anterolofio lingual que puede o no cerrar el valle, o un estilo, que es una pequeña cúspide de esmalte que se sitúa en el borde anterolingual. En algunas especies, como se explica en el capítulo 5, este estilo es muy diagnóstico debido a su frecuente presencia y su gran tamaño.

Ectolofio anterior/*anterior ectoloph*: se da el nombre de ectolofio anterior a la estructura longitudinal situada entre el anterocono y el paracono. Esta estructura puede estar formada por una cresta que parte del anterocono, por una cresta que sale del paracono o por ambas. Se les ha

nombrado tomando como referencia: (1) la cúspide a la que esté asociada y (2) la situación anterior o posterior de la cara desde la que parten. De esta manera, a la cresta asociada al anterocono que sale desde la cara posterior de éste se ha llamado “ectolofos posterior del anterocono” (Figura 4.12b). Igualmente, a la cresta que saliendo desde la cara anterior del paracono, se desarrolla en el espacio entre el anterocono y el paracono, se le ha denominado “ectolofos anterior del paracono” (Figura 4.12c). Estas crestas pueden estar presentes a la vez, formando un “ectolofos doble” (Figura 4.12c) o aparecer solamente una de ellas, formando un “ectolofos simple” (Figura 4.12b). Otro aspecto variable del ectolofos es que, en ocasiones, los ectolofos se encuentran tan desarrollados que se conectan con la cúspide hacia la que están dirigidos si son simples o entre ellos cuando son dobles; en estos casos se habla de “ectolofos completo o conectado”. Por ejemplo, la figura 4.12b sería un ectolofos anterior simple, compuesto por el ectolofos posterior del anterocono y no completo, mientras que la figura 4.12c sería un ectolofos anterior doble completo, compuesto por el ectolofos posterior del anterocono y el ectolofos anterior del paracono.



FIGURA 4.12. Tipos de ectolofos anterior

Ectolofos posterior/*posterior ectoloph*: esta estructura presenta las mismas características que el ectolofos anterior, aunque está relacionada con el paracono y el metacono. De esta manera, se podrá describir un “ectolofos posterior del paracono” (Figura 4.13b) y un “ectolofos anterior del metacono” (Figura 4.13c). Estas crestas podrán aparecer a la vez, llamándose “doble” (Figura 4.13c), o independientemente la una de la otra, recibiendo el nombre de “simple” (Figura 4.13b). De igual modo que en el ectolofos anterior, el ectolofos posterior puede formar una cresta continua que une labialmente el paracono y el metacono (Figura 4.13c) llamándose “ectolofos completo o conectado”, independientemente de que sea “simple” (Figura 4.13b) o “doble” (Figura 4.13c).




FIGURA 4.13. Tipos de ectolofos posterior

Mesolofos/*mesoloph*: en el presente trabajo se considera el mesolofos según lo figuran Mein y Freudenthal (1971b), pero es necesario aclarar los distintos estados de carácter que se pueden mencionar a lo largo del texto. La forma menos desarrollada de mesolofos se ha descrito como un “engrosamiento incipiente/*enlargement of the enamel*” (Figura 4.14); aquí se han incluido todos aquellos mesolofos que se muestran como un abombamiento de la cresta longitudinal. En segundo lugar, el estado “corto/*short*” se refiere a todos los mesolofos que se reconocen como una cresta corta (Figura 4.14). En el siguiente estado, “medio/*medium length*”, el mesolofos recorre más de la mitad de la longitud del mesoseno. Por último el mesolofos “largo/*long*” es únicamente aquel que llega hasta el borde labial del mesoseno.





FIGURA 4.14. Tipos de mesolofos. Sólo se figuran los dos estados de carácter más difíciles de distinguir: “corto” y “engrosamiento incipiente”.

Entomesolofos/*entomesoloph*: se puede considerar como la estructura opuesta al mesolofos, es decir, tiene un desarrollo similar, aunque recorre el seno, en lugar del mesoseno. Cuando está presente, es “corto” cuando no toca el borde lingual y “largo” cuando lo hace.

M2



Anteroseno


FIGURA 4.15. Anteroseno lingual

El segundo molar presenta prácticamente las mismas características que el primer molar. Únicamente cabe destacar el poco desarrollo o ausencia del anterocono en el M2 y el desarrollo de un gran anterolofos labial desde el cual parte la estructura equivalente al “ectolofos posterior del anterocono” del M1. Otro rasgo característico del M2 es el desarrollo de un anteroseno en la pared anterolingual del protocono (Figura 4.15) que tienen algunas especies.

M3

El tercer molar superior presenta una morfología peculiar debido a la reducción de la zona posterior del diente.

Neo-entolofa/*neo-entoloph*: esta cresta, descrita por Freudenthal y Daams (1988), se sitúa entre el hipocono y el protocono y las une transversalmente (Figura 4.16). En las especies más antiguas de Cricetodontini, se han descrito varios estados de carácter en función de la presencia de dos espinas asociadas a cada una de las cúspides o de una estructura continua.



Neo-entolofa

FIGURA 4.16. Neo-entolofa en el M3

Molares inferiores—Los molares inferiores son muy homogéneos. Los caracteres morfológicos que comparten son los siguientes.

Anterolofido labial/*labial anterolophid*: es una cresta que parte desde la parte anterior del diente. En el caso concreto del m1 parte desde el anterocónido. En la gran mayoría de las especies, el anterolofido está presente y bien desarrollado, uniéndose al protocónido, mientras que en las especies más modernas esta cresta se reduce, deja de estar conectada con el protocónido o incluso desaparece (Figura 4.17a).

Anterolofido lingual/*lingual anterolophid*: esta cresta aparece en muy pocos ejemplares y solamente en algunas especies. Nunca alcanza el desarrollo del anterolofido labial; cuando se




FIGURA 4.17. (a). Desarrollo del anterolofido labial en los molares inferiores; (b). Ectomesolofido en los molares inferiores; (c). Espinas del entocónido y metacónido; (d). Tipos de metalofúlido en el m1 inferior.

describe como “presente” se trata siempre de una cresta muy corta y delgada, aunque visible y distinguible.

Mesolófido/*Mesolophid*: es la misma estructura que figuran Mein y Freudenthal (1971b). Respecto a los estados de carácter, se aplican los mismos criterios que para los molares superiores.

Ectomesolófido/*ectomesolophid*: presenta los mismos estados de carácter que en los molares superiores (Figura 4.17b). Esta característica es muy importante en algunas especies, como se explica en el capítulo 7.

Espinas del entocónido y del metacónido/*spurs of the entoconid and metaconid*: Como se muestra en la figura 4.17c, son un “pinzamiento” en el esmalte de la pared posterior de las cúspides linguales. Este “pinzamiento” se aprecia incluso en dientes con un grado moderado de desgaste.

Posterosénido/*posterosenid*: este valle situado entre el posterolófido y el entocónido se encuentra en todas las especies de Cricetodontini. Presenta tres estados de carácter diferentes: puede estar abierto; puede estar cerrado por un cíngulo, es decir, una cresta situada en el borde lingual que conecta el posterolófido con el entocónido; o puede estar cerrado porque el posterolófido sea tan largo que alcance al entocónido. Este carácter es importante en las especies de *Hispanomys*, como se detalla en los apéndices del capítulo 8.

m1

El primer molar presenta caracteres particulares. A continuación, se describe cada uno de ellos.

Anterocónido/*anteroconid*: esta cúspide presenta dos características principales: su situación respecto al eje longitudinal y su forma. En algunas especies de Cricetodontini, el anterocónido está situado en el plano del eje longitudinal, mientras que en otras, está desplazado hacia el borde labial. Por otro lado, suele tener forma redondeada en la mayoría de las especies de Cricetodontini, mientras que en unos pocos taxones el anterocónido es alargado.

Metalofúlido/*metalophulid*: es la cresta de esmalte que, según su orientación, une el metacónido con el anterolofúlido o con el brazo posterior del protocónido. Por lo tanto, hay dos estructuras diferentes que están asociadas al metacónido. Mein y Freudenthal (1971b) las diferencian como “anterior” y “posterior”, asignando el estado de carácter “doble” cuando ambas están presentes en el mismo espécimen. Sin embargo, en este trabajo se sigue la nomenclatura de Rummel y Kälin (2003) que diferencian las estructuras independientemente como “metalofúlido I” (Figura 4.17d), aquel que une el metacónido con el anterolofúlido, y “metalofúlido II”, aquel que une el metacónido con el brazo posterior del protocónido (Figura 4.17d). En adelante, se hablará de “metalofúlido I+II/*metalophulid I+II*” cuando lo que se quiere destacar es que ambos metalo-

fúlicos están presentes en el mismo diente (Figura 4.17d).

m2

Número de raíces/*number of roots*: el segundo molar puede presentar dos raíces (Fig. 4.18a) o tres. En el lado anterior siempre tiene una raíz, mientras que en el lado posterior, la raíz puede dividirse. Cuando la raíz posterior está dividida se describe que el m2 tiene tres raíces. Si la separación en dos de la raíz posterior no llega hasta la mitad de la longitud de la raíz, se dice que presenta “raíz posterior parcialmente dividida” (Fig. 4.18b). Se dice que la “raíz posterior está



FIGURA 4.18. Tipos de divisiones de la raíz posterior del m2. (a) Sin dividir. (b). Parcialmente dividida. (c). Totalmente dividida.

totalmente dividida” cuando la separación sobrepasa la mitad de la longitud de la raíz o incluso llega hasta el borde de la corona dental (Fig. 4.18c).

Caracteres métricos

La talla de los molares de los cricétidos, como la de todos los roedores, se caracteriza con las variables anchura y longitud. El criterio seguido en la toma de medidas (según Daams y Freudenthal, 1988) representa la longitud y la anchura máximas de la corona dental tomadas de forma perpendicular una a la otra (Figura 4.19a). Para orientar cada ejemplar se ha utilizado como referencia la faceta de contacto entre los distintos dientes en la hilera dental. No se han medido los ejemplares rotos, muy desgastados o digeridos. Las variables métricas de cada diente se ha tomado utilizando una lupa monocular *Nikon Measurescope 10* (Figura 4.19b) provista de micrómetro digital en el material conservado en los fondos del MNCN-CSIC y de la UCM. En el caso del material conservado en NBC-Leiden o en IAU-Utrecht, las medidas se han tomado sobre fotografías realizadas con una cámara acoplada a una lupa Leica y su *software* asociado.

En primer lugar, tras la toma de datos, se ha realizado el cálculo de estadísticos descriptivos; los valores máximo, mínimo y media aritmética de las dos variables estudiadas. Cuando la muestra

supera los cinco ejemplares, se ha calculado la desviación estándar para informar del grado de dispersión que la muestra presenta. Todos estos datos se presentan a modo de tablas en cada uno de los capítulos del apartado (Paleontología Sistemática), agrupados por localidades y por especies. También se han calculado índices para conocer las proporciones de determinados elementos, o de la hilera dental. El índice, Longitud/Anchura del M3 ha sido muy útil en la caracterización de algunas especies, como se explica en el capítulo 5. Los índices Longitud m1/Longitud m3 y Longitud M1/Longitud M3 han sido tradicionalmente utilizados para distinguir unas especies de otras (Freudenthal, 1966; Rummel, 1995; Kálin y Kempf, 2009; López-Antoñanzas et al., 2010). Como se detalla en el apartado (Paleontología sistemática) han sido muy útiles en esta tesis doctoral, sobre todo en el capítulo 8.

Se han realizado análisis estadísticos con las variables métricas de las asociaciones estudiadas con dos propósitos. Por un lado, para determinar si existen diferencias significativas entre las diferentes especies y por otro, para determinar si dentro de una misma especie existen variaciones de talla entre los fósiles de distintas localidades. En los apartados metodológicos de cada uno de los capítulos del apartado (Paleontología Sistemática) se detalla qué prueba estadística se ha hecho y con qué material, por lo que a continuación se resumen, de manera muy general, los análisis que se pueden encontrar a lo largo del texto.

Para comparar dos grupos de datos entre sí, se ha elegido el test t de Student, que evalúa si dos muestras univariantes se han recogido de poblaciones con medias iguales. Para hacer esta prueba se deben cumplir tres supuestos: los datos deben ser continuos, ajustarse a distribuciones normales y tener varianzas homogéneas. Para saber si estos dos últimos supuestos se cumplen, se han realizado el test de Shapiro-Wilk y el test de Levene. Otro tipo de prueba que se ha utilizado ha sido el análisis de la varianza o ANOVA. Se utiliza para saber si varias muestras univariantes provienen de poblaciones con medias iguales o no. Los supuestos son los mismos que para la prueba t de Student, por lo que los test previos (Shapiro-Wilk y Levene) también se han realizado. En todos los tipos de pruebas, se han establecido dos umbrales para el nivel de signifi-




FIGURA 4.19. (a). Ilustración del método de toma de medidas de los molares de Cricetodontini. (b). Lupa monocular del MNCN-CSIC provista de micrómetro digital utilizado para tomar medidas.

cación, por debajo de los cuales se rechaza la hipótesis nula (que varía según el tipo de prueba). Si $p < 0,05$ se considera que hay diferencias significativas mientras que cuando $p < 0,01$, las diferencias son muy significativas.

MÉTODO DE ESTUDIO FILOGENÉTICO

Para conocer las relaciones filogenéticas de las especies de Calatayud-Daroca, se ha de tener en cuenta que éstas no forman parte de un grupo aislado y que es posible que su pariente más cercano fuera un taxón foráneo de la cuenca. Por eso, se han estudiado 44 especies más de la tribu Cricetodontini, haciendo un total de 52 especies. Además del material tipo de la especie encontradas en la Cuenca de Calatayud-Daroca, se ha observado personalmente el material de 12 poblaciones tipo (Tabla 4.2), se ha estudiado la información de publicaciones recientes que cuentan con descripciones detalladas y fotografías. En otros casos, la información se ha obtenido de publicaciones antiguas o con descripciones menos exhaustivas, por lo que la calidad de las observaciones en estas especies es menor, lo que se ha indicado convenientemente.

Los aspectos metodológicos más concretos se detallan en el capítulo 10, a continuación se describen los procedimientos generales para realizar el análisis.

En primer lugar se han elegido qué especies formarán parte de nuestro estudio. Se han seleccionado los caracteres que evidenciarán sus relaciones filogenéticas. Se han establecido las mismas y se han expresado en un cladograma. Se ha utilizado éste para corroborar o refutar hipótesis expuestas en los capítulos del apartado III (Paleontología Sistemática) o por otros autores.

El método utilizado para construir los cladogramas es el de máxima parsimonia. Este procedimiento está guiado por el principio metodológico de la parsimonia o simplicidad (Crisci, 1982; Farris, 1982; Kluge, 2001; Goloboff, 2003), el cual asume que entre hipótesis alternativas hay que elegir la que requiera un menor número de homoplasias (Morrone, 2000). El procedimiento usa el algoritmo de Wagner para construir un cladograma con un número grande de taxones y caracteres (Farris, 1970). Aunque en la práctica para más de 15-20 taxones es imposible encontrar el cladograma más parsimonioso usando este método, es un buen punto de partida para el método que se ha utilizado, la “búsqueda por permutación de ramas” (Goloboff, 1998). Después de obtener los cladogramas más parsimoniosos, se ha calculado el cladograma de consenso estricto, aquel que resulta de combinar únicamente aquellos componentes que aparezcan repetidos en todos los cladogramas originales (Nelson, 1979; Sokal y Rohlf, 1981).

La prueba estadística que se ha utilizado para determinar la confianza en los resultados del análisis cladístico ha sido el llamado soporte de ramas. Este método, propuesto por Bremer

(1994), examina cuántos pasos extra se requieren para que se pierda una rama en el cladograma de consenso. El soporte de las ramas se calcula forzando búsquedas heurísticas en las que los árboles guardados tengan cada vez un número de pasos mayor que el de los árboles más parsimoniales previamente encontrados. Así, los clados que habiendo aparecido en la búsqueda inicial desaparecen al aumentar en un paso la longitud final del árbol, se consideran clados poco apoyados y se determinarían por un valor de 1 (Morrone, 2000).

En los siguientes capítulos de los apartados III (Paleontología sistemática), IV (Bioestratigrafía) y V (Filogenia) se detallan los resultados y conclusiones al aplicar todos estos métodos en el material estudiado.

Bibliografía

de los capítulos 1-4



BIBLIOGRAFÍA (CAPÍTULOS 1-4)

- Abels, H. A., F. J. Hilgen, W. Krijgsman, R. W. Kruk, I. Raffi, E. Turco y W. J. Zachariasse. 2005. Long-period orbital control on middle Miocene global cooling: Integrated stratigraphy and astronomical tuning of the Blue Clay Formation on Malta. *Paleoceanography* 20:PA4012.
- Aguilar, J. P., M. Calvet y J. Michaux. 1994. Les rongeurs de Castelnou 6 (Pyrénées-orientales, France) et les corrélations entre faunes ibériques et françaises au Miocène moyen. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 192:109–131.
- Aguilar, J. P., V. Lazzari, J. Michaux, M. Sabatier y M. Calvet. 2007. Lo Fournas 16-M (Miocène supérieur) et Lo Fournas 16-P (Pliocène moyen), nouvelles localités karstiques à Baixas, Sud de la France. Partie I: Description et implications géodynamiques. *Géologie de la France* 1:55–62.
- Agustí, J. 1977. Descripción de *Cricetodon lavocati* Freudenthal 1966, de Sant Quirze (Trinxera). *Paleontologia i Evolució* 12:65–67.
- Agustí, J. 1980. La asociación de *Hispanomys* y *Cricetodon* (Rodentia, Mammalia) en el Mioceno superior del Vallès-Penedès (Cataluña, España). *Acta Geologica Hispanica* 15:51–60.
- Agustí, J. 1981. Roedores miomorfos del Neógeno de Cataluña. Tesis Doctoral, Universidad de Barcelona, Barcelona, 293 pp.
- Agustí, J. 1982. Tendencias evolutivas de la línea *Cricetodon-Ruscinomys* (Rodentia, Mammalia) en la Península Ibérica. *Acta Geologica Hispanica* 17:103–111.
- Agustí, J. y J. Gibert. 1982. Roedores e insectívoros del Mioceno superior de Hostalets de Pierola (Vallès-Penedès, Cataluña). Institut de Paleontologia de Sabadell. *Butlletí Informatiu* 14:19–38.
- Agustí, J. y S. Moyá-Solá. 1990. Mammal extinctions in the Vallesian (Upper Miocene). *Lecture Notes in Earth Science* 30:425–432.
- Agustí, J., I. Casanovas-Vilar y M. Furió. 2005. Rodents, insectivores and chiropterans (Mammalia) from the late Aragonian of Can Missert (middle Miocene, Vallès-Penedès Basin, Spain). *Geobios* 38:575–583.

- Anadón, P. y E. Moissenet. 1996. Neogene basins in the Eastern Iberian Range; pp. 68–76 en P. F. Friends y C. J. Dabrio (eds.), *Tertiary basins of Spain. The stratigraphic record of crustal kinematics*. World and Regional Geology 6, Cambridge University Press, Cambridge.
- Alba, D.M., S. Moyá-Solá, I. Casanovas-Vilar, J. Galindo, J. M. Robles, C. Rotgers, M. Furió, C. Angelone, M. Kohler, M. Garcés, L. Cabrera, S. Almécija y P. Obrado. 2006. Los vertebrados fosiles del Abocador de Can Mata (els Hostalets de Pierola, l’Anoia, Cataluña), una sucesión de localidades del Aragoniense superior (MN6 y MN7+8) de la cuenca del Vallès-Penedès. Campaños 2002-2003, 2004 y 2005. *Estudios Geológicos* 62:295–312.
- Alcalá, L., A. M. Alonso-Zarza, M. A. Álvarez-Sierra, B. Azanza, J. P. Calvo, J. C. Cañaveras, J. Van Dam, M. Garcés, W. Krijgsman, A. J. Van der Meulen, P. Pélaez-Campomanes, A. Pérez-González, S. Sánchez Moral, R. Sancho y E. Sanz Rubio. 2000. El registro sedimentario y faunístico de las cuencas de Calatayud-Daroca y Teruel: evolución paleoambiental y paleoclimática durante el Neógeno. *Revista de la Sociedad Geológica de España* 13:323–343.
- Alonso-Zarza, A. M. y J. P. Calvo. 2000. Palustrine sedimentation in an episodically subsiding basin: the Miocene of the northern Teruel Graben. *Palaeogeography, Palaeoclimatology, Palaeoecology* 160:1–21.
- Alonso-Zarza, A. M. y J. P. Calvo. 2002. Tajo Basin; pp. 315–320 en W. Gibbons, y T. Moreno (eds.), *The Geology of Spain*. The Geological Society, London.
- Alonso Zarza, A. M., J. P. Calvo y M. A. García del Cura. 1992. Palustrine sedimentation and associated features—grainification and pseudo-microkarst—in the Middle Miocene (Intermediate Unit) of the Madrid Basin, Spain. *Sedimentary Geology* 76:43–61.
- Álvarez-Sierra, M. A. 1983. Paleontología y bioestratigrafía del Mioceno superior del sector de la Cuenca del Duero. Estudio de los micromamíferos de la serie de Torremormojón (Palencia). Memoria de grado de licenciatura. Universidad Complutense de Madrid, Madrid, 188 pp.
- Álvarez-Sierra, M. A. 1987. Estudio sistemático y bioestratigráfico de los Eomyidae (Rodentia) del Oligoceno superior y Mioceno inferior español. *Scripta Geologica* 86:1–207.
- Álvarez-Sierra, M. A. y E. García Moreno. 1986. New Gliridae and Cricetidae (Mammalia Rodentia) from the Middle and Upper Miocene of the Duero basin, Spain. *Studia geologica salmanticensia* 22:145–189.
- Álvarez-Sierra, M. A., E. García-Moreno, N. López Martínez y R. Daams. 1987. Biostratigraphy and Paleoecological interpretation of Micromammal successions in the Upper Arago-

- nian and Vallesian (Middle-Upper Miocene) of the Duero Basin (N.Spain). *Annales de l'Institut Géologique de Hongrie* 70:517–522.
- Álvarez-Sierra, M. A., I. García-Paredes, V. Hernández-Ballarín, L.W. Van den Hoek Ostende, K. Hordijk, P. López-Guerrero, A.J. Van der Meulen, A. Oliver y P. Peláez-Campomanes. 2013. Models of historical biogeography and continental biochronology. *Spanish journal of Palaeontology* 28:129–138.
- Álvarez Sierra, M. A., J. Civis, A. Corrochano, R. Daams, C. J. Dabrio, E. García, A. González, N. López-Martínez, R. Mediavilla, R. Rivas Carballo y M. F. Valle. 1990. Un estratotipo del límite Aragoniense-Vallesiense (Mioceno medio-Mioceno superior) en la sección de Torremormojón (Cuenca del Duero, Provincia de Palencia). *Studia Geologica Salmanticensia, Actas de Paleontología*:57–64.
- Álvarez-Sierra, M. A., J. P. Calvo, J. Morales, A. M. Alonso-Zarza, B. Azanza, I. García Paredes, M. Hernández Fernández, A. J. Van der Meulen, P. Peláez-Campomanes, V. Quiralte, M. J. Salesa, I. M. Sánchez y D. Soria. 2003. El tránsito Aragoniense-Vallesiense en el área de Daroca-Nombrevilla (Zaragoza, España). *Coloquios de Paleontología Volumen Extraordinario* 1:25–33.
- Azanza, B., M. A. Alonso Zarza, M. A. Álvarez Sierra, J. P. Calvo, S. Fraile, I. García Paredes, E. Gómez, M. Hernández Fernández, A. J. Van der Meulen, D. de Miguel, P. Montoya, J. Morales, X. Murelaga, P. Peláez-Campomanes, B. Perez, V. Quiralte, M. J. Salesa, A. Sánchez Marco y D. Soria. 2004. Los yacimientos de vertebrados continentales del Aragoniense superior (Mioceno medio) de Toril, Cuenca de Calatayud-Daroca. *Geo-Temas* 6:271–274.
- Bolliger, T. V. 1994. Die Obere Süßwassermolasse in Bayern und der Ostschweiz: bio-und lithostratigraphische Korrelationen. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und histor. Geologie* 34:109–144.
- Böhme, M., J. Prieto, B. Reichenbacher, M. Aiglstorfer y A. Rocholl. 2012. Pre-Riesian hiatus (Styrian tectonic phase) and minor hiatuses in the NAFB; pp. 2–8 en M. Böhme, J. Prieto, B. Reichenbacher, M. Aiglstorfer y A. Rocholl (eds.), *The Early Miocene Molasse in southern Germany. Field Trip Guide. RCMNS Workshop May 16th–20th, 2012. Tübingen-Bad Krumbad*.
- Boon, E. 1991. Cricetiden und Sciuriden der Oberen Süßwassermolasse. Tesis Doctoral, Fakultät für Geowissenschaften der Ludwig-Maximilians-Universität, Munich, 159 pp.
- Bremer, K. 1994. Branch support and tree stability. *Cladistics* 10:295–304.

- Calvo, J. P., R. Daams, J. Morales, N. López Martínez, J. Agustí, P. Anadón, I. Armenteros, L. Cabrera, J. Civis, A. Corrochano, M. Díaz Molina, E. Elizaga, M. Hoyos, E. Martín-Suárez, J. Martínez, E. Moissenet, A. Muñoz, A. Pérez García, A. Pérez González, J. M. Portero, F. Robles, C. Santisteban, T. Torres, A. J. Van der Meulen, J. A. Vera y P. Mein. 1993. Up-to-date Spanish continental Neogene synthesis and paleoclimatic interpretation. *Revista de la Sociedad Geológica de España* 6:29–40.
- Casanovas-Vilar, I. 2007. The rodent assemblages from the Late Aragonian and the Vallesian of the Vallès-Penedès Basin (Catalonia, Spain), Tesis Doctoral, Universidad Autònoma de Barcelona, Barcelona, 286 pp.
- Casanovas-Vilar, I., I. García-Paredes y J. Van Dam. 2013. The Vallesian Rodent Record from the Vallès-Penedès Basin (Catalonia, Spain): Biostratigraphy, Diversity and Structure; pp. 132 en N. Çağatay, and C. Zabcı (eds.), 14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratethys and Black sea. Book of Abstracts. General Directorate of Mineral Research and Exploration (MTA), Istanbul.
- Casanovas-Vilar, I., S. Moyà-Solà, J. Agustí y M. Köhler. 2005. The geography of a faunal turnover: tracking the Vallesian Crisis; pp. 247–301 en A. Elewa (ed.), *Migration in organisms: climatology, geography, ecology*. Springer-Verlag Publishers, Heidelberg.
- Casanovas-Vilar, I., L. W. Van den Hoek Ostende, M. Furió y A. P. Madder. 2014. The range and extent of the Vallesian Crisis (Late Miocene): new prospects based on the micromammal record from the Vallès-Penedès basin (Catalonia, Spain). *Journal of Iberian Geology* 40:29–48.
- Casas, A. N., A. Casas, A. Pérez, S. Tena, L. Berrier, D. Garpais y T. Nalpas. 2003. Syn-tectonic sedimentation and thrust-and-fold kinematics at the intra-mountain Montalbán Basin (northern Iberian Chain, Spain). *Geodinámica Acta* 13:1–17.
- Colomer, M. y P. Santanach. 1988. Estructura y evolución del borde sur-occidental de la Fosa de Calatayud-Daroca. *Geogaceta* 4:29–32.
- Crisci, J. V. 1982. Parsimony in Evolutionary Theory Law or methodological prescription. *Journal of Theoretical Biology* 97:35–41.
- Crusafont, M. 1950. La cuestión del llamado Meótico español. *Arrahona* 1:3–9.
- Cuenca Bescós, G. 1988. Revisión de los Sciuridae del Aragoniense y del Ramblense en la fosa de Calatayud-Montalbán. *Scripta Geologica* 87:1–116.
- Daams, R. 1974. *Pseudodryomys* (Gliridae, Rodentia, Mammalia) from Early and Middle Miocene deposits in Spain. *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen*

- schappen, Series B 77:201–240.
- Daams, R. 1981. The dental pattern of the Dormice *Dryomys*, *Myomimus*, *Microdyromys* and *Peridyromys*. Utrecht micropaleontological bulletins. Special publication. Utrecht 3:1–113.
- Daams, R. 1990. Hypsodont Myomiminae (Gliridae, Rodentia) from the Miocene and the Oligocene-Miocene boundary interval of Spain. Scripta Geologica 95:1–62.
- Daams, R. 1999. *Peridyromys darocensis* and *Peridyromys sondaari*, two new species of Gliridae (Rodentia, Mammalia) from the Lower Miocene (MN3-5) of the Calatayud-Daroca Basin, Zaragoza, Spain. Deinsea 7:83–90.
- Daams, R. y M. Freudenthal. 1981. Aragonian: the stage concept versus Neogene mammal zones. Scripta Geologica 62:1–17.
- Daams, R. y M. Freudenthal. 1988. Synopsis of the Dutch-Spanish collaboration program in the Neogene of the Calatayud-Teruel basin. Scripta Geologica Special Issue 1:3–18.
- Daams, R., M. Freudenthal y M. A. Álvarez-Sierra. 1987. Ramblian: a new stage for continental deposits of early Miocene age. Geologie en Minjbouw 65:297–308.
- Daams, R., M. Freudenthal y A. Van de Weerd. 1977. Aragonian, a new stage for continental deposits of Miocene age. Newsletters on Stratigraphy 6:42–55.
- Daams, R., A. J. Van der Meulen, P. Peláez-Campomanes y M. A. Álvarez-Sierra. 1999a. Trends in rodent assemblages from the Aragonian (Early Middle Miocene) of the Calatayud-Daroca Basin (Aragón, Spain); pp. 127–139 en J. Agustí, L. Rook, and P. Andrews (eds.), Hominoid Evolution and Climate Change in Europe. 1 The Evolution of Neogene Terrestrial Ecosystems in Europe. Cambridge University Press, Cambridge.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes y W. Krijgsman. 1999b. Aragonian stratigraphy reconsidered and a re-evaluation of the middle Miocene mammal biochronology in Europe. Earth and Planetary Science Letters 165:287–294.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza y W. Krijgsman. 1999c. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North- Spain). Newsletters on Stratigraphy 37:103–139.
- Daxner-Höck, G. 2003. *Cricetodon meini* and other rodents from Mühlbach and Grund, Lower Austria (Middle Miocene, late MN5). Annalen des Naturhistorischen Museums in Wien 104A:267–291.
- De Bruijn, H. y P. Mein. 1968. On the mammalian fauna of the *Hipparion* beds in the Calatayud-

- Teruel basin (province of Zaragoza, Spain). Part V: The Sciurinae. Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen, Series B 71:73–90.
- De Bruijn, H. y E. Ünay. 1996. On the evolutionary history of the Cricetodontini from Europe and Asia Minor and its bearing on the reconstruction of migrations and the continental biotope during the Neogene; pp. 227–234 en R. L. Bernor, V. Fahlbusch, H. W. Mittmann (eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- De Bruijn, H., E. Ünay y K. Hordijk. 2013. A review of the Neogene Succession of the Muridae and Dipodidae from Anatolia, with special reference to taxa Known from Asia and/or Europe; pp. 566–582 en X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Sarac y E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a Discussion of the Evolutionary History of the Cricetodontini. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 96:151–216.
- De Jong, F. 1988. Insectivora from the Upper Aragonian and the Lower Vallesian of the Daroca-Villafeliche area in the Calatayud-Teruel basin (Spain). *Scripta geológica, Special Issue* 1:253–285.
- Domingo, L., P. L. Koch, S. T. Grimes, J. Morales y N. López-Martínez. 2012. Isotopic paleoecology of mammals and the Middle Miocene Cooling event in the Madrid Basin (Spain). *Palaeogeography Palaeoclimatology Palaeoecology* 339:98–113.
- Farris, J. S. 1970. Methods for computing Wagner Trees. *Systematic Zoology* 19:83–92.
- Farris, J. S. 1982. Outgroups and parsimony. *Systematic Zoology* 31:328–334.
- Ferreiro, E. y V. Ruiz. 1991. Memoria y Mapa geológico 1:200.000, Daroca (Hoja nº 40). Instituto Tecnológico Geominero. España, Madrid.
- Flynn, L. J., L. L. Jacobs y E. H. Lindsay. 1985. Problems in muroid phylogeny. Relationship to other rodents and origin of major groups; pp. 589–616 en P. E. A. Lockett (ed.), *Evolutionary relationships among rodents. A multidisciplinary analysis*. NATO ASI Serie A: Life Sciences.
- Flynn, L. J., M. E. Morgan, D. Pilbeam y J. C. Barry. 2014. “Endemism” relative to space, time, and taxonomic level. *Annales Zoologici Fennici* 51:245–258.
- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mittel-

- spaniens und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) Ia. Genera *Cricetodon* and *Ruscinomys* (Rodentia). Koninklijke Nederlandse Akademie Van Wetenschappen-Proceedings Series B-Physical Sciences 69:298–317.
- Freudenthal, M. y R. Daams. 1988. Cricetidae (Rodentia) from the type-Aragonian; the genera *Democricetodon*, *Fahlbuschia*, *Pseudofahlbuschia* nov. gen., and *Renzimys*; pp. 133–252 en M. Freudenthal (ed.), Biostratigraphy and paleoecology of the Neogene micromammalian faunas from the Calatayud-Teruel Basin (Spain). Scripta Geologica, Special Issue, Leiden.
- Freudenthal, M. y E. Martín Suárez. 1999. Family Muridae; pp. 401–409 en G. Rößner and K. Heißig (eds.), The Miocene land mammals of Europe. Verlag Dr. Friedrich Pfeil, München.
- García-Alix, A., 2006. Bioestratigrafía de los depósitos continentales de la transición Mio-Plioceno de la cuenca de Granada. Tesis Doctoral, Universidad de Granada, Granada, 429 pp.
- Garcés, M., J. Agustí, L. Cabrera y J. M. Pares. 1996. Magnetostratigraphy of the Vallesian (late Miocene) in the Vallès-Penedès Basin (northeast Spain). *Earth and Planetary Science Letters* 142:381–3986.
- Garcés, M., W. Krijgsman, P. Peláez-Campomanes, M. A. Álvarez Sierra y R. Daams. 2003. *Hipparion* dispersal in Europe: magnetostratigraphic constraints from the Daroca area (Spain). *Coloquios de Paleontología Volumen Extraordinario* 1:171–178.
- García-Moreno, E. 1987. Roedores y Lagomorphos del Mioceno de la Zona de la Cuenca del Duero. Sistemática, Bioestratigrafía y Paleoecología. Tesis Doctoral, Universidad Complutense de Madrid, Madrid, 258 pp.
- García Paredes, I. 2006. Patrones evolutivos de los Gliridae (Rodentia, Mammalia) del Mioceno inferior y medio del área tipo del Aragoniense (Cuenca de Calatayud-Montalbán). Tesis Doctoral, Universidad Complutense de Madrid, Madrid, 676 pp.
- García-Paredes, I. e I. Casanovas-Vilar. 2013. Vallesian Gliridae from the Valles-Penedes Basin (Catalonia, Spain) and how were they Affected by the ‘Vallesian Crisis’; pp. 127 en N. Çagatay, and C. Zabcı (eds.), 14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratethys and Black sea. Book of Abstracts. General Directorate of Mineral Research and Exploration (MTA), Istanbul.
- García-Paredes, I., P. Peláez-Campomanes y M. A. Álvarez-Sierra. 2009. Gliridae (Rodentia, Mammalia) with a simple dental pattern: a new genus and new species from the European

- Early and Middle Miocene. *Zoological Journal of the Linnean Society* 157:622–652.
- García-Paredes, I., P. Peláez-Campomanes y M. A. Álvarez-Sierra. 2010. *Microdyromis remmertii*, sp. nov., a new Gliridae (Rodentia, Mammalia) from the Aragonian type area (Miocene, Calatayud-Montalbán Basin, Spain). *Journal of Vertebrate Paleontology* 30:1594–1609.
- Goloboff, P. A. 1998. Tree searches under Sankoff parsimony. *Cladistics* 14:229–237.
- Goloboff, P. A. 2003. Parsimony, likelihood and simplicity. *Cladistics* 19:91–103.
- Gomes Rodrigues, H., L. Marivaux y M. Vianey-Liaud. 2009. Phylogeny and systematic revision of Eocene Cricetidae (Rodentia, Mammalia) from and East Asia: on the origin of cricetid rodents. *Journal of Zoological Systematics and Evolutionary Research* 48:259–268.
- Hartenberger, J. L. 1965. Les Cricetidae (Rod.) de Can Llobateres (Néogène d'Espagne). *Bulletin De La Societe Geologique De France* 7:487–498.
- Hartenberger, J. L. 1998. Description of the radiation of the Rodentia (Mammalia) from the Late Paleocene to the Miocene; phylogenetic consequences. *Comptes Rendus De L Academie Des Sciences Serie Ii Fascicule a-Sciences De La Terre Et Des Planetes* 326:439–444.
- Heißig, K., 1997. Mammal faunas intermediate between the reference faunas of MN4 and MN6 from the Upper freshwater Molasse of Bavaria; pp. 537–546 en J. P., Aguilar, S. Legendre, and J. Michaux (eds.) *Congrès BiochroM '97. Mémoires et Travaux EPHE*, Institut Montpellier, Montpellier.
- Heißig, K., 2006. Biostratigraphy of the “main bentonite horizon” of the freshwater Molasse in Bavaria. *Paleontographica* A277:93–102
- Héran, M. A., C. Lécuyer y S. Legendre. 2010. Cenozoic long-term terrestrial climatic evolution in Germany tracked by $\delta^{18}\text{O}$ of rodent tooth phosphate. *Palaeogeography, Palaeoclimatology, Palaeoecology* 285:331–342.
- Hernández-Ballarín, V., A. Oliver y P. Peláez-Campomanes. 2011. Revisión de las asociaciones de mamíferos del tránsito Aragoniense medio y superior de la Cuenca de Madrid; pp. 183–172 en A. Pérez-García, F. Gascó, J. M. Gasulla, y F. Escaso (eds.), *Viajando a Mundos Pretéritos*. Ayuntamiento de Morella, Morella, Castellón.
- Hernández-Ballarín, V., A. Oliver, I. García-Paredes y P. Peláez-Campomanes. 2010. Estudio preliminar de la fauna de roedores del yacimiento mioceno de El Cañaveral (Madrid, Espana). *Cidaris* 30:151–153.
- Hernández Fernández, M., J. A. Cárdbaba, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López Guerrero, N. Sala-Burgos, J. Morales y N. López

- Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Herráez, E. 1993. Micromamíferos (Roedores y Lagomorfos) del Mioceno del área de Madrid: Estudio sistemático y bioestratigráfico. Tesis Doctoral, Universidad Complutense de Madrid, Madrid. 338 pp.
- Hershkovitz, P. 1962. Evolution of Neotropical Cricetine rodents (Muridae) with special reference to the Phyllotine group. *Fieldiana: Zoology* 46:1–524.
- Hilgen, F. J., L. J. Lourens y J. A. Van Dam. 2012. The Neogene Period; pp. 923–978 en F. M. Gradstein, J. G. Ogg, J. G. Schmitz (eds.), *Geologic Time Scale*. Cambridge University Press, Cambridge.
- Hordijk, K. 2010. Perseverance of pikas in the Miocene. Tesis Doctoral. Medelingen van de Faculteit Geowetenschappen Universiteit Utrecht, Utrecht, 312 pp.
- Hordijk, K. y A. Van der Meulen. 2007. Presence and abundance patterns of Middle Miocene Pikas from North Spain. *Journal of Vertebrate Paleontology* 27:91A–91A.
- Julivert, M. 1954. Observaciones sobre la tectónica de la Depresión de Calatayud. *Arrahona*:3–18.
- Kälin, D. y O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 254:177–235.
- Kennett, J. y P. F. Barker. 1990. Latest Cretaceous to Cenozoic climate and oceanographic developments in the Weddell Sea, Antarctica: an ocean-drilling perspective; pp. 937–960 en P. F. Barker, J. P. Kennett (eds.), *Proceedings ODP Scientific Results*.
- Klein Hofmeijer, G. y H. De Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece) Part 8: The Cricetidae. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 91:185–204.
- Kluge, A., 2001. Philosophical conjectures and their refutation. *Systematic Biology* 50:322–330.
- Krijgsman, W., C. G. Langereis, R. Daams, y A. J. Van der Meulen. 1994. Magnetostratigraphic dating of the middle Miocene climate change in the continental deposits of the Aragonian type area in the Calatayud-Teruel basin (Spain). *Earth and Planetary Science Letters* 128:513–526.
- Krijgsman, W., M. Garcés, C. G. Langereis, R. Daams, J. van Dam, A. J. Van der Meulen, J. Agustí y L. Cabrera. 1996. A new chronology for the middle to late Miocene continental record

- in Spain. *Earth and Planetary Science Letters* 142:367–380.
- Lacomba, J. I. 1988. Rodents and Lagomorphs from a lower Vallesian fissure filling near Molina de Aragón (prov. Guadalajara, Spain); pp. 19–38 en M. Freudenthal (ed.), *Biostratigraphy and paleoecology of the Neogene micromammalian faunas from the Calatayud-Teruel Basin* (Spain). *Scripta Geologica*, Special Issue 1, Leiden.
- Lavocat, R. 1978. *Les relations faunistiques Afrique-Amérique. Aspects modernes des recherches sur l'Evolution*, Montpellier, France, 1978.
- Lear, C. H., H. Elderfield y P. A. Wilson. 2000. Cenozoic deep-sea temperatures and global ice volumes from Mg/Ca in benthic foraminiferal calcite. *Science* 287:269–272.
- López-Antoñanzas, R., P. Peláez-Campomanes, M. A. Álvarez-Sierra e I. García-Paredes. 2010. New species of *Hispanomys* (Rodentia, Cricetodontinae) from the Upper Miocene of Batallones (Madrid, Spain). *Zoological Journal of the Linnean Society* 160:725–747.
- López-Guerrero, P., M. A. Álvarez-Sierra e I. García-Paredes. 2011a. Up to date of the stratigraphical and geographical distribution of *Cricetodon*; pp. 157–158 en F. J. Sierro, y J. A. González Delgado (eds.), *Joint RCMNS-RCANS Interim Colloquium. Climate changes, bioevents and geochronology in the Atlantic and Mediterranean over the last 23 Myr*. Salamanca.
- López-Guerrero, P., I. García-Paredes y M. A. Álvarez-Sierra. 2013a. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes y P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89:573–592.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra y P. Peláez-Campomanes, P. 2013b. Cricetodontini from the Calatayud-Daroca Basin (Miocene, Spain); pp. 240 en N. Çağatay, y C. Zabcı (eds.), *14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratehys and Black sea. Book of Abstracts*. General Directorate of Mineral Research and Exploration (MTA), Istanbul.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra y P. Peláez-Campomanes. En prensa. Cricetodontini from the Calatayud–Daroca Basin (Spain): A taxonomical description and update of their stratigraphical distributions. *Comptes rendus Palevol* DOI:<http://dx.doi.org/10.1016/j.crpv.2014.07.002>.

- López-Guerrero, P., M. Álvarez-Sierra, I. García-Paredes, R. López-Antoñanzas y A. Oliver. 2009. Cricetodontini (Rodentia, Mammalia) from the Upper Aragonian and Lower Vallesian of the Toril-Nombrevilla Section (Middle and Upper Miocene, Calatayud-Daroca Basin, Zaragoza, Spain). *Journal of Vertebrate Paleontology* 29:161A–161A.
- López-Guerrero, P., M. A. Álvarez-Sierra, R. López-Antoñanzas, A. Oliver Pérez y P. Peláez-Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). *Palaeontologica Nova* 8:259–271.
- López-Guerrero, P., A. Oliver Pérez, M. A. Álvarez-Sierra, I. García Paredes, L. Van den Hoek Ostende y P. Peláez-Campomanes de Labra. 2007. Paleontología del yacimiento de vertebrados Aragoniense de Casa Montero (Madrid). *Cantera Paleontológica* 53:247–254.
- López-Guerrero, P., I. García-Paredes, L. W. Van den Hoek Ostende, J. A. Van Dam, M. A. Álvarez-Sierra, V. Hernández-Ballarín, A. J. Van der Meulen, A. Oliver y P. Peláez-Campomanes. 2011b. Cañada: a new micromammal succession from the lower Vallesian and Turolian of the Daroca area (Calatayud-Montalbán Basin, Spain). *Estudios Geológicos* 67:443–453.
- López Martínez, N. 1972. Los micromamíferos del Cuaternario de Rincón de la Victoria. *Boletín Real Sociedad Española Historia Natural (Geología)* 70:223–233.
- López Martínez, N. 1992. Técnicas de estudio de microvertebrados. Los micromamíferos y su interés bioestratigráfico; pp. 345–365 en H. Astibia (ed.), *Paleontología de vertebrados: Faunas y filogenia, aplicación y sociedad*. Servicio Editorial de la Universidad del País Vasco.
- López-Martínez, N., E. García Moreno y M. A. Álvarez-Sierra. 1986. Paleontología y bioestratigrafía (micromamíferos) del Mioceno medio y superior del sector de la Cuenca del Duero. *Studia Geologica Salmanticensia*. Universidad de Salamanca 22:191–212.
- Lourens, L. J., F. J. Hilgen, J. Laskar, N. J. Shackleton y D. Wilson. 2004. The Neogene Period; pp. 409–440 en F. M. Gradstein, J. G. Ogg, y A. G. Smith (eds.), *Geologic Time Scale 2004*. Cambridge University Press, Cambridge.
- Maddern, A., I. Casanovas-Vilar, D. M. Alba, D. DeMiguel, J. M. Robles, L.W. Van den Hoek Ostende y S. Moyá-Solá. 2013. The Abrupt Collapse of a diversity Hotspot? Reconsidering Vallesian (Late Miocene) diversity in its type area. 125 pp. en N. Çağatay, y C. Zabcí (eds.), 14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratehys and Black sea. Book of Abstracts. General Directorate of Mineral

- Research and Exploration (MTA), Istambul.
- Maridet, O. y X. J. Ni. 2013. A new cricetid rodent from the early Oligocene of Yunnan, China, and its Evolutionary implications for early Eurasian Cricetids. *Journal of Vertebrate Paleontology* 33:185–194.
- Martín-Suárez, E. 1988. Une nouvelle espèce d'*Apodemus* (Rodentia, Mammalia) du Pliocène de la dépression de Guadix-Baza (Grenade, Espagne). *Geobios* 21:523–529.
- Martínez-Salanova, J. 1987. Estudio paleontológico de los micromamíferos del Mioceno inferior de Fuenmayor (La Rioja). *Ciencias de la Tierra, Paleontología, Volumen 10*. Instituto de Estudios Riojanos, Logroño, 99 pp.
- McKenna, M. C. y S. K. Bell. 1997. *Classification of Mammals above the species level*. Columbia University Press, New York, 631pp.
- Mein, P. y M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P. y M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Michaux, J., A. Reyes y F. Catzeflis. 2001. Evolutionary History of the Most Speciose Mammals: Molecular Phylogeny of Muroid Rodents. *Molecular Biology and Evolution* 18:2017–2031.
- Minwer-Barakat, R. 2005. Roedores e insectívoros del Turoliense Superior y el Plioceno del sector de la cuenca de Guadix. Tesis Doctoral, Universidad de Granada, Granada. 535 pp.
- Montgelard, C., S. Bentz, C. Tirard, O. Verneau, y F. M. Catzeflis. 2002. Molecular Systematics of Sciurognathi (Rodentia): The Mitochondrial Cytochrome b and 12S rRNA Genes Support the Anomaluroidea (Pedetidae and Anomaluridae). *Molecular Phylogenetics and Evolution* 22:220–233.
- Morales, J., L. Alcalá, M. A. Álvarez Sierra, M. Antón, B. Azanza, E. Badiola, J. P. Calvo, P. Carrasco, S. Fraile, I. García Paredes, E. Gómez, M. Hernández Fernández, L. Merino, A. J. Van der Meulen, C. Marín Escorza, P. Montoya, M. Nieto, S. Peigné, B. Pérez, P. Peláez-Campomanes, M. Pozo, V. Quirarte, M. J. Salesa, I. Sánchez, A. Sánchez Marco, P. Silva, M. D. Soria y A. Turner. 2004. Paleontología del sistema de yacimientos de mamíferos miocenos del Cerro de los Batallones, Cuenca de Madrid. *Geogaceta* 35:139–142.
- Morrone, J. J. 2000. *El lenguaje de la cladística*. Universidad Nacional Autónoma de México, México D. F., 109 pp.

- Mourik, A., H. A. Abels, F. J. Hilgen, A. Di Stefano y W. J. Zachariasse. 2011. Improved astronomical age constraints for the middle Miocene climate transition based on high resolution stable isotope records from the Mediterranean Maltese Islands. *Paleoceanography* 26:PA1210.
- Musser, G. G. y M. D. Carleton. 2005. Superfamily Muroidea; pp. 894–1531 en D. E. Wilson, D. M. Reeder (eds.), *Mammal Species of the World: A Taxonomic and Geographic Reference*. The Johns Hopkins University Press, Baltimore.
- Nelson, G. 1979. Cladistic analysis and Synthesis-principles and definitions, with a historical note on adansons families des plantes (1763-1764). *Systematic Zoology* 28:1–21.
- Novak, R. M. 1999. *Walker's Mammals of the World*. 6th Edition. The John Hopkins University Press. Baltimore. 1936 pp.
- Oliver, A. y P. Peláez-Campomanes. 2013. *Megacricetodon vandermeuleni*, sp. nov. (Rodentia, Mammalia), from the Spanish Miocene: a New evolutionary framework for *Megacricetodon*. *Journal of Vertebrate Paleontology* 33:943–955.
- Peláez-Campomanes, P. 1993. *Micromamíferos del Paleogeno Continental Español: Sistemática, Biocronología y Paleoecología*. Tesis Doctoral. Universidad Complutense de Madrid, Madrid, 388 pp.
- Peláez-Campomanes, P., J. Morales, M. A. Álvarez Sierra, B. Azanza, S. Fraile, I. García Paredes, M. Hernández Fernández, E. Herráez, M. Nieto, B. Pérez, V. Quiralte, M. J. Salesa, I. M. Sánchez y D. Soria. 2003. Updated biochronology of the Miocene mammal faunas from the Madrid basin (Spain); pp. 431–441 en J. W. F. Reumer and W. Wessels (eds.), *Distribution and migration of tertiary mammals in Eurasia. A volume in honour of Hans de Bruijn*. *Deinsea* 10.
- Platt, N. H. y V. P. Wright. 1992. Palustrine carbonates and the florida everglades-towards an exposure index for the fresh-water environment. *Journal of Sedimentary Petrology* 62:1058–1071.
- Prieto, J. Kleinsäuger-Biostratigraphie und Paläoökologie des höheren Mittelmiozäns (MN 8) Bayerns: Spaltenfüllungen der Fränkischen Alb und Lokalitäten der Oberen Süßwassermolasse im Vergleich. Tesis Doctoral. Universidad Ludwig-Maximilian de Munich. Munich, 217pp.
- Prieto, J., M. Bohme y M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Paratethys. *Geologica Carpathica* 61:419–436.

- Prieto, J., C. Angelone, I. Casanovas-Vilar, M. Gross, J. Hír, L. W. Van den Hoek Ostende, L. C. Maul y D. Vasilyan. 2014. The small mammals from Gratkorn: an overview. *Palaeobiology and Palaeoenvironments* 94:135–162.
- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata Palasiatica* 48:27–47.
- Reig, O. A. 1984. Geographic-Distribution and evolutionary history of South-American Muroids, Cricetidae, Sigmodontinae. *Revista Brasileira De Genetica* 7:333–365.
- Rummel, M. 1995. *Cricetodon bolligeri* n. sp. ein neuer Cricetide aus dem Obermiozän von Petersbuch bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung fuer Palaeontologie und Historische Geologie* 35:109–123.
- Rummel, M. 1999. Tribe Cricetodontini; pp. 359–364 en G. Rößner and K. Heißig (eds.), *The Miocene land mammals of Europe*, München.
- Rummel, M. 2000. Miocene Cricetodontini from Petersbuch near Eichstaett. The genus *Cricetodon* Lartet 1851. *Senckenbergiana lethaea* 80:149–171.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde, Series B* 311:1–6.
- Rummel, M. y D. Kälin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Metteltmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Sanz Rubio, E. 1999. Análisis de los sistemas deposicionales carbonáticos y evaporíticos del Neógeno de la Cuenca de Calatayud (provincia de Zaragoza). Tesis Doctoral. Universidad Complutense de Madrid, Madrid, 579 pp.
- Sanz-Rubio, E., S. Sánchez-Moral, J. C. Cañaveras, H. Abdul-Aziz, J. P. Calvo, S. Cuezva, A. V. Mazo, J. M. Rouchy, C. Sesé y J. A. Van Dam. 2003. Síntesis de la cronoestratigrafía y evolución sedimentaria de los sistemas lacustres evaporíticos y carbonatados neógenos de la Cuenca de Calatayud-Montalbán. *Estudios Geológicos* 59:83–105.
- Seehuber, U. 2008. Litho- und biostratigraphische Untersuchungen in der Oberen Süßwassermolasse in der Umgebung von Kirchheim in Schwaben. Tesis Doctoral, Ludwig-Maximilians-Universität, Munich, 359 pp.
- Sesé Benito, C. 1977. Los cricétidos (Rodentia, Mammalia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno Cuaternario* 8:127–180.
- Sevilla, P. 1988. Estudio paleontológico de los Quirópteros del Cuaternario español. *Paleontologia i Evolució* 22:113–233.

- Shevenell, A., J. P. Kennett y D. W. Lea. 2004. Middle Miocene Southern Ocean cooling and Antarctic cryosphere expansion. *Science* 305:1766–1770.
- Simpson, G. C. 1945. The principles of classification and a classification of mammals. *Bulletin of the American Museum of Natural History* 85:1–350.
- Sokal, R. R. y F. J. Rohlf. 1981. Taxonomic congruence in the *Leptodomophra* reexamined. *Systematic Zoology* 30:309–325.
- SPSS. 2002. Statistical Package for the Social Sciences for Windows, Rel. 11.5.1. Chicago: SPSS Inc.
- Thiede, J. y T. O. Vorren. 1994. The Arctic Ocean and its geologic record: research history and perspectives. *Marine geology* 119:179–184.
- Ünay E., H. De Bruijn y G. Saraç. 2003. A preliminary zonation of the continental Neogene of Anatolia based on rodents; pp. 539–547 en J.W. F. Reumer and W. Wessels (eds.), *Distribution and migration of Tertiary Mammals in Eurasia. A volume in honour of Hans de Bruijn*. Deinsea, Rotterdam.
- Van Dam, J. A. 1997. The small mammals from the Upper Miocene of the Teruel-Alfambra region (Spain): Paleobiology and Paleoclimatic reconstructions. *Geologica Ultraiectina* 156:204 pp.
- Van Dam J., L. Alcalá, A. Alonso-Zarza, J. P. Calvo, M. Garcés y W. Krijgsman. 2001. The Upper Miocene mammal record from the Teruel-Alfambra region (Spain): the MN system and continental Stage/Age concepts discussed. *Journal of Vertebrate Paleontology* 21: 367–385.
- Van Dam, J., W. Krijgsman, H. A. Abels, M. A. Álvarez-Sierra, I. García-Paredes, P. López-Guerrero, P. Peláez-Campomanes y D. Ventra. En prensa. Updated chronology for middle to late Miocene mammal sites of the Daroca area (Calatayud-Montalbán Basin), Spain. DOI: <http://dx.doi.org/10.1016/j.geobios.2014.07.002>.
- Van Dam, J., H. Abdul-Aziz, M. A. Álvarez Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. Lourens, P. Mein, A. J. Van der Meulen y P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodents fauna of the Mio-Pliocene continental sediments of the Teruel-Alfambra region, Spain. *Utrecht micropaleontological bulletins. Special publication*. Utrecht 2:1–217.
- Van den Hoek Ostende, L. W. 2003. Insectivores (Erinaceomorpha, Soricomorpha, Mammalia)

- from the Ramblian of the Daroca-Calamocha area. *Coloquios de Paleontología, Volumen Extraordinario* 1:281–310.
- Van den Hoek Ostende, L. W., P. López Guerrero, P. Peláez-Campomanes, M. A. Álvarez-Sierra e I. García-Paredes. 2012. Early Late Miocene insectivores (Eulipotyphla, Mammalia) from the Cañada section (Province of Zaragoza, east Spain). *Comptes rendus Palevol* 11:495–506.
- Van der Meulen, A. J., P. Peláez-Campomanes y R. Daams. 2003. Revision of medium-sized Cricetidae from the Miocene of the Daroca-Villafeliche area in the Calatayud-Teruel basin (Zaragoza, Spain). *Coloquios de Paleontología, Volumen Extraordinario* 1:385–441.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver y P. Peláez-Campomanes. 2012. Updated Aragonian biostratigraphy: Small Mammal distribution and its implications for the Miocene European Chronology. *Geologica Acta* 10:1–24.
- Wilson, D. E. y D. M. Reeder. 2005. *Mammal Species of the World. A Taxonomic and Geographic Reference* (3rd edition), Johns Hopkins University Press. 142 pp.
- Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.
- Zachos, J., M. Pagani, L. Sloan, E. Thomas y K. Billups. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693.

III. Paleontología sistemática

Chapter 5

Cricetodon soriae



REVISION OF *CRICETODON SORIAE* (RODENTIA, MAMMALIA), NEW DATA FROM THE MIDDLE ARAGONIAN (MIDDLE MIOCENE) OF THE CALATAYUD-DAROCA BASIN (ZARAGOZA, SPAIN)

“Dolores Soria, sin duda, ha puesto una firme base para que algunos de nosotros podamos continuar investigando en uno de los temas más atractivos de la paleontología de mamíferos”

Jorge Morales

Resumen

5

En el presente capítulo, se describe el registro más antiguo de *Cricetodon* (Cricetidae, Rodentia, Mammalia) del área tipo del Aragoniense (Cuenca de Calatayud-Daroca, Zaragoza, España). Los fósiles presentes en las secciones de Las Umbrías (LUM) y Las Planas (LP) se atribuyen a *Cricetodon soriae*, especie descrita en la localidad de Somosaguas (SOM) de la Cuenca de Madrid (biozona E, MN5). El material estudiado (52 dientes yugales) pertenece a ocho localidades que abarcan un intervalo de tiempo de 14,06 a 13,80 Ma, cubriendo toda la biozona E. El análisis morfométrico del material de Calatayud-Daroca permite presentar las descripciones más específicas y detalladas de *C. soriae* hasta la fecha. Se presentan nuevas descripciones completas del material tipo de SOMN, así como una diagnosis de *C. soriae*. Se han hecho comparaciones con todas las especies de *Cricetodon* descritas hasta la fecha. Se discuten las diferencias entre *Cricetodon aureus* y *Cricetodon meini*, los resultados arrojados por el estudio de más de 100 terceros molares superiores de Vieux-Collonges hacen posible, por primera vez, distinguir los M3 de *C. aureus* y *C. meini* y comparar ambos con *C. soriae*. La presencia de características comunes indica una relación más estrecha de *C. soriae* con las especies de *Cricetodon* de Asia Menor que con los primeros representantes de *Cricetodon* en Europa central y oriental. La distribución de *C. soriae* en la Península Ibérica es coincidente con un evento de enfriamiento global y aumento de aridez en el suroeste de Europa, y con el evento de dispersión faunística llamado *Conohyus/Pliopithecus* que tuvo lugar durante la biozona local E.

Referencia

- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013. Revision of *Cricetodon soriae* (Rodentia, Mammalia), new data from the middle Aragonian (middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.



Abstract

The oldest record of *Cricetodon* (Cricetidae, Rodentia, Mammalia) from the Aragonian type area (Calatayud-Daroca basin, Zaragoza, Spain) is described. The remains from Las Umbrías (LUM) and Las Planas (LP) sections are attributed to *Cricetodon soriae*, described in the locality Somosaguas N (SOMN) from the Madrid Basin (biozone E, MN5). The studied material (including 52 jugal teeth) belongs to eight localities spanning a time interval from 14.06 to 13.80 Ma, covering the entire biozone E. A morphometric analysis of the Calatayud-Daroca material provides the most specific and detailed descriptions of *C. soriae* until now. New complete descriptions of the type material from SOMN and an emended diagnosis of *C. soriae* are presented. Comparisons with all the species of *Cricetodon* described to date have been made. Differences between *Cricetodon aureus* and *Cricetodon meini* are discussed; the results yielded by the study of more than 100 upper third molars from Vieux-Collonges make it possible, for the first time, to distinguish between *C. aureus* and *C. meini* upper M3 elements and to compare them both with *C. soriae*. The presence of common characteristics indicates a closer relationship of *C. soriae* with *Cricetodon* from Asia Minor than with the first representatives of *Cricetodon* in central and eastern Europe. The distribution of *C. soriae* in the Iberian Peninsula is coincident with the global cooling event and the increasing in aridity in south-western Europe, and the faunal dispersion event called *Conohyus/Pliopithecus* that took place

INTRODUCTION

The Calatayud-Daroca Basin, situated in the north-eastern part of the Iberian Peninsula (Fig. 5.1), is filled by a thick succession of continental deposits ranging from the Palaeogene to the early Pliocene (Daams et al., 1999a). The stratigraphical superposition of the fossiliferous levels, their abundance and richness of micromammal fossil remains, make this basin interesting for the study of rodent evolution, biostratigraphy and biochronology (Daams et al., 1998, 1999a). One of its stratigraphical sections is the stratotype of the Aragonian continental stage that comprises the uppermost part of the lower Miocene and the entire middle Miocene (Daams et al., 1977; Daams et al., 1981). New discoveries in the Calatayud-Daroca Basin have allowed the detailed description of the Aragonian stage (Daams et al., 1999b; Van der Meulen et al., 2012). The Aragonian type area has provided fossil remains which offer detailed information of rodent faunas (Álvarez-Sierra, 1987; Cuenca Bescós, 1988; De Bruijn and Saraç, 1992; García-Paredes, 2006; García-Paredes et al., 2009, 2010; Oliver et al., 2009; Peláez-Campomanes, 2001; Van der Meulen et al., 2003, 2012). It is situated near Villafeliche, between Calatayud and Daroca villages (province of Zaragoza). It is composed of four sections: Vargas, Valdemoros, Las Umbrías and Las Planas (Daams et al., 1999a). This study focuses on the older localities situated in Las Umbrías and Las Planas sections. The age estimation of the localities ranges from 14.06 to 13.8 Ma (Van Dam et al., 2006), corresponding to the local biozone E (Daams et al., 1999b; Garcés et al., 2003; Van der Meulen et al., 2012). This local biozone can be correlated to the upper part of the Neogene Mammal zone MN5 (Daams et al., 1999a, 1999b; Van der Meulen et al., 2012).

The species revised here belongs to the genus *Cricetodon* Lartet, 1851, a large-sized rodent of the family Cricetidae. *Cricetodon* includes 22 fossil species; four of them are present in the Calatayud-Daroca Basin (Freudenthal, 1966; De Bruijn et al., 1993; López-Guerrero et al., 2009). The first occurrence of this genus is *C. versteegi* from the locality of Inkonak (MN1, “Agenian”, early Miocene) in Anatolia (De Bruijn et al., 1993). The youngest record is not clear. In the Nord Alpine Foreland Basin (NAFB) the presence in the MN9 (Vallesian, late Miocene) of *Cricetodon* sp. from Nebelberg TGL III (Prieto et al., 2010) is subject to doubtful taxonomical assignment, since it is considered as *Hispanomys* sp. by Kálin and Kempf (2009). The allocation to the MN9 in the case of *Cricetodon* sp. from Petersbuch 14 (NAFB) and *C. albanensis* from La Parrilla and Buenaventura 3 (Duero Basin, Spain) remains uncertain (Rummel and Kálin, 2003; García Moreno, 1988). Excluding these occurrences, the youngest record of *Cricetodon* is *C.*

klariankae from Felsőtárkány-Felnémet 2/3 and 2/7 (MN7/8, Aragonian, middle Miocene) in the Felsőtárkány Basin, Hungary (Hír, 2007).

The taxonomical status of some species attributed to *Cricetodon* has been problematic. This is because *Cricetodon* is one of the first described fossil small-mammal genera and the diagnosis proposed by Lartet, (1851) presents very inclusive criteria. Therefore, a large number of European species have been accommodated within this genus. Some of them are now included in genera belonging to different Cricetidae tribes like *Megacricetodon minor* (Lartet, 1851) or *Democricetodon larteti* (Schaub, 1925).

Cricetodon is a very important genus in the biostratigraphy of the continental deposits from the middle and upper Aragonian (De Bruijn and Saraç, 1992; Daams et al., 1998, 1999a). The presence of *Cricetodon* is one of the criteria used for the definition of the local biozone E in Calatayud-Daroca and Madrid basins (Van der Meulen et al., 2011, 2012). The systematics and evolution of these cricetids is not clear, despite the significant research carried out in the past, and the taxonomical status of *Cricetodon* has changed through time (e.g., Freudenthal, 1966; Mein and Freudenthal, 1971a, 1971b). The first studies of the genus in the Calatayud-Daroca Basin were made by Freudenthal (1963, 1966). He described several species and proposed the first phylogeny of *Cricetodon*. This first approach was later improved by the work of Mein and Freudenthal (1971a, b) in which most of species of this genus were described in detail. Mein and Freudenthal (1971a) emended the original diagnosis of *Cricetodon* made by Lartet in 1851. Additionally, they proposed four new subgenera within it: *Hispanomys*, *Ruscinomys*, *Pseudoruscinomys* and *Cricetodon*. Because this emended diagnosis refers simultaneously to the four subgenera, when Van de Weerd (1976) erected *Hispanomys* and *Ruscinomys* as genera and synonymized *Pseudoruscinomys*, the proposed diagnosis for the subgenus by Mein and Freudenthal (1971a) should be considered as generic. Since then, the most recent emended diagnosis of *Cricetodon* has been proposed by De Bruijn et al. (1993), being widely accepted and the most used to date. As a result of the aforementioned taxonomical status changes and the fact that the limits between these genera are not clear enough, some species originally included within the genus *Cricetodon* have been placed in other Cricetodontini genera; that is the case, for example, of *Hispanomys aguirrei* (Sesé, 1977), *H. lavocati* (Freudenthal, 1966) or *H. lusitanicus* (Schaub, 1925).

Regarding the species from the biozone E in Calatayud-Daroca Basin, De Bruijn et al. (1993) pointed out that the material present in Las Umbrías 11 (previously named as Las Planas 5A2) and the remains of Las Planas 4A and 4B, suggest a new species of *Cricetodon*. Despite these preliminary conclusions and the new material discovered since then, the remains of *Cricetodon* from the biozone E of Las Umbrías and Las Planas sections have neither been completely described nor assigned to particular species until the present paper.

Different authors have focused on the study of the faunas of Cricetodontini from other regions, e.g. Anatolia in Turkey, and their relationship with the oldest cricetid records in Europe (Sen and Ünay, 1979; Klein Hofmeijer and De Bruijn, 1988; De Bruijn et al., 1993; De Bruijn and Ünay, 1996; Rummel, 1998). The efforts of the micromammal researchers of the Calatayud-Daroca Basin have been focused on the *Cricetodon* representatives from younger localities (Álvarez-Sierra et al., 2003; López-Guerrero et al., 2009); however, it is important to understand the first record of Cricetodontini from Calatayud-Daroca in order to complete the revision of the whole middle Miocene rodent faunas from this basin that is currently in progress (López-Guerrero et al., 2007; López-Guerrero et al., 2008).

Therefore, the main aims of this study are: to characterize the first representatives of *Cricetodon* in the Aragonian type area through morphological and metrical analysis of the jugal teeth, to specify their stratigraphical distribution during the middle Aragonian and to compare them morphometrically with other representatives of the genus present in Europe and Asia Minor.




FIGURE 5.1. Geological map of the Calatayud-Daroca Basin. The square marks the situation of Aragonian type area (Modified after García-Paredes et al., 2010).

MATERIAL & METHODS

during the local biozone E.

This study includes morphological and metrical analysis of 52 cheek teeth of *Cricetodon* from the Calatayud-Daroca Basin. Table 5.1 shows the different localities studied, the number of molars examined and the abbreviations used throughout this paper. The anatomical abbreviations for upper molars are M1, M2, M3 and, for lower molars, m1, m2, m3. The studied material belongs to eight localities spanning a time interval from 14.06 to 13.80 Ma. (Van Dam et al., 2006). In addition, we revised material of *C. soriae* from the type locality Somosaguas N (Madrid Basin) stored at the Universidad Complutense de Madrid (Spain); the type material of *C. bolligeri* from Petersbuch 10 stored at Bayerische Staatssammlung für Paläontologie und Geologie, Munich (Germany); *C. sansaniensis* from the type locality of Sansan stored at the Muséum d'Histoire Naturelle de Bâle (Switzerland); the type material of *C. aureus* and *C. meini* from Vieux-Collonges, and *C. albanensis* from La Grive-Saint Alban stored at the Nederlands Centrum voor Biodiversiteit-Naturalis (NCB-Naturalis), Leiden (The Netherlands); and finally, *C. jotae* from its type locality, Manchones, stored at the Instituut voor Aardwetenschappen, Utrecht University (The Netherlands). The remaining species of *Cricetodon* have been studied using data available in the published descriptions of their type

<i>Cricetodon soriae</i>										
Age (Ma)	Locality	Abb.	Collections	M1	M2	M3	m1	m2	m3	N
13.80	Las Umbrías 20	LUM20	MNCN	2	1	2	1	—	1	7
13.88	Las Planas 4C	LP4C	RGM	—	—	2	—	—	—	2
13.95	Las Umbrías 19	LUM19	MNCN	2	1	2	—	—	1	6
13.96	Las Planas 4B	LP4B	IIAUU	—	—	—	—	1	1	2
—	Las Planas 4BA	LP4BA	MNCN	1	1	2	2	1	—	7
13.98	Las Planas 4A	LP4A	IIAUU	—	—	—	—	1	—	1
14.04	Las Umbrías 16	LUM16	MNCN	—	—	—	1	2	2	5
14.06	Las Umbrías 11	LUM11	MNCN	2	2	1	2	4	3	14
14.10–13.80	Somosaguas N	SOMN/5	UCM	—	1	1	1	2	3	8

TABLE 5.1. Chronological information of the studied localities: Aragonian (Middle) Continental Stage; Mammal Neogene Zone (MN5) proposed by Mein (1975) for Europe and revised by de Bruijn et al. (1992); local biostratigraphical zone (E) after Daams et al. (1999a); numerical ages of the localities after Van Dam et al. (2006); localities, abbreviations for the localities and institutions where the fossils are stored (see institutional abbreviations). Abbreviations: Abb, abbreviations for the localities; N, number of elements of *Cricetodon soriae* studied in each locality of Calatayud-Daroca and Madrid.

Species	Locality	Age	Reference
' <i>Cricetodon</i> ' <i>fandli</i>	Gratkorn	MN7/8	Prieto et al., 2010
' <i>Cricetodon</i> ' <i>klariankae</i>	Felsőtárkány–Felnémet 2/3	MN7/8	Hir, 2007
<i>C. bolligeri</i>	Petersbuch 10	MN7/8	Ziegler et al., 2005
<i>C. jumaensis</i>	Petersbuch 18	MN7/8	Rummel, 2001
<i>C. cariensis</i>	Sarıçay	MN7	Sen and Ünay, 1979
<i>C. albanensis</i>	La Grive–Saint Alban	MN7	Mein and Freudenthal, 1971a
<i>C. engesseri</i>	Chräzerentobel 505m	MN7	Rummel and Kálin, 2003
<i>C. sansaniensis</i>	Sansan	MN6	De Bruijn et al., 1992
<i>C. pasalarensis</i>	Paşalar	late MN6	Peláez-Campomanes and Daams, 2002
<i>C. jotae</i>	Manchones	MN6, biozone G2	De Bruijn et al., 1992
<i>C. hungaricus</i>	Hasznos	MN6	De Bruijn et al., 1993
<i>C. caucasicus</i>	Belometchetskaya	late MN6	De Bruijn et al., 1992
<i>C. soriae</i>	Somosaguas	MN5, biozone E	Hernández-Fernández et al., 2006
<i>C. meini</i>	Vieux–Collonges	MN4–MN5	De Bruijn et al., 1992
<i>C. candirensis</i>	Çandır	MN5	De Bruijn et al., 2003
<i>C. aureus</i>	Vieux–Collonges	MN4–MN5	De Bruijn et al., 1992
<i>C. volkeri</i>	Dingshanyanchi Fm.	middle Miocene	Wu et al., 2009
<i>C. aliveriensis</i>	Aliveri	MN4	Klein Hofmeijer and De Bruijn, 1988
<i>C. wanhei</i>	Songlinzhuang	early Miocene	Qiu, 2010
<i>C. tobieni</i>	Horlak 1a	MN3–MN4	Tobien, 1978
<i>C. kasapligili</i>	Keseköy	MN3	De Bruijn et al., 1993
<i>C. versteegi</i>	Kilçak 3a	MN1	De Bruijn et al., 1993

TABLE 5.2. List of the comparison species studied, type locality and their ages. The references in where the dating for the sites is described are given on the fourth column.

localities. All the species of *Cricetodon* published to date are summarized in Table 5.2, including their type localities and ages.

Length and width measurements for each specimen, given in mm, have been taken using a Nikon Measuroscope 10 microscope with digital micrometer. The procedure used to measure the maximum length and width is shown in Figure 5.2.

All the measurements, the number of measured specimens, minimum, maximum and mean values are given in Table 5.3. In order to compare the studied material with the large-sized species of *Cricetodon*, Table 5.4 includes measurements taken from the literature. In the case of the *C. aureus* M3, we applied our morphological criteria to separate *C. aureus* from *C. meini* and, afterwards, we recalculated the minimum, mean and maximum values using Mein and Freudenthal (1971b) dataset of the type series included in the collection database of the NCB-Naturalis, Leiden (The Netherlands).

The nomenclature used for the structures present in the occlusal surface, modified after

Mein & Freudenthal (1971b), is summarized in Figure 5.3. The photographs were taken with an environmental Scanning Electron Microscope FEI Quanta 200. The image processing was made using Adobe Photoshop software. The molars are figured as if they were from the right side; reversed images have been indicated by an underlined label.

Institutional Abbreviations—IAUU, Instituut voor Aardwetenschappen, Utrecht University, The Netherlands; MNCN, Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain; RGM, NCB-Naturalis, former Rijksmuseum van Geologie en Mineralogie, Leiden, The Netherlands; UCM, Universidad Complutense de Madrid, Spain.






FIGURE 5.2. Measurement method for length and width used in the present paper. The upper draws represent left upper teeth (from M1 to M3) and the lower draws represent right lower m3, m2 and m1. Abbreviations: L, length; W, width.

M1



M3



m1




FIGURE 5.3. Terminology of the parts of the cheek teeth of *Cricetodon*. Only the M1, M3 and the m1 morphology has been drawn; however, this nomenclature can be applied for the rest of the upper and lower molars. (Modified after Mein and Freudenthal, 1971b).

SYSTEMATIC PALEONTOLOGY

Order RODENTIA Bowdich, 1821

Family CRICETIDAE Fischer, 1817

Subfamily CRICETODONTINAE Stehlin and Schaub, 1951

Tribe CRICETODONTINI Simpson, 1945

Genus *CRICETODON* Lartet, 1851

Diagnosis

Original Diagnosis (Lartet, 1851:20, translated from French)—“[Rodents with tubercular cheek teeth. Same tooth pattern and number of teeth as rats: six for each mandible. Molars present fewer tubercles than those of domestic rats. They are more related to the molars of the hamsters. However, they differ from the latter by the presence of an additional tubercle when the first upper and lower molars are considered. The most significant similarity with hamsters concerns the presence of an arterial hole on the exterior condyle of the humerus that is absent in the case of domestic rats.]”

Emended Diagnosis (Mein and Freudenthal, 1971a:17, translated from French)—“[Medium to large-sized Cricetodontini, the upper incisors are smooth or lined by a ‘dipodoid’ type groove. The M1 has four or five roots, the posterior root of m2 can be simple or bifurcate; the M1 has a simple or bifurcate anterocone, the upper molars show a trend in developing ectolophs; the M1 often shows protostyls and parastyls; lower molars often have protostylids, ectostylids and ectomesolophids; the m1 always has a descending labial ridge from the higher part of the anteroconid; molars are brachyodonts or hypsodonts, but the height of the crown of an unworn tooth never exceeds the length of the roots of a worn teeth.]”

Remarks—As explained in the introduction, this emended diagnosis refers simultaneously to the subgenera, *Hispanomys*, *Ruscinomys*, *Pseudoruscinomys* and *Cricetodon* sensu Mein and

Freudenthal (1971a).

Diagnosis for the subgenus (Mein and Freudenthal, 1971a:17, translated from French)— “[Medium to large-sized Cricetodontini, the upper incisors are smooth, the ectolophs of the upper molars are absent or weak; the molars are brachyodont; the anterocone on the upper M1 is simple or bifid; the upper M1 has four roots; the lower m2 has two or three.]”

Emended Diagnosis (De Bruijn et al., 1993:177)—“Small, medium or large-sized Cricetodontinae with low-crowned cheek teeth. Cusps of cheek teeth ‘inflated’. Anteroconid of m1 bearing a small rounded cusp. Majority of the m1 with double metalophulid or metalophulid II only. Anterocone of M1 single or double. Labial branch of the anterolophid of the m2 incorporated in the metalophulid I. The M1 and M2 with three or four roots. Free posterior arm of the hypoconid absent in all lower molars. Occlusal surface of all cheek teeth worn flat in senile specimens.”

Type Species—*Cricetodon sansaniensis* Lartet, 1851

Other Species Included—See Table 5.2.

CRICETODON SORIAE López Martínez, Cárđaba, Salesa, Hernández Fernández, Cuevas González and Fesharaki, 2006

(Figs. 5.1–5.7; Tables 5.1 and 5.3)

Cricetodon soriae n. sp. López Martínez, Cárđaba, Salesa, Hernández Fernández, Cuevas González and Fesharaki, 2006 in Hernández Fernández et al., 2006: 275, figs. 10–11 (original description).

Cricetodon meini Freudenthal, 1963:74 (only for Las Planas 4A and 4B) *Cricetodon* n. sp. 3 in De Bruijn, Fahlbusch, Saraç and Ünay, 1993: 206, Pl.17

Material

Original Type Locality— Somosaguas N (SOMN), Madrid Basin, Spain.

New Localities—See Table 5.1.

Stratigraphic Range—middle Miocene (middle Aragonian, upper part of the MN5, local biozone E).

Geographical Range—central basins of the Iberian Peninsula.

Diagnosis

Original Diagnosis (López Martínez, Cárdbaba, Salesa, Hernández Fernández, Cuevas González and Fesharaki, 2006 in: Hernández Fernández et al., 2006:275, translated from Spanish)—“[Medium-sized *Cricetodon*, with a relatively short and wide m1 and a relatively developed m3; reduced ectoloph, mesoloph and mesolophid; posterior metalophid in m1. It differs from *C. aureus* by the dental proportions and its smaller size; it differs from *C. jotae* by its more robust m1, the posterior metalophid in the m1, less development of the ectoloph on the M2, the larger absolute width in all elements and the larger absolute and relative size of M2, m2 and m3.]”

Emended Diagnosis—Medium to large-sized *Cricetodon*. Upper and lower molars without styls or stylids. M1 with anterior and posterior protolophule; anterior ectoloph on the M1 absent. Anterior ectoloph on the M2 is weak and it can be observed as an enamel spur; posterior ectoloph well-distinguished and short, never connected to the paracone. High-crowned M3 with a short and disconnected posteroloph; well-developed hypocone; neo-entoloph hardly developed or absent; and with a continuous entoloph. Short and wide m1, metalophulid I, always present, may be disconnected: metalophulid II always present. The lower third molar is large and longer than the m1; the length m1/ length m3 index is more or less 1; variable mesolophid on the m3, it may be short and connected to the metaconid, long and connected either to the metaconid or to the entoconid, or double connected to both the metaconid and the entoconid.

Differential Diagnosis—*Cricetodon soriae* differs from *C. aureus* Mein and Freudenthal, 1971b in: larger size of the m3; more developed hypocone and more degree of hypsodonty on the M3; lack metalophulid I alone on the m1; presence of entoconid spur on the m1. Differs from *C. caucasicus* Argyropulo, 1938, *C. kasapligili* De Bruijn, Fahlbusch, Saraç and Ünay, 1993 and *C. tobieni* De Bruijn, Fahlbusch, Saraç and Ünay, 1993, in: split anterocone; four-rooted M1; larger size. Differs from *C. sansaniensis* Lartet, 1851, in: deeply split anterocone and double protolophule. Differs from *C. versteegi* De Bruijn, Fahlbusch, Saraç and Ünay, 1993 in: less developed anterolophs and ectolophs; larger size. Differs from *C. aliveriensis* Klein Hofmeijer and De Bruijn, 1988 and *C. pasalarensis* (Tobien, 1978) in: larger size. Differs from *C. meini* Freudenthal, 1963 in: larger size; m1 with metalophulids type I and type II always present together on the same specimen. Differs from *C. albanensis* Mein and Freudenthal, 1971a, *C. bolligeri* Rummel, 1995, *C. jumaensis* Rummel, 2001 and *C. engesseri* Rummel and Kálin, 2003 in: m1 with metalophulids type I and type II always present together on the same specimen. Differs from *C. jumaensis* Rummel, 2001 and *C. bolligeri* Rummel, 1995 in: absence of styls on the M1. Differs from *C. candirensis* (Tobien, 1978) and *C. hungaricus* (Kordos, 1986) in: less developed anterolophs and ectolophs. Differs

from *C. cariensis* (Sen and Ünay, 1979) in: absence of enamel-coated valleys on the upper molars; ectolophs more developed. Differs from *C. jotae* Mein and Freudenthal, 1971a in: the larger size; m1 with metalophulids type I and type II always present together on the same specimen; less developed anterolophs and ectolophs. Differs from *C. klariankae* Hír, 2007 in: the less developed ectolophs. Differs from “*Cricetodon*” *fandli* Prieto et al., 2010 in: the less developed ectolophs; absence of lingual spur of the anterolophule.

CRICETODON SORIAE FROM THE CALATAYUD-DAROCA BASIN

Description of the Studied Material

M1—Material: LUM11:2; LP4BA:1; LUM19:2; LUM20:2. This molar has four roots. The tooth is very wide and short, having a rounded outline. A shallow groove on the anterior wall divides the anterocone into two parts. The connection between the protocone and anterocone is through the anterolophule. This lophule can present as an enlargement of the enamel. A posterior protolophule connects the paracone with the posterior branch of the protocone. An anterior protolophule connects the paracone with the anterolophule in most cases; it is absent only in LP4BA and LUM20. The anterior ectoloph is absent. The posterior ectoloph is simple, never complete. The mesoloph is short in LUM11 and LUM19, and it is incipient in LUM20. The mesoloph-ectoloph connection is absent. The metalophule is directed backwards. The labial posteroloph is quite thin but well delimited. The lingual posteroloph is absent. The anterosinus is closed by either a cingulum (LUM11) or two ridges attached to the protocone and the hypocone respectively (LUM19). The protosinus and the mesosinus are closed by a cingulum. The sinus is either open (one out of two specimens in LUM20) or closed by a cingulum, but never by a styl. The enamel on the base of the protocone and the hypocone is rough (Fig. 5.4).

M2—Material: LUM11:2; LP4BA:1; LUM19:1; LUM20:1. The tooth has four roots. It is short and wide, but with an outline more rectangular than in the M1. The anterocone is reduced to an enlargement of the enamel on the anteroloph. The loph that connects the anteroloph with the paracone is reduced and generally lower than the anteroloph (LUM19 and LUM20). This loph can be absent (LUM11 and LP4BA). The posterior ectoloph is simple and disconnected. Only a few specimens present mesoloph and it is short (LUM11) or incipient (LP4BA). It shows neither enamel coated valley, nor an entomesoloph. Both labial and lingual posterolophs are present but weak. The anterosinus is always closed. The protosinus is well developed and it is closed by a cingulum in most specimens, although in one case (LUM20) is closed by a styl. The mesosinus, which is well developed, can be closed, either by a cingulum (LUM11, LP4BA, one out of two

specimens in LUM19, and LUM20) or by two ridges (one out of two specimens in LUM19) but never by a styl. The sinus can be either open (LP4BA and LUM20) or closed by a cingulum (LUM11) (Fig. 5.4).

M3—Material: LUM11:1; LP4BA:2; LUM19:2; LP4C:2; LUM20:2. It has a subtriangular and rounded outline with three roots. Both labial and lingual anterolophs are present, but the lingual branch is weaker than the labial one. The anterior ectoloph is absent. The posterior ectoloph can be simple (LUM11, LP4BA, LUM19 and one out of two specimens in LP4C) or double (one out of two specimens in LP4C and LUM20). It is never complete. Most of the specimens have a mesoloph and its length is variable. It may be short (one out of two specimens in LP4BA and one out of two specimens in LUM20), medium length, about half the distance between longitudinal ridge and labial border (LUM11) or long, reaching the labial border (LP4C). The mesoloph-ectoloph connection is absent. The entoloph is continuous and there is no trace of the neo-entoloph. The labial posteroloph is usually well developed but short and it never connects to the metalophule or the metacone. It is absent in only one specimen from LP4C. The sinus is anterior or transversal (LUM11) (Fig. 5.4).

m1—Material: LUM11:2; LUM16:1; LP4BA:2; LUM20:1. This molar has an elongated shape and two roots. The anteroconid is situated on the longitudinal axis of the occlusal surface. The labial anterolophid is a well-developed ridge that connects the anteroconid with the protoconid in some specimens (LUM11 and LUM16); on the rest (LP4BA and LUM20) this ridge does not reach the protoconid. The lingual anterolophid is present in only one specimen (LUM16). In all cases, both metalophulid I and II are present together on the same specimen. The longitudinal ridge bears a mesolophid. This mesolophid can be short (LUM16 and one out of two specimens in LP4BA) or incipient. The ectomesolophid is present in only one specimen from LUM11. The entoconid spur is absent. The labial posterolophid is well developed (LUM11) or it can be absent (LUM20). The posterolophid is an oval cusp rather than a ridge, due to a constriction on the posterior wall. A low cingular ridge blocks the mesosinusid in one case (LP4BA), but in the rest, this valley is open. The sinusid is directed forwards and it can be blocked either by a cingular ridge (LUM11 and LP4BA) or by two ridges (LUM20) but never by a stylid. The posterosinusid is blocked by a cingular ridge. The protoconid and the hypoconid show rough enamel (Fig. 5.5).

m2—Material: LUM11:4; LUM16:3; LP4A:1; LP4BA:1; LP4B:1. The molar is almost rectangular in shape and the anterior and the posterior parts have similar width. The labial branch of the anterolophid is well developed. The lingual anterolophid is absent. The mesolophid is variable in length; it can be short (LP4A; LP4BA and LP4B) or long (one out of three specimens in LUM11), but it never reaches the lingual border; in some cases (two out of three specimens in LUM11, LUM16; LP4BA; LP4B), the mesolophid is curved and projected towards the metaconid, reaching it. The lingual posterolophid is constricted behind the hypocone; its lingual

part is cuspidate. The ectomesolophid is present in one case from LUM11. There is no anterosinusid and the mesosinusid is blocked by a cingular ridge in most cases, although it can also be open (one out of two specimens in LUM11 and one out of two specimens in LUM16). In most teeth, a cingular ridge is connecting the hypoconid with the protoconid at the labial border of the sinusid, whereas in one specimen from LUM11 the cingular ridge starts from the protoconid and it does not reach the hypoconid. The sinusid is never blocked by a stylid. A weak labial posterosinusid is formed on the posterior wall of the hypoconid. The lingual posterosinusid is equally blocked by a cingular ridge (LUM11 and LUM16) or by the structure resulting from the fusion of the lingual posterolophid and the entoconid (LP4A; LP4BA and LP4B). The enamel is rough on the base of the hypoconid and the protoconid (Fig. 5.5).

m3—Material: LUM11:3; LUM16:2; LP4B:1; LUM19:1; LUM20:1. This molar has a triangular but rounded outline. The labial branch of the anterolophid is long and it reaches the protoconid. The lingual branch of the anterolophid is absent. The mesolophid is always present and well developed. It may reach the lingual border (LUM11 and LUM16) or it may be of a medium length (LP4B and LUM19). In one case (LUM20), the mesolophid is short and connected with the metaconid. It can be connected both to the metaconid and to the entoconid in some cases (two out of three specimens in LUM11). The ectomesolophid is absent. The lingual posterolophid is high and extends to the entoconid. It can be somewhat constricted behind the hypoconid, its lingual part is cuspidate and a weak valley on the posterior wall of the tooth is developed. The hypolophid, which is transversal, can be connected to the anterior branch of the hypoconid. Some specimens have the mesolophid connected with the ectolophid (two out of three specimens in LUM11). In some cases, (LUM11, LUM16 and LUM19), a low cingular ridge connects the hypoconid with the protoconid on the sinusid, but in the rest (LP4B and LUM20), the sinusid is open. It never presents stylids. The mesosinusid is open in most specimens, although in one case from LUM16, it is blocked by a low cingular ridge. This structure is also present at the end of the lingual posterosinusid. In some cases (LP4B and LUM20) the lingual posterolophid projects to the entoconid closing it. The enamel of the hypoconid and the protoconid is rough (Fig. 5.5).

Discussion

The most important features of the studied cricetid material are: medium to large size, upper molars with three or four roots, and split anterocone on the M1. Most of the m1 have metalophid I and II, the anteroconid bearing a small rounded cusp and the lower molars often with ectomesolophids. These features agree with the proposed diagnosis of the genus *Cricetodon* sensu De Bruijn et al. (1993).

Furthermore, the studied material possesses features like: brachydont molars, with short or

absent ectolophs that fit into the group 5 defined by Prieto et al. (2010) for *Cricetodontini*.

Among the Calatayud-Daroca remains, we observe differences in the degree of the development of mesolophs and mesolophids in all elements. These structures are less developed on the upper molars, where they are short or reduced to an “incipient mesoloph” (an enlargement of the enamel on the longitudinal ridge) whereas, they are more developed on the lower molars. The mesolophids can be medium or long in the material from the older localities (LUM11, LUM16), although in the youngest locality (LUM20), it can form a short crest. The posterior ectolophs on the M3 also show variability in their development. Simple ectolophs on LUM11 and LP4A that develop a second branch on LP4C and LUM20 are present.

Another type of differences is in the frequency of occurrence of the cingular ridges that are blocking the main valleys. Structures like the cingulum that blocks the sinus and labial posteroloph on the M2 are present on the older localities (LUM11, LP4BA), whereas the youngest locality (LUM20) shows the sinus open and the labial posteroloph is absent. We suggest that the difference in frequency of particular states of character among the localities cannot be truly evaluated as the number of specimens in the sample is very low. This means that it is difficult to see the complete range of morphological variability of the species at a single locality. Nevertheless, differences among the most diagnostic features, like the size and degree of the development of the ectolophs and posterolophs on the upper molars, the presence of double protolophule and short posterolophs on the M3, or the presence of both metalophulids on the m1 have not been found. Thus, we consider that the studied material belongs to the same species.

CRICETODON SORIAE FROM THE MADRID BASIN

Cricetodon soriae has been described by López Martínez et al. (2006) on the locality Somosaguas N. This site was dated around 14.1 to 13.8 Ma (Luis and Hernando, 2000) corresponding to local biozone E (middle Aragonian) from the Madrid Basin (Hernández Fernández et al., 2006; Domingo et al., 2009a). In order to clarify the morphology of the type material of *C. soriae* from Somosaguas N, we have described it under the same morphological criteria as the material from Calatayud-Daroca. One of the paratypes, SOMN-53, was described by López Martínez et al. (2006) as an M2; however, following a comprehensive study of the material, we have identified this specimen as an M3. The different elements are not completely represented in *C. soriae* material from SOMN. The available material from the type locality corresponds to one M2, one M3, two m1, one m2 and two m3.

Descriptions of the Material from the Type Locality Somosaguas Norte.

M2—The only available specimen presents a square-like contour. The posterior ectoloph is simple and disconnected. The specimen does not present a mesoloph. It shows neither a mesoloph-ectoloph connection nor entomesoloph. The labial posteroloph is present but weak. The anterosinus is always closed. The mesosinus and the sinus are well developed and closed by a cingulum. The sinus is curved forwards (Fig. 5.6).

M3—This element is represented by only one fragment. The lingual anteroloph is present. The anterior ectoloph is absent. The posterior ectoloph is simple and not complete. The mesoloph is a slight enlargement of the enamel from the longitudinal ridge. There is no enamel-coated valley. The entoloph is continuous. The labial posteroloph is present but short and it is not connected to the metalophule or to the metacone (Fig. 5.6).

m1—The two specimens have an elongated outline in the occlusal view. The anteroconid is situated on the longitudinal axis of the occlusal surface. The labial anterolophid is a well-developed ridge and it does not reach the protoconid. The lingual anterolophid is absent. The metalophulid I is present, but not connected to the anteroconid. The metalophulid II connects the metaconid with the posterior branch of the protoconid. The longitudinal ridge bears a slight enlargement as a short mesolophid. The ectomesolophid is absent. The entoconid is not constricted. The labial posterolophid is not present. The posterolophid is an oval cusp rather than a ridge, due to a constriction behind the hypoconid. The anterosinusid and mesosinusid are not blocked by cingulids. The sinusid is directed forwards and blocked by a cingular ridge. The posterosinusid is open (Fig. 5.6).

m2—The outline of the only m2 present is almost rectangular and the width of the anterior and posterior parts is similar. The labial branch of the anterolophid is well developed and reaches the protoconid. The lingual anterolophid is absent. The mesolophid is short and reaches the metaconid. The lingual posterolophid is constricted on the posterior wall. The ectomesolophid is absent. There is no anterosinusid and the mesosinusid is closed by a ridge. A cingular ridge, descending from the protoconid, blocks the sinusid. The hypoconid presents a weak labial posterosinusid. The lingual posterosinusid is closed by a cingular ridge (Fig. 5.6).

m3—The labial branch of the anterolophid is long, reaching the protoconid. The lingual branch of the anterolophid is present, but weak. The mesolophid is present and well developed in the two studied specimens; it never reaches the lingual border (medium length) and is connected to the metaconid. The ectomesolophid is absent. The lingual posterolophid is high and extends to the entoconid. The hypolophulid, which is transversal, can be connected to the anterior branch of the hypoconid. None of the specimens have an enamel-coated valley. The sinusid is transversal and it is either closed by a low cingulum or by a small stylid. The mesosinusid is open. There is no labial posterosinusid. A small cingular ridge is present at the end of the lingual posterosinusid (Fig. 5.6).




FIGURE 5.4. *Cricetodon soriae*, from Calatayud-Daroca. Upper molars. Las Umbrías 11. A, inverted left M3, LUM11-321; B, inverted left M2, LUM11-3; C, right M1, LUM11-1; Las Umbrías 19. E, right M3, LUM19-212; F, right M1, LUM19-209. Las Umbrías 20. G, right M3, LUM20-221; H, right M2, LUM20-224; I, right M3, LUM20-222. Scale bar equals 1 mm.




FIGURE 5.5. *Cricetodon soriae*, from Calatayud-Daroca. Las Umbrías 11. A, inverted left m1, LUM11-6; B, right m2, LUM11-9; C, inverted left m2, LUM11-4; D, inverted left m3, LUM11-11; E, inverted left m2, LUM11-10; F, inverted left m3, LUM11-12. Las Umbrías 16. G, inverted left m3, LUM16-242; H, right m2, LUM16-241; I, right m3, LUM16-243. Las Umbrías 20. J, right m1, LUM20-223; K, right m3, LUM20-225. Las Umbrías 19. L, inverted left m3, LUM19-214. Scale bar equals 1 mm.




FIGURE 5.6. *Cricetodon soriae*, from Somosaguas N. A, left mandible, SOMN-2109 (Holotype); B, left m1, SOMN-2109 (i); C, left m2, SOMN-2109 (ii); D, left m3, SOMN-2109 (iii); E, inverted right m3, SOMN-50; F, left M2, SOMN-51; G, left M3, SOMN-53. Scale bars equals 0.5 mm and 1mm.

Discussion

These specimens from SOMN share several characters with the material from Calatayud-Daroca: the m1 with both metalophulids I and II present, the sinusid on the m2 closed by a cingulum and the m2 also has a medium mesolophid that is connected to the metaconid. The M2 has no anterior ectoloph and the posterior one is simple and not connected to the metacone. But the most important similarities are present on the m3. The m3 from both SOMN and Calatayud-Daroca is large. The values of the length m1/length m3 index are between 1.02 and 1.06 for *C. soriae* from SOMN and our specimens show values of 0.99. That means that the m3 is approximately as long as the m1 in both cases.

Based on the aforementioned characters, we conclude that there are morphometrical criteria enough to assign the material from Calatayud-Daroca to *C. soriae*. Although the Madrid and Calatayud-Daroca basins show differences in the abundance of some genera (Peláez-Campomanes et al., 2003; Hernández-Ballarín et al., 2011), the similarities on their rodent assemblages have made possible to recognize the local biozonation built for Calatayud-Daroca by Daams et al. (1999a) in the Madrid Basin (Peláez-Campomanes et al., 2003). In addition, these basins have been included in the southern biogeographical province due to the similarity between the faunas of both basins (Agustí et al., 1984; Agustí, 1990; Álvarez-Sierra et al., 1990; Alberdi and Azanza, 1997; Morales et al., 1999; Gómez-Cano et al., 2011).

COMPARISON WITH OTHER SPECIES OF *CRICETODON*

As a result of this study, our knowledge about *C. soriae* has increased. This allows us to make the comparisons with the other *Cricetodon* in Europe and Asia Minor.

A comparison of the measurements shows a clear difference in size between our material, and the small-sized *Cricetodon* i.e. *C. caucasicus*, *C. versteegi*, *C. pasalarensis*, *C. aliveriensis*, *C. kasapligili*, *C. tobieni*, *C. jotae* and *C. meini* (Argyropulo, 1938; Freudenthal, 1963; Mein and Freudenthal, 1971a, 1971b; Tobien 1978; Klein Hofmeijer and De Bruijn, 1988; De Bruijn et al., 1993; Daxner-Höck, 2003; Rummel, 2001; Markovic, 2008).

We have found several morphological differences in the species that are similar in size to *Cricetodon* from Calatayud Daroca. *C. candirensis* has upper molars with well-developed anterolophs and ectolophs (Tobien, 1978; Kordos, 1986). *Cricetodon cariensis* has enamel-coated valleys on the upper teeth, and less-developed ectolophs (Sen and Ünay, 1979) than our material. *Cricetodon bolligeri*, *C. engeseri* and *C. albanensis* possess type I metalophulids exclusively on the

m1 (Rummel, 1995; Rummel and Kälin, 2003; Kälin and Kempf, 2009), whereas metalophulid type II is always present in the material from Calatayud-Daroca. Although *C. jumaensis* presents the same kind of metalophulid (I+II type) as *C. soriae* in the m1 (7 out of 19), metalophulids type I without type II (4 out of 19) or type II alone (5 out of 19) are also found in the type locality of Petersbuch 18 (Rummel, 2000; Rummel, 2001). Furthermore, a styl is present on the holotype and the hypolophulid on the m3 is forward directed (Rummel, 2001), whereas *C. soriae* never have styls and the hypolophulid on the m3 is transversal.

Cricetodon sansaniensis was described by Lartet (1851) but detailed descriptions were given by Baudelot (1972). According to her descriptions, *C. sansaniensis* presents a single or weakly divided anterocone. The ectolophs in this species are absent in the half of the specimens from the type locality. However, a weak projection of the enamel in the paracone or the anterocone can be distinguished in the other half of the specimens (Baudelot, 1972; Maridet and Sen, 2012). By contrast, the material from Calatayud-Daroca never shows an anterior ectoloph on the M1. *Cricetodon klariankae*, defined in Hungary has anterior ectolophs and double posterior ectolophs on the M1, together with a reduced hypocone on the M3 (Hír, 2007). “*Cricetodon*” *fandli* presents a very well-developed ectolophs which reach the paracone and metacone respectively in the upper molars (Prieto et al., 2010).

Cricetodon soriae has been compared with the representatives of the genus from the central and eastern Europe basins that are close in age. The Swiss Molasse of the NAFB, Uzwill-Nutzenbuech dated at 14.9 Ma by Kälin and Kempf (2009) contains *Cricetodon* aff. *aureus*. This material shows features such as an entomesoloph and styls on the M1, but it has no anterior protolophule (Rummel and Kälin, 2003). On the German Molasse of the NAFB, the first occurrence of the genus corresponds to *Cricetodon* aff. *meini* from Eberhausen (Boon, 1991; De Bruijn et al., 1993). This fossil site is dated in 15.2 Ma by Abdul Aziz et al. (2010). This material has differences with *C. soriae* such as the presence of solely metalophulid type I and ectomesolophid on the m1 (Boon, 1991; de Bruijn et al., 1993). *Cricetodon hungaricus*, from Hasznos, Hungary (Kordos, 1986; de Bruijn et al., 1993) shows different features in comparison with *C. soriae*, such as the presence of metalophulid type II without type I, shorter mesolophids on the m3, and shorter M3 and m3 (de Bruijn et al., 1993). In Serbia, the species *C. meini* has been found in the MN6 localities of Bele Vode and Lazarevac (Markovic, 2008). The specimens present features such as absence of anterior protolophule of the M1, open valleys in the M2 and m2, styls and metalophulid type II without type I and a reduced M3 (Markovic, 2008) that are not present in *C. soriae*. *Cricetodon meini* from Mühlbach (MN5) in Austria differs from *C. soriae* in its smaller size, it has neither anterior protolophule nor posterolophid on the M1 and it shows a reduced hypocone and paracone on the M3, and, the m3 is shorter than the m1 (Daxner-Höck, 2003).

***Cricetodon* from Vieux-Collonges**

Cricetodon aureus and *C. meini* were described in the middle Miocene French locality of Vieux-Collonges (Mein and Freudenthal, 1971b). The authors (1971b) stated that it was difficult to distinguish between the third molars of *C. aureus* and *C. meini* from Vieux-Collonges because of the large overlap in size. They studied the material stored at the Faculté des Sciences de Lyon, the Muséum d'Histoire Naturelle de Lyon and the NCB-Naturalis. They included detailed descriptions of each element of *C. aureus* and *C. meini*. However, the metrical and morphological descriptions of the third molars were reduced to the specimens included in the maxillas and mandibles. The authors pointed out: “[We have not tried to distinguish both forms on the third molars. We have only a few specimens on the mandible and maxilla that we know for sure is *C. aureus*]” (Mein and Freudenthal, 1971b: 26; translated from French) and “[We have not given the minimum for the third molars because of the difficulty of distinguishing the specimens of both species]” (Mein and Freudenthal, 1971b: 27; translated from French).

As we have pointed out before, in the discussion of the material from SOMN, we consider that the morphometrical features of the third molar have a high diagnostic value in *C. soriae*. First and second molars of *C. meini* are clearly smaller than *C. soriae*, whereas *C. aureus* is closer in size to *C. soriae* (Table 5.4). In order to evaluate the differences in size and morphology between the *C. aureus* and *C. soriae*, we have carried out a morphometrical analysis of 137 M3 elements identified as either *C. aureus* or *C. meini* stored at the RGM. Descriptive statistics presented on Table 5.4 for the M3 of *C. aureus* were calculated using the original measurements of Mein and Freudenthal (1971b) from the database of the NCB-Naturalis collections, but using the new specific assignation for the M3 based on the morphological criteria explained in this paper. The observed differences on the morphology of *C. aureus*, including the upper third molar are discussed here. The frequency and percentage of the most diagnostic features are presented on Figure 5.7.

In general, *C. aureus* presents the following features: absent entomesoloph on the M1; some M2 with styls and complete posterior ectolophs; ectomesolophid on the m1, with some m1 displaying exclusively type I or type II metalophulid; stylids on the m2; lingual anterolophid on the m3 (Mein and Freudenthal, 1971b) that are absent in *C. soriae*. Moreover, the mesolophs and mesolophids present in *C. aureus* are longer than the ones from Calatayud-Daroca material. The lower third molar shows a single short mesolophid that is never connected either to the metaconid or the entoconid, whereas *C. soriae* shows a long mesolophid that can be connected to either one of these cusps or to both. Our analysis determinates that the degree of hypsodonty in LUM11 is approximately 1.66 mm and about 1.20–1.30 mm in *C. aureus*. The M3 of *C. aureus* is smaller and less hypsodont than the specimens from Calatayud-Daroca. In addition, we observe that the neo-entoloph is absent in all the specimens (n=9) of *C. soriae* from Calatayud-Daroca, whereas

6 out 22 specimens of *C. aureus* show developed enamel spurs on both hypocone and protocone. This neo-entoloph can be disconnected (Fig. 5.7).

Regarding the posteroloph of the M3, we have described four character states for this structure: absent, short, connected to the metalophule and connected to the metacone with absent metalophule. Most of the M3 of *C. soriae* (87.5%) show a short posteroloph, a deep valley (posterosinus) delimited by a metalophule and a posteroloph is present. The M3 of *C. aureus* present short posteroloph in 45.5% of the specimens, although, this ridge is absent in 31.8% of the specimens. Furthermore, the posterosinus is not as deep as in the specimens from Spain (Fig. 5.7). In general, the mesolophs in *C. soriae* are less developed than in *C. aureus*. The M3 of *C. soriae* presents a well-developed hypocone and thus the posterior part of the teeth is elongated. On the other hand, *C. aureus* shows a more reduced hypocone, with a posterior part of its M3 also reduced.

Sarica-Filoreau (2002) proposed that the shape of the M3 can be expressed in terms of the Length/Width index. She recognized two groups: 1) rounded: Length/Width index between 1.00–1.10 and 2) and elongated: Length/Width index 1.10–1.20. For the Calatayud-Daroca material, most (five out of seven specimens) of the values for Length/Width index are approximately 1.10 (Fig. 5.7), which means that the external outline is close to an elongated form (Figs. 5.3. 5.4 to 5.6). In the case of *C. aureus*, the specimens show a large dispersion of index values (0.98–1.15), although we note that most of the specimens (16 out of 25) have an index of between 0.98 and 1.09. These specimens are characterized by a rounded shape.




FIGURE 5.7. Frequency and percentage (in parentheses) of the character states described for the M3 of *C. soriae* from the middle Aragonian of the Calatayud-Daroca Basin in comparison with *C. aureus* and *C. meini* from Vieux-Collonges.

FINAL REMARKS & CONCLUSIONS

The oldest occurrence of *C. soriae* in the Calatayud-Daroca basin is LUM11. As we pointed out above, De Bruijn et al. (1993) suggested the possibility of a new species in LUM11. The reasons they give to justify their proposal were: the intermediate size between the large species (*C. aureus* and *C. sansaniensis*) and the smaller species (*C. meini* and *C. jotae*); the low degree of hypsodonty and the combination of two features, the double protolophule and the lack of the anterior ectoloph on the M1.

After our morphological analysis of *C. soriae*, we can complete the information about the most important features of this species; these are absence of styli and stylids, low development of the mesoloph-ids, presence of anterior and posterior protolophules on the M1; absence of the anterior ectoloph on the M1 and M3; well-developed posteroloph on the M3; presence of a metalophulid II or I and II together; presence of incomplete metalophulid I; the m3 longer than the m1; the large size of the m3; long and double mesolophids on the m3.

Based partly of these aforementioned characters, we observe that there is a morphometrical resemblance between *C. soriae* and the species of *Cricetodon* from Anatolia. This idea is in agreement with the hypothesis of De Bruin et al. (1993) that suggests that the species present in LUM11 and in LP4A and LP4B, may be related to the Anatolian species of *Cricetodon*. They support that hypothesis based on a number of features such as the presence of anterior protolophule on the M1 and the fact that the m3 is longer than the m1. Furthermore, we find relevant to add to these common features, the presence of an incomplete metalophulid I on the m1 that we described in *C. soriae*. Moreover, De Bruijn et al. (1993) observed this incomplete metalophulid I on the m1 on the species from Anatolia such as *C. versteegi* and *C. tobieni*.

It is worth noting that any further study on *Cricetodon* must include a cladistic study in order to evaluate whether *C. soriae* share a common ancestry with the species from Asia Minor, since e.g. the phylogenetic study by Bi (2005) did not include a number of species, *C. soriae* among them.

The first occurrence of *C. soriae* is coincident with the faunal dispersion event called *Conohyus/Pliopitechus* described around 14 Ma ago (Van der Made, 1999). Several studies indicate the initiation of an episode of global cooling and aridity (Zachos et al., 2001; Böhme et al., 2011) corresponding with this period of time. For the Iberian Peninsula, the paleoclimatical context

corresponds to a cooling event and a decrease in the amount of precipitation (Van der Meulen and Daams 1992; Domingo et al., 2009a; Böhme et al., 2011). Such changes on the environmental conditions suggest that the Iberian Peninsula might be an optimum region for the establishment of *Cricetodon* representatives like *C. soriae*.

CONCLUSIONS

Both the morphological and metrical features of the studied material lead us to identify it as *Cricetodon soriae*. This species is the oldest representative of the genus in the Iberian Peninsula. The number of localities containing *C. soriae* increases by eight and thus the species can be found not only in the Madrid Basin but also in Calatayud-Daroca. The metrical and morphological analysis performed here adds to the information about the size and morphology of the scarce material from the type locality (SOMN) and allows us to emend the diagnosis of *C. soriae*. Moreover, detailed comparisons with the rest of the species of *Cricetodon* including the first representatives of *Cricetodon* in central and eastern Europe have been performed here. In order to evaluate the differences with *C. aureus*, we firstly have studied the type material of Vieux-Collonges and then identified the upper third molars as belonging to two species *C. aureus* and *C. meini*. Finally, we found several morphological features that *C. soriae* shares with the Anatolian species of *Cricetodon*. The first occurrence of *C. soriae* is coincident with a mammal migration event across Europe. This migration could be related to an environmental change into a period of cooling and an increase in the aridity of the Iberian Peninsula.

	Locality	Length					Width				L/W
		Min	Mean	Max	N		Min	Mean	Max	N	
M1	LUM19		3.16		1			2.07		1	1.53
	LUM11		3.30		1		2.13	2.14	2.14	2	1.54
M2	LUM20		2.52		1			2.30		1	1.10
	LP4BA		2.59		1			2.17		1	1.19
	LUM11		2.47		1			2.11		1	1.17
M3	LUM20	2.19	2.22	2.24	2		1.90	1.92	1.95	2	1.16
	LP4C		2.09		1			1.93		1	1.08
	LUM19		2.03		1			1.87		1	1.09
	LP4BA	1.93	1.96	1.99	2		1.88	1.94	1.99	2	1.01
	LUM11		2.24		1					1	
m1	LUM20		2.74		1			1.94		1	1.41
	LP4BA							1.71		1	
	LUM11		2.65		1			1.82		1	1.46
m2	LP4B		2.55		1			1.96		1	1.30
	LP4A							2.03		1	
	LUM16		2.38		1			1.94		1	1.23
m3	LUM11	2.49	2.53	2.57	3		1.93	2.04	2.14	2	1.24
	LUM20		2.78		1			2.20		1	1.26
	LUM19							1.98		1	
	LP4B		2.25		1			1.88		1	1.20
	LUM16	2.53	2.63	2.73	2		1.86	1.94	2.02	2	1.36
	LUM11	2.64	2.69	2.75	3		1.96	2.02	2.08	3	1.33

TABLE 5.3. Length and width of the upper and lower molars taken of *Cricetodon soriae* from Calatayud-Daroca. Units are given in mm. Abbreviations: Min, minimum values; Max, maximum values; N, number of specimens; L/W, Length/Width index.

	Species	Length					Width				L/W
		Min	Mean	Max	N		Min	Mean	Max	N	
M1	<i>Cricetodon klariankae</i>	3.30	3.39	3.55	8		2.15	2.21	2.27	9	1.53
	<i>Cricetodon sansaniensis</i>	2.99	3.38	3.68	34		2.00	2.19	2.35	33	1.54
	<i>Cricetodon albanensis</i>	3.18	3.48	3.77	50		1.99	2.15	2.31	50	1.62
	<i>Cricetodon aureus</i>	3.19	3.34	3.67	13		1.98	2.12	2.35	13	1.57
M2	<i>Cricetodon klariankae</i>	2.65	2.80	2.92	11		1.85	2.03	2.17	11	1.38
	<i>Cricetodon sansaniensis</i>	2.26	2.56	2.77	26		1.94	2.15	2.41	26	1.19
	<i>Cricetodon albanensis</i>	2.44	2.61	2.77	50		1.92	2.06	2.20	50	1.20
	<i>Cricetodon aureus</i>	2.45	2.55	2.65	8		2.01	2.09	2.18	8	1.22
M3	<i>Cricetodon klariankae</i>	1.76	1.91	2.05	11		1.62	1.76	1.90		1.09
	<i>Cricetodon sansaniensis</i>	1.91	2.09	2.31	26		1.73	1.96	2.16	23	1.07
	<i>Cricetodon albanensis</i>	1.94	2.11	2.30	50		1.77	1.92	2.05	50	1.10
	<i>Cricetodon aureus</i>	1.97	2.10	2.23	25		1.88	1.97	2.07	25	1.07
m1	<i>Cricetodon klariankae</i>	2.65	2.74	2.80	13		1.67	1.79	1.95	13	1.53
	<i>Cricetodon soriae</i>		2.65					1.86		1	1.42
	<i>Cricetodon sansaniensis</i>	2.59	2.86	3.08	29		1.74	1.89	2.08	29	1.51
	<i>Cricetodon albanensis</i>	2.70	2.94	3.20	50		1.76	1.87	1.97	50	1.57
	<i>Cricetodon aureus</i>	2.61	2.73	2.89	12		1.73	1.81	1.90	12	1.51
m2	<i>Cricetodon klariankae</i>	2.45	2.69	2.87	16		1.75	1.93	2.05	16	1.39
	<i>Cricetodon soriae</i>		2.58					2.05		1	1.26
	<i>Cricetodon sansaniensis</i>	2.55	2.71	2.89	25		1.93	2.14	2.37	32	1.27
	<i>Cricetodon albanensis</i>	2.45	2.65	2.86	50		1.90	2.06	2.23	50	1.29
	<i>Cricetodon aureus</i>	2.50	2.61	2.65	17		1.91	2.03	2.21	17	1.29
m3	<i>Cricetodon klariankae</i>	2.32	2.41	2.50	9		1.75	1.85	2.00	9	1.30
	<i>Cricetodon soriae</i>	2.37	2.50	2.64			2.03	2.08	2.12	2	1.20
	<i>Cricetodon sansaniensis</i>	2.39	2.65	3.00	20		1.81	2.02	2.20	20	1.31
	<i>Cricetodon albanensis</i>	2.49	2.64	2.80	50		1.85	1.98	2.10	50	1.25
	<i>Cricetodon aureus</i>	2.19	?	2.77	7		1.75	?	2.16	7	—

TABLE 5.4. Length and width of the upper and lower molars of large species of *Cricetodon*. Units are given in mm. Measurements for *C. sansaniensis*, *C. klariankae* and *C. albanensis* the M1, M2, m1, m2 and m3 of *C. aureus* and *C. soriae* from SOMN given by Maridet and Sen, 2012; Hír, 2007; Mein and Freudenthal, 1971a, 1971b and Hernández Fernández et al., 2006. Descriptive statistics presented for M3 of *C. aureus* was calculated using the dataset of the type series included in the collection database of the NCB-Naturalis, Leiden (The Netherlands), applying the new criteria of present paper to classify the M3. Abbreviations: Min, minimum values; Max, maximum values; N, number of specimens; L/W, Length/Width index.

LITERATURE CITED

- Abdul Aziz, H., M. Böhme, A. Rocholl, J. Prieto, J. R. Wijbrans, V. Bachtadse, and A. Ulbig. 2010. Integrated stratigraphy and $^{40}\text{Ar}/^{39}\text{Ar}$ chronology of the Early to Middle Miocene Upper Freshwater Molasse in western Bavaria (Germany). *International Journal of Earth Sciences* 99:1859–1886.
- Agustí, J. 1990. The Miocene Rodent Succession in Eastern Spain: A zoogeographical appraisal; pp. 375–404 in E. H. Lindsay, V. Fahlbusch, and P. Mein (eds.), a zoogeographical, appraisal in European Neogene Mammal Chronology. NATO ASI Series. Series A: Life Sciences, 180. Plenum Press, New York.
- Agustí, J., S. Moyà-Solà, and J. Pons Moyà. 1984. Mammal distribution dynamics in the eastern margin of the Iberian Peninsula during the Miocene. *Paléobiologie continentale* 14:33–46.
- Alberdi, M. T., and B. Azanza. 1997. Comentarios al artículo Magnetostratigrafía preliminar de los materiales pliocenos de la cubeta de Villarroja (Sierra de Cameros, La Rioja), de E. Pueyo Morer, A. Muñoz Jiménez y J. M. Parés (1996). *Geogaceta* 22:7–10.
- Álvarez-Sierra, M. A. 1987. Estudio sistemático y bioestratigráfico de los Eomyidae (Rodentia) del Oligoceno superior y Mioceno inferior español. *Scripta Geologica* 86:1–207.
- Álvarez-Sierra, M. A., J. Civis, A. Corrochano, R. Daams, C. J. Dabrio, E. García, A. González, N. López Martínez, R. Mediavilla, R. Rivas Carballo, and M. F. Valle. 1990. Un estratotipo del límite Aragoniense-Vallesiense (Mioceno medio-Mioceno superior) en la sección de Torremormojón (Cuenca del Duero, Provincia de Palencia). *Studia Geologica Salmanticensia. Actas de Paleontología Volumen Extraordinario* 57–64.
- Argyropulo, A. I. 1938. On the fauna of Tertiary Cricetidae of the USSR. *Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS* 20:223–226.
- Baudelot, S. 1972. Etude des chiroptères, insectivores et rongeurs du Miocene de Sansan, Toulouse. Ph.D. dissertation, Université Paul Sabatier de Toulouse, Toulouse, 364 pp.
- Bi, S. 2005. Evolution, systematics and functional anatomy of Cricetodontini (Cricetidae, Rodentia, Mammalia) from the Northern Juggar Basin, northwestern China. Ph.D. dissertation,

- Howard University, Washington, 156 pp.
- Boon, E. 1991. Cricetiden und Sciuriden der Oberen Süßwassermolasse. Ph.D. dissertation, Fakultät für Geowissenschaften der Ludwig-Maximilians-Universität, München, 159 pp.
- Bowdich, T. E. 1821. An Analysis of the Natural Classification of Mammals for the Use of Students and Travelers. J. Smith, Paris, 115 pp.
- Böhme, M., M. Winklhofer, and A. Ilg. 2011. Miocene precipitation in Europe: Temporal trends and spatial gradients. *Palaeogeography, Palaeoclimatology, Palaeoecology* 304:212–218.
- Cuenca Bescós, G. 1988. Revisión de los Sciuridae del Aragoniense y del Ramblense en la fosa de Calatayud-Montalbán. *Scripta Geologica* 87:1–116.
- Daams, R., and M. Freudenthal. 1981. Aragonian: the stage concept versus Neogene mammal zones. *Scripta Geologica* 62:1–17.
- Daams, R., M. Freudenthal, and A. Van de Weerd. 1977. Aragonian, a new stage for continental deposits of Miocene age. *Newsletters of Stratigraphy* 6:42–55.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, and W. Krijgsman. 1999a. Aragonian stratigraphy reconsidered and a re-evaluation of the middle Miocene mammal biochronology in Europe. *Earth and Planetary Science Letters* 165:287–294.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999b. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.
- Daams, R., L. Alcalá, M. A. Álvarez-Sierra, B. Azanza, J. A. Van Dam, A. J. Van der Meulen, J. Morales, M. Nieto, P. Peláez-Campomanes, and D. Soria. 1998. A stratigraphical framework for Miocene (MN4–MN13) continental sediments of Central Spain. *Comptes rendus de l'Académie des sciences, Séries IIA* 327:625–631.
- Daxner-Höck, G. 2003. *Cricetodon meini* and other rodents from Mühlbach and Grund, Lower Austria (Middle Miocene, late MN5). *Annalen des Naturhistorischen Museums in Wien* 104A:267–291.
- De Bruijn, H., and G. Saraç. 1992. Early Miocene rodent faunas from the Eastern Mediterranean area. Part II. *Mirabella* (Paracricetodontinea, Muroidea). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 95:25–40.
- De Bruijn, H., and E. Ünay. 1996. On the evolutionary history of the Cricetodontini from Europe

- and Asia Minor and its bearing on the reconstruction of migrations and the continental biotope during the Neogene; pp. 227–234 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Sarac, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a discussion of the evolutionary history of the Cricetodontini. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 96:151–216.
- De Bruijn, H., R. Daams, V. Fahlbusch, G. Daxner-Höck, L. Ginsburg, P. Mein, and J. Morales. 1992. Report of the RCMNS working group on fossil mammals, Reisenburg 1990. *Newsletters of Stratigraphy* 26:65–118.
- Domingo, L., J. Cuevas-González, S. T. Grimes, M. Hernández Fernández, and N. López-Martínez. 2009. Multiproxy reconstruction of the palaeoclimate and palaeoenvironment of the Middle Miocene Somosaguas site (Madrid, Spain) using herbivore dental enamel. *Palaeogeography, Palaeoclimatology, Palaeoecology* 272:53–68.
- Fischer, G. 1817. *Adversaria zoologica*, Mémoires de la Société Impériale des Naturalistes de Moscou 1917:357–472.
- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mittelspaniens und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 69:298–317.
- Garcés, M., W. Krijgsman, P. Peláez-Campomanes, M. A. Álvarez-Sierra, and R. Daams. 2003. *Hipparion* dispersal in Europe: magnetostratigraphic constraints from the Daroca area (Spain). *Coloquios de Paleontología Volumen Extraordinario* 1:171–178.
- García Moreno, E. 1988. The Miocene rodent biostratigraphy of the Duero basin (Spain): a proposition for a new Aragonian/Vallesian limit. *Paleontologia i Evolució* 22:103–112.
- García-Paredes, I. 2006. Patrones evolutivos de los Gliridae (Rodentia, Mammalia) del Mioceno inferior y medio del área tipo del Aragoniense (Cuenca de Calatayud–Montalbán). Ph.D. dissertation, Universidad Complutense de Madrid, Madrid, 676 pp.
- García-Paredes, I., P. Peláez-Campomanes, and M. A. Álvarez-Sierra. 2009. Gliridae (Rodentia, Mammalia) with a simple dental pattern: a new genus and new species from the European Early and Middle Miocene. *Zoological Journal of the Linnean Society* 157:622–652.

- García-Paredes, I., P. Peláez-Campomanes, and M. A. Álvarez-Sierra. 2010. *Microdyromys remerti*, sp. nov., a new Gliridae (Rodentia, Mammalia) from the Aragonian Type Area (Miocene, Calatayud-Montalbán Basin, Spain). *Journal of Vertebrate Paleontology* 30:1594–1609.
- Gómez Cano, A. R., M. Hernández Fernández, and M. A. Álvarez-Sierra. 2011. Biogeographic provincialism in rodent faunas from the Iberocitanian Region (southwestern Europe) generates severe diachrony within the Mammalian Neogene (MN) biochronologic scale during the Late Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 307:193–204.
- Hernández-Ballarín, V., A. Oliver, and P. Peláez-Campomanes. 2011. Revisión de las asociaciones de mamíferos del tránsito Aragoniense medio y superior de la Cuenca de Madrid; pp. 183–172 in A. Pérez-García, F. Gascó, J. M. Gasulla, and F. Escaso (eds.), *Viajando a Mundos Pretéritos*. Ayuntamiento de Morella, Morella, Castellón.
- Hernández Fernández, M., J. A. Cárdena, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőtárkány-Felnémet (Northern Hungary). *Fragmenta Paleontologica Hungarica* 24–25:16–24.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254:177–235.
- Kordos, L. 1986. Upper miocene hamsters (Cricetidae, Mammalia) of Hasznos and Szentendre. *Magyar Allami Földtani Intézet Jelentése Az* 1984:523–553.
- Klein Hofmeijer, G., and H. De Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece) Part 8: The Cricetidae. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 91:185–204.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d'une récapitulation des diverses espèces d'animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d'autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. J. A. Portes, Auch, 41 pp.
- López-Guerrero, P., M. A. Álvarez-Sierra, R. López-Antoñanzas, A. Oliver Pérez, and P. Peláez-Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior

- (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). *Palaeontologica Nova* 8:259–271.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra, R. López-Antoñanzas, and A. Oliver. 2009. *Cricetodontini* (Rodentia, Mammalia) from the upper Aragonian and lower Vallesian of the Toril-Nombrevilla section (Middle and Upper Miocene, Calatayud-Daroca basin, Zaragoza, Spain). *Journal of Vertebrate Paleontology* 29(3, supplement):161A.
- López-Guerrero, P., A. Oliver Pérez, M. A. Álvarez-Sierra, I. García Paredes, and P. Peláez-Campomanes. 2007. El registro de los *castoridae* (Rodentia, Mammalia) del Aragoniense y Vallesiense (Mioceno Medio y Superior) de las cuencas centrales españolas; pp. 118–119 in J.C. Braga, A. Checa, and M. Company (eds.), *XXIII Jornadas de la Sociedad Española de Paleontología* (Caravaca de la Cruz 3–6 de Octubre de 2007). Libro de resúmenes. Instituto Geológico y Minero de España & Universidad de Granada, Granada.
- Luis, A., and J. M. Hernando. 2000. Los microvertebrados fósiles del Mioceno Medio de Somosaguas Sur (Pozuelo de Alarcón, Madrid, España). *Coloquios de Paleontología* 51:69–86.
- Maridet, O., and S. Sen. 2012. Les *Cricetidae* du gisement de Sansan; pp. 29–64 in S. Peigné and S. Sen (eds.), *Les mammifères du gisement de Sansan. Mémoires du Muséum national d'Histoire naturelle*, Paris.
- Markovic, Z. 2008. Rodents of middle Miocene localities of Lazarevac village and Bele vode (central Serbia). *Bulletin of the Natural History Museum* 1:79–98.
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des *Cricetidae* (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les *Cricetidae* (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Morales, J., M. Nieto, P. Peláez-Campomanes, D. Soria, M. A. Álvarez-Sierra, L. Alcalá, L. Amezua, B. Azanza, E. Cerdeño, R. Daams, S. Fraile, J. Guillem, M. Hoyos, L. Merino, I. de Miguel, R. Monparler, P. Montoya, B. Pérez, M. J. Salesa, and I. M. Sánchez. 1999. Vertebrados continentales del Terciario de la cuenca de Loranca (Provincia de Cuenca); pp. 237–260 in E. Aguirre, and I. Rábano (eds.), *La Huella del Pasado: Fósiles de Castilla-La Mancha*. Junta de Comunidades de Castilla-La Mancha, Toledo.
- Oliver, A., P. López-Guerrero, I. García-Paredes, M. A. Álvarez-Sierra, and P. Peláez-Campomanes. 2009. Evolution of *Megacricetodon* tooth pattern through geometric morphometric analysis. *Journal of Vertebrate Paleontology* 29(3, supplement):158A.
- Peláez-Campomanes, P. 2001. Revision of the Aragonian (Miocene) *Atlantoxerus* (Sciuridae).


- Journal of Paleontology 75:418–426.
- Peláez-Campomanes, P., J. Morales, M. A. Álvarez-Sierra, B. Azanza, S. Fraile, I. García Paredes, M. Hernández Fernández, E. Herráez, M. Nieto, B. Pérez, V. Quiralte, M. J. Salesa, I. M. Sánchez, and D. Soria. 2003. Updated biochronology of the Miocene mammal faunas from the Madrid basin (Spain); pp. 431–441 in J. W. F. Reumer and W. Wessels (eds.), *Distribution and Migration of Tertiary Mammals in Eurasia. A Volume in Honour of Hans de Bruijn*. Deinsea, Rotterdam.
- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.
- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata Palasiatica* 48:27–47.
- Rummel, M. 1995. *Cricetodon bolligeri* n. sp. ein neuer Cricetide aus dem Obermiozän von Petersbuch bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 35:109–123.
- Rummel, M. 1998. Die Cricetiden aus dem Mittel- und Obermiozän der Türkei. *Documenta naturae* 123:1–300.
- Rummel, M. 2000. Miocene Cricetodontini from Petersbuch near Eichstaett. The genus *Cricetodon* Lartet 1851. *Senckenbergiana Lethaea* 80:149–171.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde B* 311:1–6.
- Rummel, M., and D. Kálin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Mitelmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Sarica-Filoreau, N. 2002. Faunes de rongeurs neogenes et quaternaires des grabens d'Anatolie occidentale, systematique, biochronologie et implications tectoniques. Ph.D. dissertation, Muséum national d'Histoire naturelle, Paris, 327 pp.
- Schaub, S. 1925. Die hamsterartigen Nagetiere des Tertiärs und ihre lebenden Verwandten. *Abhandlungen der Schweizerischen paläontologischen Gesellschaft* 45:1–114.
- Sen, S., and E. Ünay. 1979. Sur quelques Cricetodontini (Rod.) du Miocène moyen d'Anatolie. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* 82:293–301.
- Sesé, C. 1977. Los cricétidos (Rodentia, Mammalia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno Cuaternario* 8:127–80.

- Simpson, G. C. 1945. The principles of classification and a classification of mammals. *Bulletin of the American Museum of Natural History* 85:1–350.
- Stehlin, H. G., and S. Schaub. 1951. Die Trigonodontie der simplicidentaten Nager. *Schweizerische paläontologische Abhandlungen* 67:1–385.
- Tobien, H. 1978. New species of Cricetodontini (Rod. Mam.) from the Miocene of Turkey. *Mainzer Geowissenschaftliche Mitteilungen Geologisches Landesamt Rheinland-Pfalz* 6:209–219.
- Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. Van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodent faunas of the Mio–Pliocene continental sediments of the Teruel-Alfambra region, Spain. *Utrecht Micropaleontological Bulletin, Special Publication* 2:1–217.
- Van der Made, J. 1999. Intercontinental relationship Europe–Africa and the Indian Subcontinent; pp. 359–364 in G. Rößner and K. Heißig (eds.), *The Miocene Land Mammals of Europe*. Verlag Dr. Friedrich Pfeil, München.
- Van der Meulen, A. J., and R. Daams. 1992. Evolution of Early–Middle Miocene rodent faunas in relation to long-term palaeoenvironment changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 93:227–253.
- Van der Meulen, A. J., P. Peláez-Campomanes, and R. Daams. 2003. Revision of medium-sized Cricetidae from the Miocene of the Daroca-Villafeliche area in the Calatayud-Teruel basin (Zaragoza, Spain). *Coloquios de Paleontología Volumen Extraordinario* 1:385–441.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, and P. Peláez-Campomanes. 2012. Updated Aragonian biostratigraphy: Small Mammal distribution and its implications for the Miocene European Chronology. *Geologica Acta* 10:1–24.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, P. López-Guerrero, V. Hernández-Ballarín, and P. Peláez-Campomanes. 2011. Biostratigraphy or Biochronology? Lessons from the Early and Middle Miocene small Mammal events in Europe. *Geobios* 44:309–321.
- Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.

- Zachos, J., M. Pagani, L. Sloan, E. Thomas, and K. Billups. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693.
- Ziegler, R., T. Dahlmann, J. W. F. Reumer, and G. Storch. 2005. Germany; pp. 61–98 in L. W. Van den Hoek Ostende, C. S. Doukas and J. W. F. Reumer (eds.), *The Fossil Record of the Eurasian Neogene Insectivores (Erinaceomorpha, Soricomorpha, Mammalia) Part I Scripta Geologica special issue, 5*. Leiden.

Chapter 6

C. sansaniensis, C. jotae



CRICETODON FROM THE MIDDLE AND LATE ARAGONIAN OF THE CALATAYUD-DAROCA BASIN: A TAXONOMICAL DESCRIPTION

“La formation de ces ectolophes est liée à un accroissement rapide de la taille de telle sorte qu'on pourrait être tenté de placer ces espèces au voisinage de C.sansaniensis, mais son origine à partir de C. cf. jotae paraît vraisemblable, aussi nous le considérons comme un groupe de la lignée moyenne ayant acquis secondairement une grande taille”

Pierre Mein & Mathijs Freudenthal

Resumen

6

Los yacimientos de fósiles del Aragoniense superior de la Cuenca de Calatayud-Daroca (Zaragoza, España) han proporcionado abundantes asociaciones de roedores; la mayoría de ellas contienen representantes de la tribu Cricetodontini. El presente estudio de las localidades de Mioceno medio de la Cuenca de Calatayud-Daroca muestra la presencia de dos especies diferentes: *Cricetodon sansaniensis* (presente en la biozona local F, MN6, Aragoniense medio) y *Cricetodon jotae* (presente en las biozonas locales G1, G2 y parte de la G3, MN6-MN7/8, Aragoniense medio y superior). Los dieciocho yacimientos estudiados se han datado entre 13,76 Ma y ~12,57 Ma. *Cricetodon sansaniensis* tiene ectolofos poco desarrollados y ambos metalofúlidos, I y II, en el mismo m1, así como mesolofos y mesolófidos que, de acuerdo con las tendencias descritas para las especies de Cricetodontini, son rasgos basales. La descripción detallada de *Cricetodon jotae* ha permitido enmendar su diagnosis. *Cricetodon jotae* muestra una combinación de caracteres basales (como el metalofúlido II) y caracteres derivados (como la ausencia de mesolofos; la presencia de ectolofos dobles; mesolófidos poco desarrollados o ausentes; ausencia de ectomesolófido y el anterolófido labial del m1 poco desarrollado o ausente). Este trabajo completa la revisión de las especies de *Cricetodon* presentes en Calatayud-Daroca.

Referencia

Este capítulo, con algunas modificaciones, recoge una parte de la publicación:

López-Guerrero, P., I., García-Paredes, M. A., Álvarez-Sierra, and P. Peláez-Campomanes. 2014. Cricetodontini from the Calatayud-Daroca Basin (Spain): a taxonomical description and up-to-date of their stratigraphical distributions. *Comptes Rendus Palevol.* en prensa.



Abstract

Upper Aragonian fossil sites from the Calatayud-Daroca Basin (Zaragoza, Spain) have yielded fairly large rodent assemblages; most of them contain representatives of the tribe Cricetodontini. The study of the localities from the middle Miocene of the Calatayud-Daroca Basin reveals that the fossils belong to two different species: *Cricetodon sansaniensis* (local biozone F, MN6, middle Aragonian) and *Cricetodon jotae* (local biozone G1, G2 and partly G3, MN6-MN7/8, middle-upper Aragonian). The age of the eighteen sites studied spans a time range from 13.76 Ma to ~12.57 Ma. *Cricetodon sansaniensis* has weak ectolophs and both metalophulids, I and II, in the same m1, and mesolophs/ids that, according with the trends described for Cricetodontini, are basal traits. The emended diagnosis of *C. jotae* is also presented here. *Cricetodon jotae* displays a combination of basal character (metalophulid II) and derivate characters (absence of mesolophs; presence of double ectolophs; weak or absent mesolophids; absence of ectomesolophid and weak or absent labial anterolophid of the m1). This work completes the revision of the species of *Cricetodon* from the Calatayud-Daroca Basin.

INTRODUCTION

In the Calatayud-Daroca Basin the tribe Cricetodontini is represented by the genera *Cricetodon* and *Hispanomys* (Álvarez-Sierra et al., 2003; López-Guerrero et al., 2008, 2009, 2013a, 2013b). Details of the oldest species from the middle and late Aragonian (middle Miocene), are well known since *Cricetodon soriae* López Martínez et al., in: Hernández Fernández et al., 2006 was revised by López-Guerrero et al. (2013a), and the new species *Cricetodon nievei* López-Guerrero et al., 2014 has been recently described. However, since the works of Mein and Freudenthal (1971a), the material of the species from the late Aragonian has not been reviewed. Therefore, in order to complete the knowledge of the Cricetodontini from the Calatayud-Daroca basin, the taxonomical and morphometrical information has to be updated. Therefore, in this paper, the material of *Cricetodon* found in the earliest part of the late Aragonian from the Calatayud-Daroca Basin is described and discussed. It includes the study of new material belonging to this genus from eighteen sites, which has been assigned to two different species. These fossil localities cover the local biozones F, G1, G2 and the lower part of the G3 (Daams et al., 1999), correlated with MN6 and lower part of the MN7/8.

The study of the localities from the middle Miocene of the Calatayud-Daroca Basin reveals that the material belong to two different species: *Cricetodon sansaniensis* Lartet, 1951 (local biozone F, MN6, middle Aragonian) and *Cricetodon jotae* Mein and Freudenthal, 1971a (local biozone G1, G2 and partly G3, MN6-MN7/8, middle-upper Aragonian).

Cricetodon sansaniensis was described in the French locality of Sansan and it is widespread across the central and southwestern European basins. This species has been related with *C. aureus* and *C. albanensis* by different authors (Mein and Freudenthal, 1971a; Maridet and Sen, 2012) due to its large size. *Cricetodon sansaniensis* was already cited as *Cricetodon* cf. *sansaniensis* in the Calatayud-Daroca Basin, in Armantes VI (Freudenthal, 1966; Mein and Freudenthal, 1971a). Since then, more fossils belonging to this species have been found in the Calatayud-Daroca Basin, which allows to improve the knowledgement of the morphological variability of this species in the basin.

Cricetodon jotae was described by Mein and Freudenthal (1971a). These authors refer to Freudenthal's studies (1963 and 1966) for the metrical information and the comparison with other species is briefly discussed. Although it is a common and abundant species in the Calatayud-Daroca Basin, detailed morphometric descriptions have been never provided in a systematic study.

Therefore, the main aims of this study are to characterize the representatives of *Cricetodon* from the late Aragonian through morphological and metrical analysis of the cheek teeth and to specify their stratigraphic distribution in the Calatayud-Daroca Basin.

In the case of *Cricetodon jotae*, we also provide the morphological study of all the assemblages from the Calatayud-Daroca Basin, in order to know the intraspecific variability. Using this information, we have compared *C. jotae* with other representatives of the genus present in Europe and Asia Minor.

MATERIAL & METHODS

We have studied 503 upper and lower molars belonging to 18 sites. Table 6.1 contains all the information about the different localities studied. Within the total sample, 89 specimens from five localities belong to *C. sansaniensis* Lartet, 1851 and the other 414 molars correspond to *C. jotae* Mein and Freudenthal, 1971a, recovered from 13 localities (Table 6.1). We observed directly the material of several species summarized in table 6.2. The information about the other species of Cricetodontini from the Calatayud-Daroca Basin has been found in López-Guerrero et al. (2008, 2009, 2013a, 2014). The numerical ages of the localities are after Daams et al. (1999) and Van Dam et al. (2006) and Van Dam et al. (in prep.). The nomenclature used is the one modified by López-Guerrero et al. (2013a). Length and width measurements for each specimen were taken (in mm) using a Nikon Measuroscope 10 microscope with digital micrometer, following the procedure suggested by López-Guerrero et al. (2013a). Tables 6.3 and 6.4 summarize the descriptive statistics for each locality. Figure 6.1 shows length and width scatter-diagrams of *C. sansaniensis* and the species with similar size (*C. klariankae* Hír, 2007; *C. engesseri* Rummel and Kälin, 2003; *C. cariensis* Sen and Ünay, 1979; *C. aureus* Mein and Freudenthal, 1971b; and *C. albanensis* Mein and Freudenthal, 1971a). Length and width scatter-diagrams for *C. jotae* and the species with similar size (*C. meini* Freudenthal, 1963 and *C. bolligeri*, Rummel, 1995) are presented in figure 6.2. The photographs from figures 6.3 and 6.4 were taken with an environmental Scanning Electron Microscope FEI Quanta 200 at the Museo Nacional de Ciencias Naturales-CSIC, Madrid (Spain) except for those from ARM7 and VT2C, taken with a digital camera attached to a Leica Microscope at Naturalis Biodiversity Center, Leiden (The Netherlands).

Institutional Abbreviations—**BSPG**, Bayerische Staatssammlung für Paläontologie und Geologie, Munich (Germany); **IAUU**, Instituut voor Aardwetenschappen, Utrecht University (The Netherlands); **MNCN**, Museo Nacional de Ciencias Naturales-CSIC, Madrid (Spain); **NHNB**, Muséum d'Histoire Naturelle de Bâle (Switzerland); **PM**, Páasztó Múzeum, Páasztó (Hungary); **RGM**, Naturalis Biodiversity Center, former Rijksmuseum van Geologie en Mineralogie, Leiden (The Netherlands); **UCM**, Facultad de Geología, Universidad Complutense de Madrid (Spain).

	Locality	Abb.	Collections	M1	M2	M3	m1	m2	m3	N
Age (Ma)	<i>Cricetodon sansaniensis</i>									
13.68	Valalto 2C	VT2C	RGM	3	5	5	3	—	4	20
13.70	Valalto 2B	VT2B	RGM	—	—	—	2	1	3	6
	Armantes 7	ARM7	IAUU	5	3	5	1	1	2	17
13.75	Las Umbrías 21	LUM21	MNCN	8	8	4	5	4	5	34
13.76	Las Umbrías 22	LUM22	MNCN	—	2	2	3	2	3	12
	Total			16	18	16	14	8	17	89
	<i>Cricetodon jotae</i>									
	Toril 1	TOR1	RGM	4	4	4	3	3	3	21
	Alcocer 3	AC3	RGM	—	2	—	2	—	—	4
	Alcocer 2	AC2	RGM	—	—	—	4	4	3	11
	Alcocer 1	AC1	RGM	1	—	1	—	1	—	3
	Villafeliche 9	VL9	RGM	1	2	2	2	3	2	12
	Arroyo del Val VI	ARV6	IAUU	19	10	15	19	24	22	109
13.08	Las Planas 5K	LP5K	RGM	3	5	1	5	2	—	16
13.16	Las Planas 5L	LP5L	RGM	—	2	—	3	—	1	6
	Borjas	BOR	RGM	6	4	2	5	4	3	24
	Manchones	MAN	IAUU/RGM	30	33	20	37	35	30	185
	Valalto 1	VT1	MNCN/RGM	1	—	—	1	1	2	5
13.55	Las Planas 5C	LP5C	MNCN	3	2	2	3	2	—	12
13.56	Las Planas 5B	LP5B	RGM	3	1	—	—	—	2	6
	Total			71	65	47	84	79	68	414

TABLE 6.1. Studied material of *Cricetodon sansaniensis* and *C. jotae* from the Calatayud-Daroca Basin. Numerical ages of the localities after Van Dam et al. (2006), localities, abbreviations for the localities, and institutions where the fossils are stored (see institutional abbreviations). Abbreviations: Abb, abbreviations for the localities; N, number of specimens.

Species	Type locality	Institution
<i>C. soriae</i>	Somosaguas (Spain)	UCM
<i>C. bolligeri</i>	Petersbuch 10 (Germany)	BSPG
<i>C. fandli</i>	Gratkorn (Austria)	BSPG
<i>C. sansaniensis</i>	Sansan (France)	MHNB
<i>C. aureus</i>	Vieux-Collonges (France)	RGM
<i>C. meini</i>	Vieux-Collonges (France)	RGM
<i>C. albanensis</i>	La Grive-Saint Alban (France)	RGM
<i>C. klariankae</i>	Felsőtárkány-Felnémet	PM
<i>C. jotae</i>	Manchones (Spain)	IAUU

TABLE 6.2. Information of the material directly studied for comparisons (see institutional abbreviations).

SYSTEMATIC PALEONTOLOGY

Order RODENTIA Bowdich, 1821

Family CRICETIDAE Fischer, 1817

Subfamily CRICETODONTINAE Stehlin and Schaub, 1951

Tribe CRICETODONTINI Simpson, 1945

Genus *CRICETODON* Lartet, 1851

CRICETODON SANSANIENSIS Lartet, 1851

(Figs 6.1; 6.3; Tables 6.1–6.3)

Material

Original Type Locality—Sansan, France, (MN6, middle Miocene)

Stratigraphic Range in the Calatayud-Daroca Basin—local biozone F (MN6, upper Aragonian, middle Miocene).

Geographical Range—central and southwestern Europe.

Diagnosis

Emended diagnosis (Maridet and Sen, 2012: 30)—“Large-sized *Cricetodon*; M1 with an anterocone either undivided or divided at its top, but complete on the 3/4 of the crown height. In M1, ectolophs are absent in half of the specimens, weakly developed on the posterior surface of the anterocone and/or paracone in the other half of the specimens. The mesocone presents a triangular shape with a short mesoloph in 1/4 of the M1s, slightly more frequent in M2 and M3. The anterior surface of both the paracone and the metacone are notched by a vertical groove. The m1 has an undivided anteroconid; the metaconid has a posterior metalophulid in more than half

of the specimens, double or only anterior for the other specimens. Mesolophid absent or rare in m1, more frequent in m2 and m3, but always short. M1 with four roots, and m2 with 2 roots, with the posterior one either with a double pulp cavity or divided at its extremity for about half of the specimens.”.

CRICETODON SANSANIENSIS FROM THE CALATAYUD-DAROCA BASIN

Description of the Studied Material

M1—This molar has four roots. A shallow groove on the anterior wall divides the anterocone in two parts, but not covering the entire height of the cusp (Fig. 6.3.A; 6.3.D). The anterolophule is disconnected in two teeth from ARM7 (Fig. 6.3.A). The protolophule is posteriorly directed; it is anterior in two specimens (1/5 in LUM21 and 1/5 in ARM7); it is present but disconnected in one specimen from LUM21. The anterior ectoloph is frequently absent (2/3 in VT2C; 3/4 in LUM21, and 2/4 in ARM7); it is composed by the posterior ectoloph of the anterocone (Fig. 6.3.A) in all teeth but one from VT2C, which is double, composed by the posterior ectoloph of the anterocone and the anterior ectoloph of the paracone. The anterior ectoloph is never complete. The posterior ectoloph is frequent (Fig. 6.3.D); it is absent only in LUM21 (1/4). It is always composed by the posterior ectoloph of the paracone, and never complete. The mesoloph is mostly absent (Fig. 6.3.A); it is short (1/2 in VT2C and 1/4 in ARM7; Fig. 6.3.D) or reduced to an enlargement of the enamel, called incipient, (3/4 in ARM7 and 3/4 in LUM21). The anterosinus is open in one specimen from LUM21; closed by a cingulum (4/4 in ARM7; 3/4 in LUM21 and 1/3 in VT2C; Fig. 6.3.A; 6.3.D) or by a style (2/3 in VT2C). The mesosinus is open (1/4 in LUM21) or blocked by a cingulum (Fig. 6.3.A; 6.3.D). The sinus can be either open (3/4 in LUM21; 1/4 in ARM7; Fig. 6.3.D), closed by a cingulum (1/2 in VT2C; 1/4 in ARM7 and 1/4 in LUM21; Fig. 6.3.A) or blocked by a style (1/4 in ARM7 and 1/2 in VT2C).

M2—The anterior ectoloph is reduced (Fig. 6.3.E; 6.3.G) and, generally, lower than the anteroloph; it can be absent (1/3 in ARM7). The posterior ectoloph is composed by the posterior ectoloph of the paracone and it is not complete (Fig. 6.3.E; 6.3.G); it is rarely absent (1/2 in LUM22 and 1/3 in ARM7). The mesoloph is mostly absent (Fig. 4.3.E; 4.3.G); it can be short (1/4 in VT2C) or incipient (2/6 in LUM21 and 1/4 in VT2C). The labial posteroloph is always present and the lingual posteroloph is absent in LUM22 and in VT2C (1/4). The anterosinus is always closed by a cingulum. The protosinus is weak but distinguishable in 1/7 in LUM21 (Fig. 6.3.G). The mesosinus, which is well developed, can be open (1/2 in LUM22; 2/6 in LUM21, and 3/3 in ARM7) or closed by a cingulum (4/4 in VT2C, 1/2 in LUM22 and 4/6 in LUM21; Fig.

6.3.E; 6.3.G). The sinus can be open (3/4 in VT2C; 1/2 in LUM22; 2/6 in LUM21, and 1/2 in ARM7; Fig. 6.3.E; 6.3.G) or closed, either by a cingulum (1/2 in LUM22; 4/6 in LUM21, and 1/2 in ARM7) or by a style (1/4 in VT2C).

M3—Both labial and lingual anterolophs are present, but the lingual branch is weaker than the labial one (Fig. 6.3.C; 6.3.F; 6.3.H). The anterior ectoloph is absent (Fig. 6.3.C; 6.3.F; 6.3.H). All specimens except for 2/4 from ARM7 have a posterior ectoloph; it is formed by the posterior ectoloph of the paracone (2/4 ARM7; 1/3 in LUM21; 2/2 in LUM22, and 2/4 in VT2C; Fig. 6.3.C; 6.3.F; 6.3.H). The posterior ectoloph is composed by both the posterior ectoloph of the paracone and the anterior ectoloph of the metacone in some cases (1/4 in LUM21 and 2/4 in VT2C); it is absent in two out of four specimens from ARM7. The posterior ectoloph is complete in two molars (1/4 in VT2C and 1/4 in LUM21). Most of the specimens have a mesoloph; it is absent in two teeth (1/5 in VT2C and 1/2 in LUM22). The length of the mesoloph is variable; it can be short (1/5 in VT2C and 3/4 in ARM7; Fig. 6.3.H), about half the distance between longitudinal ridge and labial border (3/5 in VT2C; 2/4 in LUM21, and 1/4 in ARM7; Fig. 6.3.C), or reaching the labial border (1/2 in LUM22 and 2/4 in LUM21; Fig. 6.3.F). An the enamel-coated valley can be present (1/4 in VT2C; 1/2 in LUM22 and 2/4 in LUM21). The axioloph is disconnected (2/3 in VT2C) or continuous (Fig. 6.3.C; 6.3.F; 6.3.H). The labial posteroloph is well developed but short and it is never connected to the metalophule or the metacone (Fig. 6.3.C; 6.3.F; 6.3.H); it is absent in 2/4 specimens from LUM21.

m1—Only the metalophulid I is found in some specimens (1/2 in LUM2; 1/3 in LUM22; Fig. 6.3.I; 6.3.L); both metalophulids, I and II, are present together in the remaining teeth. The metalophulid I is incomplete in one case (LUM21; Fig. 6.3.I). The mesolophid is rare; only three teeth from LUM21 have one; they are incipient (Fig. 6.3.I), short and medium, respectively. The ectomesolophid is present in one specimen from LUM22. The entoconid spur and the labial posterolophid are absent. The mesosinusid can be open or blocked; it is closed by a low cingular ridge (1/2 in VT2C and 1/3 in LUM22; Fig. 6.3.I) or by a stylid (1/2 in VT2B). The sinusid is blocked by a cingular ridge (2/2 in VT2C; 1/2 in VT2B; in ARM7 and 2/3 in LUM21) or it remains open (Fig. 6.3.I; 6.3.L). The posterosinusid is blocked by a cingular ridge in one specimen (1/2 in VT2B); it is open in the rest of the teeth.

m2—The labial branch of the anterolophid is well developed and it reaches the protoconid (Fig. 6.3.J; 6.3.M). Only the tooth from ARM7 possesses a lingual anterolophid. A short mesolophid (Fig. 6.3.J; 6.3.M) is usually present; it is absent in one tooth. In some specimens (4/5 in LUM21), the mesolophid is curved and projected towards the metaconid, reaching it. One fossil from LUM22 shows an ectomesolophid (Fig. 6.3.J). There is no anterosinusid and the mesosinusid is always blocked by a cingular ridge. The sinusid is blocked by a cingulid; in one out of four teeth from LUM21 this cingular ridge is split in two crests. The lingual posterosinusid is

open (2/2 in LUM22 and 2/4 in LUM21), blocked by a cingular ridge (2/4 in LUM21), or by the joint of the lingual posterolophid and the entoconid (1/1 in ARM7).

m3—The labial branch of the anterolophid is long and it reaches the protoconid (Fig. 6.3.K; 6.3.N). The lingual branch of the anterolophid is present in two out of four molars from LUM21. The mesolophid can be short (1/4 in VT2C; 1/3 in VT2B; 1/2 in LUM22; 3/5 in LUM21, and 2/2 in ARM7; Fig. 6.3.N) or have medium length (2/4 in VT2C; 1/2 in LUM22, and 1/5 in LUM21; Fig. 6.3.K). In two fossils (1/3 in VT2B and 1/5 in LUM21) the mesolophid is double and short. In one tooth (1/3 in LUM22) it is connected with the metaconid, being absent in the remaining specimens (1/4 in VT2C and 1/3 in VT2B). The lingual posterolophid can be constricted behind the hypoconid (Fig 6.3.K); its lingual part is cuspidate and displays a weak valley on the posterior wall. The sinusid is blocked, in most cases, by a low cingular ridge or by a stylid (1/2 in VT2B); it remains open in two specimens (1/2 in VT2B and 1/2 in LUM21; Fig 6.3.K; 6.3.N). The mesosinusid is open or blocked by a low cingular ridge. A cingulid is also present at the end of the lingual posterosinusid (1/2 in LUM22 and 1/4 in LUM21). The lingual posterolophid projects towards the entoconid, in some teeth (2/2 in VT2C; 2/2 in VT2B; 2/2 in ARM7, and 1/4 in LUM21). In the remaining specimens (1/2 in LUM22 and 2/4 in LUM21), the lingual posterosinusid is open (Fig. 6.3.K; 6.3.N).

Discussion

The described specimens fit within the size range of *Cricetodon sansaniensis* Lartet, 1851 (Fig. 6.1), one of the largest species of *Cricetodon* recovered until now. However, the measurements of the m1 are slightly smaller than the mean value of those from Sansan (Fig. 6.1). The incompletely divided anterocone, the presence of anterior ectoloph, the labially curved anterior ectoloph, and the poorly developed or absent posterior ectoloph on the M1 are characteristic traits of *C. sansaniensis* from Sansan (Baudelot, 1972; Maridet and Sen, 2012). These characters are also found in the studied fossils. In addition, the large mesoloph of the M3 and the presence of metalophulid I and II together in the m1 are features commonly found in the specimens from Sansan (Baudelot, 1972; Maridet and Sen, 2012) and also present in the Calatayud-Daroca Basin. The studied material has no lingual spur of the anterolophule, neither two crests at the edge of the anterosinus and mesosinus of the M1, and the material from Sansan has both structures. Apart from that, one of the m1 from LUM22 displays ectomesolophid, whereas this feature is shown by 11 out of 29 specimens from Sansan (Maridet and Sen, 2012); nevertheless, the relative frequency of this character in both areas cannot be compared due to the small sample size of the assemblages from Calatayud-Daroca with respect to that from Sansan.

Part of the studied material were already cited in the Calatayud-Daroca Basin, that from

ARM7 as *Cricetodon* cf. *sansaniensis* in Freudenthal (1966) and Mein and Freudenthal (1971a), and the molars from VT2C and VT2B as *Cricetodon* sp. in Daams and Freudenthal (1981). The fossils from LUM21 and LUM22 are described and assigned for the first time. *Cricetodon sansaniensis* is widespread across the central and southwestern European basins. Apart from Sansan and the localities of the present study, it has recorded in another five fossil sites (Bolliger, 1994; Sach, 1999; Ginsburg and Bulot, 2000; Kälin and Kempf, 2009). Besides, *Cricetodon* cf. *sansaniensis* has been reported 12 times in the literature, although some of them might be not refer to *C. sansaniensis* (García Moreno, 1988; Bolliger, 1994; Sach, 1999; Ginsburg and Bulot, 2000; Rummel, 2000; Rummel and Kälin, 2003; Seehuber, 2008; Kälin and Kempf, 2009). Apart from Niderwis, *C. sansaniensis* is recorded in other Swiss localities from the local biozone Öschgraben ranging from 14.2 Ma to 14.1 Ma (Kälin and Kempf, 2009); therefore, the youngest known record is from Switzerland. The latest known record of the species is uncertain. The assignment to *C. sansaniensis* of the fossils described by García Moreno (1988) in the *Megacricetodon lopezae* zone from the Duero Basin (~local biozone G in Calatayud-Daroca Basin) is questionable and the specimens are under revision. Not considering the Duero record, the youngest occurrence is in the Calatayud-Daroca Basin, in which *C. sansaniensis* is present in the biozone F, between 13.76 Ma and 13.68 Ma (Van Dam et al., 2006; Daams et al., 1999).

Cricetodon sansaniensis from the Calatayud-Daroca Basin has some character states (such as weak ectolophs and both metalophulids, I and II, on the same m1, and mesolophs/ids) which places it among the basal representatives of the tribe according to the trends described for Cricetodontini (Agustí, 1982; De Bruijn et al., 1993; Hír, 2007; Maridet and Sen, 2012).

Due to its large size, *Cricetodon sansaniensis* has been included by some authors (Mein and Freudenthal, 1971a; Maridet and Sen, 2012) in the anagenetic lineage *C. aureus*–*C. sansaniensis*–*C. albanensis*. According to Mein and Freudenthal (1971a), *Cricetodon sansaniensis* has been considered the descendant of *C. aureus* because it possess more developed ectolophs and has metalophulids I more frequently. Agustí (1982) proposed *C. sansaniensis* from ARM7 as the ancestor of *Hispanomys lavocati* from the Vallès-Penedès Basin (named by this author as *Cricetodon* (*Pararuscinomys*) *lavocati*). He supported his hypothesis in the large size of both species, but since then, many other large-sized species of Cricetodontini (such as *Cricetodon soriae* and *Cricetodon engesseri*) have been described and they have to be taken into consideration in any further phylogenetic analysis.




FIGURE 6.1. Length/Width scatter-diagram of the cheek teeth of several large-sized species of *Cricetodon*. Dotted circles denote the range of *C. sansaniensis* from Calatayud-Daroca and dashed lines indicate the maximum and minimum values of the compared species in its type localities. Source of data: Mein and Freudenthal, 1971a (*C. albanensis*), 1971b (*C. aureus*); Sen and Ünay, 1979 (*C. cariensis*); Rummel and Kälin, 2003 (*C. engesseri*); Sen and Maridet, 2012 (*C. sansaniensis*).




FIGURE 6.2. *Cricetodon sansaniensis* from Calatayud-Daroca. Upper molars. Armantes 7: A, inverted right M1, AR7 (58)-542; A, left M3, AR7 (58)-531. Valalto 2C: C, inverted right M3, RGM-266419. Las Umbrías 21: D, inverted right M1, LUM21-179; E, inverted right M2, LUM21-156; F, inverted right M3, LUM21-69. Las Umbrías 22: G, inverted right M2, LUM22-115; H, inverted right M3, LUM22-116. Lower molars: Las Umbrías 21: I, left m1, LUM21-161; J, inverted right m2, LUM21-164; K, left m3, LUM21-166. Las Umbrías 22: L, inverted right m1, LUM22-120; M, inverted right m2, LUM22-122; N, left m3, LUM22-124.

CRICETODON JOTAE Mein and Freudenthal, 1971

(Figs 6.2–6.3; Tables 6.1–6.3)

Material

Original Type Locality—Manchones, Calatayud-Daroca Basin, Spain, local biozone G2, (MN6 upper Aragonian, middle Miocene)

Stratigraphic Range in the Calatayud-Daroca Basin—local biozone G1-local biozone G3 (MN6-MN7/8, upper Aragonian, middle Miocene).

Geographical Range—Southwestern Europe.

Diagnosis

Original Diagnosis (Mein and Freudenthal, 1971a:18, translated from French) —“[Bigger than *C. meini*; the posterior metalophulid is less frequent than in *C. meini*, the anterior metalophulid, however, is more frequent; the ectolophs are better developed, but none of them form a complete connection]”.

Emended Diagnosis—The upper molars show poor development of the mesolophs and posterolophs. The anterior ectoloph of the M1 can be composed by the anterior ectoloph of the paracone or by the posterior ectoloph of the anterocone. The M1 always has a posterior ectoloph, composed by the posterior ectoloph of the paracone, by the anterior ectoloph of the metacone, or by both. The ectolophs are not complete. The lingual posteroloph is common, but short. The main valleys are frequently blocked by cingula and styles. The lower molars have short or absent mesolophids, ectomesolophids and lingual anterolophids. Most of the m1 display a metalophulid I and a metalophulid II together in the same specimen, but some m1 have only a metalophulid I or a metalophulid II. The m2 possess two or three roots; the posterior root can be either partially or completely split.

Differential Diagnosis—Differs from *C. caucasicus* Argyropulo, 1938; *C. versteegi* De Bruijn et al., 1993; *C. tobieni* De Bruijn et al., 1993; *C. aliveriensis* Klein Hofmeijer and De Bruijn, 1988 and *C. kasapligili* De Bruijn et al., 1993 in the larger size of all elements. Apart from that, *C. caucasicus* has a non-divided anterocone; *C. versteegi* backwards directed sinusids; *C. tobieni* and *C. aliveriensis* more developed anterolophs on the M2 and *C. kasapligili* does not have metalophulid I in the m1.

Differs from *C. engesseri* Rummel and Kälin, 2003; *C. soriae* López Martínez et al. in Hernández Fernández et al., 2006; *C. hungaricus* (Kordos, 1986); *C. jumaensis* Rummel, 2001; '*Cricetodon*' *klariankae* Hír, 2007; *C. albanensis* Mein and Freudenthal, 1971a; *C. sansaniensis* Lartet, 1851 and *C. aureus* Mein and Freudenthal, 1971b in the smaller size of all elements. Additionally, *C. engesseri* possess a developed posterolophid on the M3; *Cricetodon albanensis* does not have metalophulid II; *C. jumaensis* and *C. sansaniensis* display a more basal pattern on the m3, with larger mesolophids. '*Cricetodon*' *klariankae* was originally ascribed to *Cricetodon* but recently its generic assignation was questioned (Prieto et al., 2014); it possesses a posterior ectoloph of the M1 mostly double and it is complete. *Cricetodon aureus* has metalophulids II without metalophulid I more frequently than *C. jotae*.

Differs from *C. pasalarensis* (Tobien, 1978) in: anterocone divided by a groove that cover the entire height of the crown; presence of anterior ectoloph; absence of roughness on the basis of the hypocone and the absence of lingual anterolophid in the m3. Differs from *C. candirensis* (Tobien, 1978); *C. cariensis* (Sen and Ünay, 1979) and '*Cricetodon*' *fandli* Prieto et al., 2010 in: ectolophs not complete; absence of enamel coated valley; presence of non-interrupted metalophulid I. Differs from *C. bolligeri* Rummel, 1995 in: absence of ectomesolophid and thick stylids on the m1; anterolophule always connected to the anteroconid on the m1 and the poor development of the mesolophid on the m3. Differs from *C. meini* Freudenthal, 1963 in: absence of neo-entoloph and less-developed mesoloph on the M3. Differs from *C. wanbei* Qiu, 2010 in: absence of neo-entoloph on the M3; forwards directed hypolophulid on the m1. Differs from *C. volkeri* Wu et al., 2009 in: absence of labial spur of the anterolophule; anterocone well divided and short posteroloph on the M1; absence of neo-entoloph on the M3.

CRICETODON JOTAE FROM THE CALATAYUD-DAROCA BASIN

Description of the Type Locality MAN

M1—This molar has four roots. The anterocone shows a furrow which reaches the base of the crown (Fig. 6.4.A; 6.4.D). A posterior protolophule connects the paracone with the posterior branch of the protocone. The anterior ectoloph is frequent (21/27); it is formed by the anterior ectoloph of the paracone in one tooth or by the posterior ectoloph of the anterocone (20/27; Fig. 6.4.A; 6.4.D). One specimen has a well-developed anterior ectoloph reaching the paracone. All the fossils have posterior ectoloph; the posterior spur of the paracone is always present (Fig. 6.4.A; 6.4.D) and in three out of 29 cases it is associated with the anterior spur of the metacone. None of the fossils have the posterior ectoloph complete. The mesoloph is short (9/23), highly reduced

(6/23; Fig. 6.4.D) or absent (8/23; Fig. 6.4.A). A ridge-like structure is developed in the anterior wall of the protocone (14/24); it is absent in the remaining specimens (Fig. 6.4.A; 6.4.D). The labial posteroloph is quite thin but well delimited; it can be absent (4/27). Most molars possess a lingual posteroloph (18/27; Fig. 6.4.D). The anterosinus is closed by a cingulum (20/24), being low in one tooth (Fig. 6.4.A; 6.4.D), and remaining open in three specimens. The mesosinus is blocked by a cingulum (17/23; Fig. 6.4.A; 6.4.D), by a low cingulum (2/23) or it is open (4/23). Several specimens display an open sinus (18/24; Fig. 6.4.A; 6.4.D); it is closed by a cingulum (5/24) or by two crests (1/24). The protosinus can be blocked by a style (5/28) or by a cingulum (22/28) and it remains open in one specimen.

M2—The anterior ectoloph is reduced and generally lower than the anteroloph; it is absent in 12 out of 31 teeth (Fig. 6.4.B; 6.4.E). The posterior ectoloph is always present; it is simple and not connected to the metacone (Fig. 6.4.B; 6.4.E). The mesoloph is frequently absent (17/28; Fig. 6.4.B; 6.4.E), but it can be incipient (11/28). The protocone bears a spine in most molars (18/20). The ectomesoloph is found in only one out of eleven cases. The labial posteroloph is common (26/27; Fig. 6.4.B; 6.4.E), as well as the lingual posteroloph (18/23; Fig. 6.4.E). The well-developed mesosinus is closed by a cingulum in 21 out of 25 specimens (Fig. 6.4.B; 6.4.E); it is open in four. The sinus is open in 23 out of 33 specimens (Fig. 6.4.B; 6.4.E). The protosinus is always weak but distinguishable (Fig. 6.4.B; 6.4.E).

M3—It has reduced hypocone. The lingual anteroloph is distinguishable but weak in 13 out of 18 cases (Fig. 6.4.C). A well-developed labial anteroloph is displayed in 15/18 molars. The anterior ectoloph is absent. The posterior ectoloph is always present; it is formed by both posterior ectoloph of the paracone and anterior ectoloph of the metacone (9/16; Fig. 6.4.C) or only by the posterior ectoloph of the paracone (Fig. 6.4.F). In six specimens the posterior ectoloph is complete (Fig. 6.4.C). Six out of nine teeth have mesoloph; it is short in five teeth and long, reaching the labial border, in one. The axioloph is continuous. Most M3 possess a posteroloph (14/18); it is usually weak and short (Fig. 6.4.F), and ends free. The sinus is long, curved and mostly directed forwards, although it is transversal in one specimen.

m1—The labial anterolophid is a well-developed ridge displayed by most m1 (30/33); it can be connected to the protoconid (10/33), or not (20/33; Fig. 6.4.J). Metalophulid I is found in eight out of 37 teeth, and metalophulid II in five. Twenty-three out of 37 specimens show both metalophulids, I and II (Fig. 6.4.J). Only one incipient mesolophid is observed. The entomesolophid is barely present (2/23); one of them is incipient and the other one is a long crest. Some protoconids bear a spur. The entoconid spur and the labial posterolophid are absent. The hypolophulid is directed forwards (Fig. 6.4.J). The mesosinusid is closed by a cingular ridge (11/35), by a stylid (1/35), or it remains open (23/35; Fig. 6.4.J). The sinusid is wide and short and it is slightly directed forwards (Fig. 6.4.J); it can be blocked by a low cingular ridge (26/34), by two crests attached to

the hypoconid and the protoconid (3/34), or by a stylid (2/34). The sinusid remains open in three teeth. The posterosinusid is blocked by a cingular ridge (4/27) or it is open (Fig. 6.4.J).

m2—This molar shows two (3/9) or three roots (4/9), the posterior one being partially split in two out of four specimens, and completely split in the other two. The lingual anterolophid is found in only three teeth (3/31); it is short and does not reach the metaconid in two of them. The mesolophid is present in 20 out of 33 fossils and it is mostly short (13/20; Fig. 6.4.K); it can be curved and projected towards the metaconid, reaching it in two out of twenty specimens. A short and isolated crest is placed near the mesolophid in five out of twenty molars. There is no anterosinusid and the mesosinusid is blocked by a cingular ridge (9/32) or it remains open (Fig. 6.4.K). In most of the teeth, the sinusid is closed by a cingulid (19/31); it can be blocked by two crests attached to the hypoconid and the protoconid (2/31), by a stylid (1/31), or it can remain open (9/31; Fig. 6.4.K). The lingual posterosinusid can be blocked by a cingular ridge (5/25) or the lingual posterolophid can be directly connected to the entoconid (7/25); it is open in 13 out of 25 specimens (Fig. 6.4.K).

m3—The labial branch of the anterolophid is long and it reaches the protoconid (25/27). In two teeth the anterolophid is extended through the basis of the protoconid towards the sinusid. The lingual branch of the anterolophid is mostly absent (Fig. 6.4.L); two out of ten specimens have it, but without reaching the metaconid. The mesolophid is barely present (8/28; Fig. 6.4.L); it is short (7/28) or long (1/28), but it does not exceed the half length of the mesosinusid (medium). The lingual posterolophid is large and extends towards the entoconid. The anteriorly directed hypolophid is connected to the anterior branch of the hypoconid (Fig. 6.4.L). The sinusid is open (17/27) or blocked by a cingulum (10/27; Fig. 6.4.L). The mesosinusid is mostly open (27/29), but it is closed by a cingulum in two teeth. The lingual posterosinusid is open (20/28; Fig. 6.4.L) or blocked by the lingual posterolophid that projects towards the entoconid closing it.

***Cricetodon jotae* from other sites in the Calatayud-Daroca Basin.**

Cricetodon jotae is a species commonly present in the Calatayud-Daroca Basin but despite the abundance of fossils of this species, detailed morphometric descriptions have been never presented in a systematic study. Here we offer the morphology of the rest of the assemblages of *C. jotae* found in the Calatayud-Daroca Basin in order to know all the intraspecific variability of the species. The fossils of *C. jotae* are homogeneous in morphology and size (Fig 6.2; Table 6.4). However, there is certain degree of variability in several features that are absent in the type locality, which are the following grouped by element:

M1—The anterior ectoloph is composed by the posterior ectoloph of the anterocone and

the anterior ectoloph of the paracone in ARV6 (2/15). Three localities have molars without a posterior ectoloph of the paracone: BOR (1/5), ARV6 (8/17) and TOR1 (1/4). A labial spur of the anterolophule is shown in LP5B (1/2) and TOR1 (1/3). The mesoloph is short in all the localities except for LP5K, where it can be long (1/2). A style is present at the edge of the mesosinus in BOR (2/5), ARV6 (1/17) and TOR1 (1/3), and at the sinus in BOR (1/5), LP5K (1/2), ARV6 (4/6), VILL9, ALC1 and TOR1 (1/3). The protosinus is blocked by two crests in LP5K (1/2) and ARV6 (1/16).

M2— The posterior ectoloph is absent in BOR (1/3) and ALC3, and a style is blocking the mesosinus in LP5C (1/2).

M3—The posterior ectoloph of the paracone is absent in a specimen from ARV6 (1/13). The mesoloph exceeds the half of the length of the mesosinus, but it does not reach the labial border in ARV6 (2/13). The mesoloph and ectoloph can be connected in BOR (1/2), and the neo-entoloph is found in ARV6 (1/13) and in the only M3 from ALC1.

m1—The lingual anterolophid is present and reaches the metaconid in LP5K (1/4), ARV6 (1/15), and ALC3. The mesolophid is short in specimens from BOR (1/4), ALC2 (2/3) and TOR1 (1/2).

m2—The labial anterolophid is short or absent in two out of twenty m2 from ARV6; rarely, the mesolophid exceeds the half of the length of the mesosinusid, but without reaching the labial border, in ARV6 (2/21); the entomesolophid is present but incipient in ARV6 (2/21) and VILL9 (1/4). The sinusid is closed by a stylid and a crest attached to the hypoconid in ARV6 (1/20).

m3—The lingual anterolophid is present in one out of 17 cases from ARV6. The mesolophid is connected to the metaconid in ARV6 (1/20) and is double in BOR (1/20). The sinusid is closed by a cingulid in LP5B (1/2), BOR (1/2), LP5L, ARV6 (1/19) and in ALC2 (1/2); closed by two crests in ARV6 (1/19) and in VILL9 (1/2); and a stylid is shown in ARV6 (1/19) and in TOR1 (1/3).

The scatter-diagrams (Fig. 6.2) show a large overlap in the distributions of the measurements among the studied sites, suggesting a similar size of the fossils of *Cricetodon jotae* in the Calatayud-Daroca Basin.

Discussion

Cricetodon from Manchones was firstly described by Freudenthal (1963) as *Cricetodon meini*. A few years later, in 1966, he improved this study providing new measurements and descriptions

of the fossils from that locality. The last study of the type assemblage is presented in the revision of the rodent faunas of the Cenozoic made by Mein and Freudenthal (1971a). They described a new species, *C. jotae* Mein and Freudenthal, 1971a in the material from Manchones and Arroyo de Val 6. The authors refer to Freudenthal's studies (1963 and 1966) for the metrical information and the comparison with other species is shortly described.

Outside the Calatayud-Daroca Basin, *Cricetodon jotae* is also registered in two Portuguese localities, Povóia do Santarem and Chloes (MN6; Antunes and Mein, 1977), as well as in the French locality of La Grive L7 (MN7; Mein and Freudenthal, 1971a). It is nowhere else recorded, thus the geographical range of *C. jotae* is restricted to the southwestern Europe.

Morphologically, *Cricetodon jotae* displays characters, such as the metalophulid II, which are considered basal for the tribe Cricetodontini (Mein and Freudenthal., 1971a), and also has features such as: absence of mesoloph; presence of double ectoloph; low-developed or absent mesolophids; absence of ectomesolophid and weak or absent labial anterolophid of the m1, defined as derived by many authors (Agustí, 1982; López-Antoñanzas and Mein, 2009). This mosaic distribution of the characters, combining derived and basal traits, was described in other species of Cricetodontini from the late Aragonian like *Hispanomys aguirrei* (Sesé Benito, 1977) and *Cricetodon nievei* López-Guerrero et al., 2014.

Only few authors have discussed the phylogenetic relationships of *C. jotae*. Mein and Freudenthal (1971a) suggested that *C. jotae* could be related with *C. meini*. The evolutionary trends that they described are: an increase of the hypsodonty and the development of the ectolophs. On the other hand, according to Agustí (1982), *Hispanomys aguirrei* could be the descendant of *C. jotae*, assuming a development which resulted in more-developed ectolophs, increased hypsodonty and the reduction of the cingular formations (Agustí, 1982).

Due to the above referred mosaic-type morphology of *C. jotae* and *H. aguirrei*, it is difficult to establish their phylogenetic relationships. A cladistic analysis could be used to evaluate them, but it will be considered in chapter 8.




FIGURE 6.2. Length/Width scatter-diagram of the cheek teeth of *Cricetodon jotae* from Calatayud-Daroca (including all specimens), *C. meini* from Vieux-collonges and *C. bolligeri* from Petersburg 10. Dashed lines indicate the maximum and minimum values of the compared species in its type localities. Source of data: Mein and Freudenthal, 1971b (*C. meini*), Rummel, 2001 (*C. bolligeri*).




FIGURE. 6.4. *Cricetodon jotae* from Calatayud-Daroca. Upper molars. Manchones: A, inverted right M1, MA-2183; B, inverted right M2, MA-2199; C, left M3, MA-2213; D, inverted right M1, MA-2188; E, left M2, MA-2196; F, inverted right M3, MA-2220. Las Planas 5C: G, inverted right M1 LP5C-5; H, left M2, LP5C-8; I, inverted right M3, LP5C-11. Lower molars. Manchones: J, left m1, MA-2225; K, left m2, LP5C-2245; L, left m3, MA-2253. Las Planas 5C: M, inverted right m1, LP5C-19; N, left m2, LP5C-15; O, inverted right m2, LP5C-16.

FINAL REMARKS & CONCLUSIONS

After this comprehensive study, all the material of *Cricetodon* from the Calatayud-Daroca Basin has been definitively assigned at species level. *Cricetodon sansaniensis* was previously found in ARM7 and now is also recovered from other four localities. Similarly, eleven new occurrences of *C. jotae* have been described. The metrical and morphological analysis allow us to clarify the doubtful assignation of the scarce material from ARM7 to *C. sansaniensis*, and to emend the diagnosis of *C. jotae*. Moreover, the increase of the sample size lets us to improve the comparison with the rest of the species of *Cricetodon*.

Cricetodon sansaniensis from Calatayud-Daroca Basin has weak ectolophs and both metalophulids, I and II, on the same m1, and mesolophs/ids that, according with the trends described for Cricetodontini, are basal traits. *Cricetodon sansaniensis* has been related with *C. aureus* and *C. albanensis*. However, we consider that this relationship has to be re-evaluated, since there are some species recently described (*C. soriae* and *C. engesseri*) that could be descendant of *C. sansaniensis* as well.

Regarding *C. jotae*, we observe a combination of basal character (metalophulid II) and derivate features (absence of mesolophs; presence of double ectolophs; low-developed or absent mesolophids; absence of ectomesolophid and weak or absent labial anterolophid of the m1). This combination of primitive and derivate character could be the result of mosaic evolution. This particularity in the morphology hinder the assessment of its phylogenetic relationships. Chapter 10 treats this questions by means of a cladistic analysis and an evaluation of the phylogeny of *C. jotae* will be presented there.

Locality		Length				Width			
		Min	Mean	Max	N	Min	Mean	Max	N
M1	ARM7	3.05	3.19	3.26	3	2.04	2.16	2.28	1
	LUM21		3.39		1	2.03	2.14	2.25	2
M2	VT2C	2.44	2.49	2.54	4	2.01	2.06	2.14	4
	ARM7	2.35	2.37	2.39	2	1.93	1.94	1.96	2
	LUM21	2.42	2.51	2.59	3	1.93	2.02	2.09	3
	LUM22		2.38		2	1.88	1.97	2.06	2
	VT2C	1.86	1.88	1.89	2	1.83	1.84	1.84	2
M3	ARM7	1.95	2.06	2.17	2	1.88	1.90	1.93	2
	LUM21	1.86	1.92	1.98	2	1.76	1.87	1.98	2
	LUM22		2.01		1		1.86		1
	VT2C						1.78		1
m1	VT2B					1.69	1.73	1.78	2
	ARM7		2.55		1		1.76		1
	LUM21		2.67		1		1.79		1
	LUM22		2.60		1		1.75		1
	VT2C						1.78		1
m2	ARM7		2.73		1		1.96		1
	LUM21		2.72		1	1.99	2.08	2.16	2
	LUM22		2.79		1		2.17		1
	VT2C						2.07		1
m3	VT2B					1.94	2.01	2.07	2
	ARM7	2.34	2.54	2.74	2	1.77	1.94	2.10	2
	LUM21	2.52	2.53	2.54	2	1.77	1.87	1.96	2
	LUM22	2.60	2.62	2.63	2	1.92	1.96	1.99	2
	VT2C						2.07		1

TABLE 6.3. Length and width of the studied material of *Cricetodon sansaniensis*. Units are given in mm. Abbreviations: Min, minimum values; Max, maximum values; N, number of specimens.

	NM Zone	Local	Locality	Length				Width			
		biozone		Min	Mean	Max	N	Min	Mean	Max	N
M1	MN7/8	G3	TOR1	2.62	2.80	2.89	3	1.82	1.87	1.92	3
	MN7/8	G3	AC1		3.12		1		2.07		1
	MN7/8	G3	VL9		2.95		1				
	MN6	G2	ARV6	2.76	2.99	3.29	13	1.83	1.98	2.12	16
	MN6	G2	LP5K		2.98		1	1.94	2.01	2.07	2
	MN6	G2	BOR	2.82	3	3.14	4	1.98	1.99	2	4
	MN6	G2	MAN	2.69	2.94	3.18	19	1.80	1.99	2.18	19
	MN6	G1	VT1		3.14		1		2.03		1
	MN6	G1	LP5C		2.93		1	1.82	1.87	1.97	3
	MN6	G1	LP5B		3.01		1	1.83	1.93	2.03	2
M2	MN7/8	G3	TOR1	2.19	2.23	2.27	2	1.95	1.96	1.97	2
	MN7/8	G3	VL9		2.28		1		1.91		1
	MN6	G2	ARV6	2.02	2.26	2.43	8	1.71	1.92	2.11	9
	MN6	G2	LP5K	2.17	2.24	2.36	3	1.90	1.93	1.99	3
	MN6	G2	LP5L		2.28		1		1.81		1
	MN6	G2	BOR	2.17	2.21	2.26	3	1.86	1.88	1.90	3
	MN6	G2	MAN	2.05	2.23	2.41	8	1.69	1.87	2.01	8
	MN6	G1	LP5C	2.23	2.26	2.29	2	1.81	1.88	1.94	2
M3	MN7/8	G3	TOR1		2.55		1		1.73		1
	MN7/8	G3	AC1		1.74		1		1.76		1
	MN7/8	G3	VL9	1.65	1.69	1.73	2		1.72		2
	MN6	G2	ARV6	1.58	1.75	1.89	15	1.50	1.67	1.85	15
	MN6	G2	LP5K		1.85		1		1.78		1
	MN6	G2	BOR	1.89	1.90	1.92	2	1.73	1.79	1.85	2
	MN6	G2	MAN	1.65	1.83	2	18	1.58	1.71	1.83	18
	MN6	G1	LP5C		1.81		1		1.71		1
m1	MN7/8	G3	TOR1	1.67	1.76	1.86	3	1.68	1.73	1.80	3
	MN7/8	G3	AC3		2.46		1		1.62		1
	MN7/8	G3	AC2						1.66		1
	MN7/8	G3	VL9		2.36		1		1.55		1
	MN6	G2	ARV6	2.39	2.50	2.69	14	1.57	1.65	1.76	15
	MN6	G2	LP5K	2.31	2.49	2.59	3	1.60	1.69	1.75	4
	MN6	G2	LP5L		2.41		1		1.67		1
	MN6	G2	BOR	2.45	2.48	2.51	2	1.57	1.64	1.71	3
	MN6	G2	MAN	2.28	2.48	2.68	35	1.50	1.65	1.82	35
	MN6	G1	VT1		2.47		1		1.62		1
	MN6	G1	LP5C		2.68		1		1.70		1
	MN6	G1	LP5B								
m2	MN7/8	G3	TOR1		2.46		1		1.83		1
	MN7/8	G3	AC2		2.35		1	1.81	1.88	1.95	2
	MN7/8	G3	AC1						2.02		1
	MN7/8	G3	VL9	2.29	2.44	2.60	4	1.79	1.92	2.03	4
	MN6	G2	ARV6	2.35	2.46	2.59	17	1.78	1.92	2.05	18
	MN6	G2	LP5K	2.38	2.41	2.44	2	1.91	1.92	1.93	2
	MN6	G2	BOR	2.36	2.41	2.46	2	1.85	1.86	1.86	2
	MN6	G2	MAN	2.22	2.39	2.65	27	1.71	1.87	2.07	28
	MN6	G1	LP5C	2.36	2.49	2.61	2	1.86	1.87	1.87	2
	MN6	G1	LP5B								
	MN6	G1	VT1								
	MN6	G1	AC1								
m3	MN7/8	G3	TOR1	2.24	2.32	2.39	2	1.76	1.83	1.89	2
	MN7/8	G3	AC2		2.17		1		1.76		1
	MN7/8	G3	VL9	1.96	2.08	2.19	2	1.75	1.76	1.77	2
	MN6	G2	ARV6	2.13	2.27	2.39	15	1.69	1.82	2	16
	MN6	G2	LP5L		2.49		1		1.92		1
	MN6	G2	BOR	2.28	2.36	2.44	2	1.77	1.80	1.83	2
	MN6	G2	MAN	2.12	2.32	2.60	10	1.66	1.82	2	10

TABLE 6.4. Length and width of the studied material of *Cricetodon jotae*. Units are given in mm. Abbreviations: Min, minimum values; Max, maximum values; N, number of specimens.

LITERATURE CITED

- Agustí, J. 1982. Tendencias evolutivas de la línea *Cricetodon-Ruscinomys* (Rodentia, Mammalia) en la Península Ibérica. *Acta Geologica Hispanica* 17:103–111.
- Antunes, M., and P. Mein. 1977. Contributions à la paléontologie du miocène moyen continental du bassin du Tage. III. Mammifères-póvoa de Santarem, Pero Filho et Choes (Secorio) conclusiones generales. *Ciências da Terra (UNL)* 2:143–165.
- Argyropulo, A. I. 1938. On the fauna of Tertiary Cricetidae of the USSR. *Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS* 20:223–226.
- Baudelot, S. 1972. Etude des Chiroptères, Insectivores et Rongeurs du Miocène de Sansan (Gers). 364 pp. Ph.D. dissertation, Université Paul Sabatier, Toulouse, France.
- Bolliger, T. V. 1994. Die Obere Süßwassermolasse in Bayern und der Ostschweiz: bio- und lithostratigraphische Korrelationen. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und histor. Geologie* 34:109–144.
- Bowdich, T. E. 1821. *An Analysis of the Natural Classification of Mammals for the Use of Students and Travelers*. J. Smith, Paris, 115 pp.
- Daams, R., and M. Freudenthal. 1981. Aragonian: the stage concept versus Neogene mammal zones. *Scripta Geologica* 62:1–17.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, and W. Krijgsman. 1999. Aragonian stratigraphy reconsidered and a re-evaluation of the middle Miocene mammal biochronology in Europe. *Earth and Planetary Science Letters* 165:287–294.
- De Bruijn, H., V. Fahlbusch, G. Sarac, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a discussion of the evolutionary history of the Cricetodontini. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 96:151–216.
- Fischer, G. 1817. *Adversaria zoologica*, *Mémoires de la Société Impériale des Naturalistes de Moscou* 1917:357–472.
- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mittelspaniens und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.

- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B 69:298–317.
- García Moreno, E. 1988. The Miocene rodent biostratigraphy of the Duero basin (Spain): a proposition for a new Aragonian/Vallesian limit. *Paleontologia i Evolució* 22:103–112.
- Ginsburg, L. and C. Bulot. 2000. Le cadre stratigraphique du site du Sansan; pp. 39–67 in Ginsburg, L. (eds.), *La faune miocène du Sansan et son environnement*. Memoires du Museum National d'Histoire Naturelle. Paris.
- Hernández Fernández, M., J. A. Cárdena, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőtárkány-Felnémet (Northern Hungary). *Fragmenta Paleontologica Hungarica* 24–25:16–24.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254:177–235.
- Kordos, L. 1986. Upper miocene hamsters (Cricetidae, Mammalia) of Hasznos and Szentendre. *Magyar Allami Földtani Intézet Jelentése Az* 1984:523–553.
- Klein Hofmeijer, G., and H. De Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece) Part 8: The Cricetidae. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B 91:185–204.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d'une récapitulation des diverses espèces d'animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d'autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. J. A. Portes, Auch, 41 pp.
- López-Antoñanzas, R., and P. Mein. 2009. First detailed description of *Hispanomys bijugatus* Mein & Freudenthal, 1971 (Rodentia, Cricetodontinae) from the Upper Aragonian of La Grive-Saint Alban (France): Biostratigraphical implications. *Geobios* 42:783–796.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89: 573–592.

- López-Martínez, N. 1997. Lignées évolutives, formes intermédiaires et évolution en mosaïque: exemple des lagomorphes européens; pp. 105–120 in J. P., Aguilar, S. Legendre, and J. Michaux (eds.), Congrès BiochroM '97. Mémoires et Travaux EPHE, Institut Montpellier, Montpellier.
- Maridet, O., and S. Sen. 2012. Les Cricetidae du gisement de Sansan; pp 29–64 in S. Peigné and S. Sen (eds.), Les mammifères du gisement de Sansan. Mémoires du Muséum national d'Histoire naturelle, Paris.
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.
- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata Palasiatica* 48:27–47.
- Rummel, M. 1995. *Cricetodon bolligeri* n. sp. ein neuer Cricetide aus dem Obermiozän von Petersbuch bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 35:109–123.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde B* 311:1–6.
- Rummel, M., and D. Kálin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Mittelmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Sen, S., and E. Ünay. 1979. Sur quelques Cricetodontini (Rod.) du Miocène moyen d'Anatolie. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* 82:293–301.
- Sesé Benito, C. 1977. Los cricétidos (Rodentia, Mammalia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno Cuaternario* 8:127–80.
- Simpson, G. C. 1945. The principles of classification and a classification of mammals. *Bulletin of the American Museum of Natural History* 85:1–350.
- Stehlin, H. G., and S. Schaub. 1951. Die Trigonodontie der simplicidentaten Nager. *Schweizerische paläontologische Abhandlungen* 67:1–385.
- Tobien, H. 1978. New species of Cricetodontini (Rod. Mam.) from the Miocene of Turkey.

Mainzer Geowissenschaftliche Mitteilungen Geologisches Landesamt Rheinland-Pfalz 6:209–219.

Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.

Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.

Chapter 7

Cricetodon nievei



NEW CRICETODONTINI FROM THE MIDDLE MIOCENE OF EUROPE: AN EXAMPLE OF MOSAIC EVOLUTION

“Nieves had a strong charming personality. She was enthusiastic, curious, active, fascinating. She was the synonym of bravery, rigorousness, excellence, talent and hard work.”

Marián Álvarez-Sierra, Humberto Astibia Ayerra, Manuel Hernández Fernández & Salvador Moyà Solà

Resumen

7

En este capítulo se describe una nueva especie de Cricetodontini (Cricetidae, Rodentia, Mammalia), *Cricetodon nievei* sp. nov., en la sección de Toril (Toril 3A, Toril 3B, Toril2) y en la localidad de Las Planas 5H. Estos yacimientos pertenecen a la biozona local G3 (Aragoniense superior, final del Mioceno medio) de la Cuenca de Calatayud-Daroca. La nueva especie muestra un patrón en mosaico consistente en una combinación de caracteres primitivos y derivados: los molares superiores tienen un patrón basal tipo *Cricetodon*—con caracteres tales como ectolofos cortos y no completos—, mientras que los molares inferiores comparten con los representantes más antiguos de *Hispanomys* varias características derivadas—e.g., ausencia de metalofúlido II—. También se discute la asignación genérica del nuevo taxón y se compara éste con todas las especies de *Cricetodon* y *Hispanomys* descritas hasta la fecha. El patrón de evolución en mosaico de los caracteres dentales se reconoce también en otras especies de Cricetodontini de Europa durante el Aragoniense superior, si bien con diferente combinación de caracteres morfológicos. En ese intervalo de tiempo, la diversidad de la tribu aumenta, incluyendo especies con morfología más compleja y mayor variabilidad intraespecífica que los representantes más antiguos. La nueva especie propuesta es morfológicamente cercana a los Cricetodontini registrados fuera de la Cuenca de Calatayud-Daroca, especialmente *C. albanensis* e *H. decedens* de Francia. Por último, se analiza el contexto paleoecológico de la nueva especie; la distribución estratigráfica de *Cricetodon nievei* sp. nov., coincide con cambios en las comunidades faunísticas que están posiblemente relacionados con un aumento de la humedad detectado en la cuenca.

Referencia

- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes and Peláez-Campomanes, P. 2013. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 82:573–792.



Abstract

A new species of Cricetodontini (Cricetidae, Rodentia, Mammalia), *Cricetodon nievei* sp. nov. from the Toril section (Toril 3A, Toril 3B, Toril 2) and Las Planas 5H is described. All this sites belong to the local biozone G3 (late Aragonian, late middle Miocene) from the Calatayud-Daroca Basin (Zaragoza, Spain). The new species displays a mosaic pattern consisting in a combination of primitive and derived characters: the upper molars have a basal *Cricetodon*-like pattern—short and not complete ectolophs—, whereas the lower molars share several derived features with the older representatives of *Hispanomys*—absence of metalophulid II. The generic assignment of the new taxon is further discussed and it is compared with all the species of *Cricetodon* and *Hispanomys* described until date. This mosaic evolutionary pattern of dental characters is also recognized in other species of Cricetodontini from Europe during the late Aragonian, although involving different combination of morphological characters. At this time, the diversity of the tribe increases, including species with complex morphology and higher intraspecific variability than the older representatives. The new species proposed here is morphologically close to Cricetodontini recorded outside the Calatayud-Daroca Basin, especially *C. albanensis* and *H. decedens* from France. Finally, the palaeoecological context of the new species is discussed; the stratigraphical distribution of *Cricetodon nievei* sp. nov. coevals changes in the faunal assemblages which are possibly related to an increase of humidity detected in the Calatayud-Daroca Basin.

INTRODUCTION

The tribe Cricetodontini (Rodentia) is commonly present in the European Miocene; it provides relevant information regarding the biochronology, biostratigraphy and palaeoecology of the continental deposits (Van der Meulen and Daams, 1992; Daams et al., 1999c, 1999a; Van der Meulen et al., 2011, 2012). The representatives of Cricetodontini are considered to be immigrants in Asia Minor around the Oligocene-Miocene boundary (De Bruijn and Ünay, 1996). During the early Miocene (Mammal Neogene zones MN1 and MN2, sensu De Bruijn et al., 1992 and Steininger 1999), the tribe remains modest in the number of both species and specimens. The diversification and geographical expansion into continental Europe start later on, at MN5. During the late Aragonian (MN7/8), the tribe reaches its maximum diversity (De Bruijn and Ünay, 1996; Rummel 1999). In the middle Aragonian the species of Cricetodontini display a widely extended geographical distribution; all of them are included in the genus *Cricetodon*, showing a primitive stage-of-evolution according to De Bruijn and Ünay (1996).

During the latest part of the middle Miocene (late Aragonian) new combinations of morphological dental characters among the species occur. The intraspecific variability and the diversity seem to be higher than in older assemblages (De Bruijn and Ünay, 1996). The taxa exhibit a mosaic morphology that consists in the development of different rates of change in the jugal teeth. In this interval, the transition from *Cricetodon* to *Hispanomys*, in southwestern Europe (Agustí, 1980; De Bruijn et al., 1993; Aguilar et al., 1994; López-Guerrero et al., 2008; López-Antoñanzas & Mein, 2009), and to *Byzantinia* in Turkey (De Bruijn et al., 1993; De Bruijn and Ünay, 1996) took place. Therefore, the evolutionary history of the tribe Cricetodontini gathers important taxonomical changes during the late Aragonian.

In this paper we have studied the fossils of a new species found in the upper Aragonian from the Calatayud-Daroca Basin (Daams et al., 1999b; Álvarez-Sierra et al., 2003). The rodent fauna from this basin has been extensively studied by many authors (Daams and Freudenthal, 1981, 1988; Álvarez Sierra, 1987; Daams et al., 1998; Peláez-Campomanes, 2001; Van der Meulen et al., 2003; García-Paredes, 2006; Oliver et al., 2009; García-Paredes et al., 2009, 2010; López-Guerrero et al., 2013; Oliver and Peláez-Campomanes, 2013). Specifically, our study is part of an ongoing systematic and phylogenetic revision of the Cricetodontini from this basin (López-Guerrero et al., 2008, 2009, 2011, 2013).

The tribe Cricetodontini is represented by six species in the late Aragonian from Calatayud-Daroca Basin (Freudenthal, 1966; Mein and Freudenthal, 1971a; López-Guerrero et al., 2008, 2009): *Cricetodon* cf. *sansaniensis* Lartet, 1851, *Cricetodon jotae* Mein and Freudenthal, 1971, *Hispanomys aguirrei* (Sesé Benito, 1977), *Hispanomys lavocati* (Freudenthal, 1966), *Hispanomys nombrevillae* (Freudenthal, 1966), and the new species described here. Their stratigraphical distributions, according to the local biozonation proposed by Daams et al. (1999c), are as follows: *Cricetodon* cf. *sansaniensis* is present in the local biozone F (Freudenthal, 1966); *Cricetodon jotae* in the G1 (pers. obs.), G2 (Mein and Freudenthal, 1971a) and the early G3 (pers. obs.); *Hispanomys aguirrei*, *Hispanomys lavocati* and the new species, in the G3 (Lopez-Guerrero et al., 2008), and finally, *Hispanomys nombrevillae* is found in the H (López-Guerrero et al., 2009).

Here we describe the new taxon that shows a particular pattern with a combination of basal and derived features. We also compare it in detail with the species of Cricetodontini with close morphology and discuss the context and status with regards to other representatives of the tribe in Europe.

MATERIAL & METHODS

The studied material includes 274 upper and lower molars. Table 7.1 contains the number of specimens studied for each fossil site and the abbreviations used. They are housed at the Museo Nacional de Ciencias Naturales-CSIC (Madrid, Spain) and at the Naturalis Biodiversity Center (Leiden, The Netherlands). The studied fossils belong to four localities from the Toril-Nombrevilla section—TOR2, TOR3A, TOR3B—and Las Planas section—LP5H. These sections are located in the Calatayud-Daroca Basin, southern part of the province of Zaragoza (Fig. 7.1).

<i>Cricetodon nievei</i> sp. nov.									
Locality	Abb.	Collections	M1	M2	M3	m1	m2	m3	N
Toril 3B	TOR3B	MNCN	8	12	4	10	8	6	48
Toril 3A	TOR3A	MNCN	18	30	27	36	24	24	159
Toril 2	TOR2	MNCN	4	5	2	—	4	2	17
Las Planas 5H	LP5H	RGM	7	9	4	10	8	12	50
		Total	37	56	37	56	44	44	274

TABLE 7.1. Material of *Cricetodon nievei* sp. nov. from the Calatayud-Daroca Basin, studied localities, abbreviations for the localities and institutions where the fossils are stored (see institutional abbreviations). Abb= abbreviations for the localities; N= number of specimens.

All the sites belong to the local biozone G3 from the late Aragonian (Álvarez-Sierra et al., 2003) and they can be correlated to the Mammal Neogene zone MN7/8 (Daams et al., 1999b). The locality LP5H has been correlated to the chron C5Ar.1r (Daams et al., 1999a) that spans a time range of 12.474 Ma–12.735 Ma (Hilgen et al., 2012). According to Van Dam et al. (2006), the numerical age of LP5H is 12.57 Ma using the local magnetostratigraphical correlation. Additionally, Álvarez Sierra et al. (2003) and Garcés et al. (2003) correlated LP5H with the localities Toril 2, Toril 3A and Toril 3B.

The new material has been compared with several taxa of Cricetodontini from different localities hosted in other institutions: *Cricetodon soriae* from the type locality Somosaguas stored at the Universidad Complutense de Madrid (Spain); type material of *C. bollikeri* from Petersburg 10 and '*Cricetodon*' *fandli* from Gratkorn stored at Bayerische Staatssammlung für Paläontologie




FIGURE 7.1. Geological map of the Calatayud-Daroca Basin, the square marks the situation of the area of Daroca (Modified after García-Paredes et al., 2010). The orthophotograph shows the position of Toril and Nombrevilla sections.

und Geologie, Munich (Germany); *C. sansaniensis* from the type locality of Sansan stored at the Muséum d'Histoire Naturelle de Bâle (Switzerland); type material of *C. aureus* and *C. meini* from Vieux-Collonges, and *C. albanensis* from La Grive–Saint Alban stored at the Naturalis Biodiversity Center (NBC), Leiden (The Netherlands); *C. klariankae* from the type locality Felsőtárkány–Felnémet stored at Municipal Museum of Pásztó, Pásztó (Hungary), *C. jotae* from its type locality, Manchones, stored at the Instituut voor Aardwetenschappen, Utrecht University (The Netherlands) and finally, the type material of *Hispanomys aguirrei* from Escobosa de Calatañazor stored at the Museo Nacional de Ciencias Naturales, CSIC, Madrid (Spain).

The nomenclature used for the structures present in the occlusal surface, modified after Mein and Freudenthal (1971b), is summarized in Figure 7.2. Notation M1, M2, M3/m1, m2, m3 is used for upper/lower molars, respectively. Maximum length and maximum width, given in mm, were measured using a Nikon Measuroscope 10 microscope with digital micrometer following López-Guerrero et al. (2013). Descriptive statistics of measurements are given in Table 7.2. Length/width scatter-diagrams of *C. nievei* sp. nov. are shown in Figure 7.3. The scatter-diagrams include *C. jotae* from its type locality, Manchones (Calatayud-Daroca Basin), due to the geographical and temporal proximity, and because of the large sample size which allows straightforward comparisons with the new species. Among the assemblages of *C. nievei* sp. nov. size have been compared by performing Levenne's and ANOVA tests in localities with at least five data per dental element (Table 7.3). Tables 7.4 and 7.5 present the measurements of the species of *Cricetodon* and *Hispanomys* (type localities) from the late Aragonian used in the discussion. The

photographs were taken with an Environmental Scanning Electron Microscope FEI Quanta 200 in environmental mode at the Museo Nacional de Ciencias Naturales-CSIC (Madrid, Spain). Descriptive and analytical statistics have been carried out with IBM SPSS v. 19.0.0 (IBM SPSS 2010). The molars are figured as if they were from the right side; reversed images have been indicated by an underlined label.

Repository Institution—All specimens are housed in the Museo Nacional de Ciencias Naturales-CSIC (Madrid, Spain) with a repository number prefixed by TOR.

Institutional abbreviations—MNCN, Museo Nacional de Ciencias Naturales-CSIC, Madrid, Spain; RGM, Naturalis Biodiversity Center, former Rijksmuseum van Geologie en Mineralogie, Leiden, The Netherlands.




FIGURE 7.2. Terminology of the parts of the cheek teeth of *Cricetodon*. Only the M1, M3 and the m1 morphology has been draw; however, this nomenclature can be applied for the rest of the upper and lower molars. (Modified after Mein and Freudenthal, 1971b).

SYSTEMATIC PALEONTOLOGY

Order RODENTIA Bowdich, 1821

Family CRICETIDAE Fischer, 1817

Subfamily CRICETODONTINAE Stehlin and Schaub, 1951

Tribe CRICETODONTINI Simpson, 1945

Genus *CRICETODON* Lartet, 1851

CRICETODON NIEVEI sp. nov.

(Figs 7.4–7.6; Tables 7.1–7.3)

1981 *Cricetodon* from LP5H—Daams and Freudenthal, p. 4, fig. 1.

1984 *Cricetodon* from LP5H—Daams and Van der Meulen, p. 246, fig. 2.

1988 *Cricetodon* spp. from LP5H—Daams and Freudenthal, p. 14, fig. 8.

2003 *Cricetodon* spp. from TOR2, Toril3A and TOR3B—Álvarez-Sierra, Calvo, Morales, Alonso-Zarza, Azanza, García Paredes, Hernández Fernández, Van der Meulen, Peláez-Campomanes, Quiralte, Salesa, Sánchez and Soria, p. 30, table 1.

2004 *Cricetodon* sp.—Azanza, Alonso Zarza, Álvarez Sierra, Calvo, Fraile, García Paredes, Gómez, Hernández Fernández, Van der Meulen, de Miguel, Montoya, Morales, Murelaga, Peláez-Campomanes, Pérez, Quiralte, Salesa, Sánchez Marco and Soria, p. 273, table 2.

2009 *Cricetodon* aff. *jotae*—López-Guerrero, Álvarez-Sierra, García-Paredes, López-Antoñanzas and Oliver, p. 161A.

Material

Holotype—Left m1, TOR3A-321 (Fig. 7.4H).

Paratypes—M1: TOR3A-118–TOR3A-131, TOR3A-268–TOR3A-270, TOR3A-278; M2: TOR3A-138–TOR3A-166, TOR3A-287; M3: TOR3A-218–TOR3A-237,

TOR3A-260–TOR3A-236; m1: TOR3A-168–TOR3A-197, TOR3A-317–TOR3A-322; m2: TOR3A-198–TOR3A-217, TOR3A-323–TOR3A-326; m3: TOR3A-238–TOR3A-259, TOR3A-267, TOR3A-297.

Type horizon—Local biozone G3, upper Aragonian, middle Miocene.

Type locality—TOR3A, province of Zaragoza, Calatayud-Daroca Basin, Spain.

Other localities—LP5H, TOR2 and TOR3B (see Table 7.1).

Stratigraphic range—Local biozone G3, MN7/8, upper Aragonian, middle Miocene.

Etymology—Dedicated to our mentor Dr. Nieves López Martínez who made a great contribution to the palaeontology of micromammals.

Diagnosis

Diagnosis—Medium-sized *Cricetodontini* with thick enamel. The main valleys of the upper molars are almost closed by large cingular ridges and/or styles; the latter are especially frequent in the protosinus. Upper molars always have a short mesoloph except for the M3 where the mesoloph can be absent. M1 with four roots and the anterocone well divided by a groove. The anterior ectoloph is frequently present, simple and not complete. The posterior ectoloph is always present, simple and not complete. The posteroloph, in the M3, is absent. The lower molars present a short mesolophid. The metalophulid I on the m1 is always present and the metalophulid II is absent. The m1 and m2 frequently have a well-developed ectomesolophid which can reach the labial border. The m2 is generally birradiculated; in some cases, its posterior root is partly divided.

Differential diagnosis—*Cricetodon nievei* sp. nov. differs from *C. caucasicus* Argyropulo, 1938, *C. versteegi* De Bruijn, Fahlbusch, Saraç, and Ünay, 1993, *C. tobieni* De Bruijn, Fahlbusch, Saraç, and Ünay, 1993, *C. aliveriensis* Klein Hofmeijer and De Bruijn, 1988, *C. kasapligili* De Bruijn, Fahlbusch, Saraç, and Ünay, 1993, and *C. wanhei* Qiu, 2010 by its larger size.

Cricetodon nievei sp. nov. differs from *C. engesseri* Rummel and Kälin, 2003, *C. soriae* López Martínez, Cárdbaba, Salesa, Hernández Fernández, Cuevas González and Fesharaki, 2006 in Hernández Fernández et al. 2006, *C. jumaensis* Rummel, 2001, *C. klariankae* Hír, 2007, *C. albanensis* Mein and Freudenthal, 1971a, *C. sansaniensis* Lartet, 1851 and *C. aureus* Mein and Freudenthal, 1971b by its smaller size.

Cricetodon nievei sp. nov. differs from *C. pasarlariensis* (Tobien, 1978) by the deeply split anterocone in the M1; its M2 and M3 with less-developed lingual anterolophs and the absence of

metalophulid II in the m1. It differs from *C. candirensis* (Tobien, 1978) and *C. hungaricus* (Kordos, 1986) by its less-developed ectolophs; the presence of metalophulid I on the m1; the absence of the connections in an 'X-shaped' pattern among the four main cusps in the m3. Differs from *C. cariensis* (Sen and Ünay, 1979) by its well-developed ectolophs—which do not form a continuous structure—and the absence of enamel-coated valley. Differs from *C. bolligeri* Rummel, 1995 by the absence of anterior protolophule on the M1, and its less developed ectomesolophid on the m1. Differs from *C. meini* Freudenthal, 1963 by its presence of posterior ectoloph; the absence of anterior protolophule on the M1 and metalophulid II on the m1. Differs from *C. jotae* Mein and Freudenthal, 1971a by the presence of styles on the protosinus of the M1; the presence of mesolophs on the upper molars; its protocone connected to the longitudinal crest through the posterior arm of the protocone; the absence of metalophulid II. Differs from *C. volkeri* Wu, Meng, Ye, Ni, Bi and Wei, 2009 by its M1 with four roots and less-developed posteroloph; M2 with less-developed mesolophs; larger m2. Differs from '*Cricetodon*' *fandli* Prieto, Böhme and Gross, 2010 by the M1 without double, and complete anterior ectoloph and m1 without metalophulid II.

Cricetodon nievei sp. nov. differs from *Hispanomys dispectus* Agustí, 1980 by its M1 with poor-developed ectolophs, its sinuous contour; the absence of enamel-coated valley and the m1 without metalophulid II. Differs from *H. daamsi* Agustí, Casanovas-Vilar and Furió, 2005 and *H. lavocati* (Freudenthal, 1966) by its smaller size; the presence of mesoloph and poor-developed and not complete ectolophs on the upper molars. Differs from *H. aguirrei* (Sesé Benito, 1977) by the absence of metalophulid II on the m1; the poor-developed ectolophs on the M1. Differs from *H. bijugatus* Mein and Freudenthal, 1971a by the m1 without metalophulids II and the absence of enamel coated valley. Differs from *H. decedens* (Schaub, 1925) by its sinuous contour on the M1 and the absence of metalophulid II. Differs from *H. castelnovi* Aguilar, Clavet and Michaux, 1994 by its bigger size. Differs from *H. nombrevillae* (Freudenthal, 1966); *H. adroveri* Agustí, 1986; *H. baixasi* Aguilar, Michaux and Lazzari, 2007; *H. peralensis* Van de Weerd, 1976; *H. moralesi* López-Antoñanzas, Peláez-Campomanes, Álvarez-Sierra and García-Paredes, 2010; *H. mediterraneus*, Aguilar 1982; *H. thaleri* (Hartemberger, 1965), and *H. aragonensis* (Freudenthal, 1966) by its non-complete ectolophs.

CRICETODON NIEVEI FROM THE CALATAYUD-DAROCA BASIN

Description of the Type Locality TOR3A

M1—Material: 18 specimens. It has four roots and a sinuous outline. The enamel is thick (Fig. 7.4B). The anterocone is divided in two parts by a shallow groove on the anterior wall of

the tooth. A labial anteroloph is present in five out of 14 molars; this anteroloph is joined to the labial part of the anterocone and is extended through the protosinus, but without reaching the protocone (Fig. 7.4B, 4C). The anterolophule is connected to the lingual part of the anterocone. A short labial spur of the anterolophule is present in one specimen. The protolophule is always posterior. The posterior ectoloph of the anterocone is present in most of the cases (10/12; Fig. 7.4A, 4B). In one case, this anterior ectoloph is well developed and it reaches the paracone. The posterior ectoloph is always present; it is simple, composed by the posterior ectoloph of the paracone, in most specimens (11/12; Fig. 7.4A, 4C) and it is double, composed by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone, in one case. None of them is complete. The mesoloph is present in all specimens but one (14/15), being short in most cases (13/15; Fig. 7.4A, 4B, 4C); it is reduced to an enlargement of the enamel (incipient) in one case. A ridge-like structure is present in the anterior wall of the protocone in six out of 14 specimens (Fig. 7.4A, 4B), being absent in the remaining ones (8/14; Fig. 7.4C). The enamel coated valley is absent. The entomesoloph is present in some specimens (2/13) and it is short (Fig. 7.4B). The labial posteroloph is always present, small, but well delimited (Fig. 7.4A, 4B, 4C). The lingual posteroloph is mostly present (8/10; Fig. 7.4A; 4C). The anterosinus is closed by a well-developed and large cingulum (10/12; Fig. 7.4A, 4B, 4C); by a style (1/12) or it can also remain open (1/12). The mesosinus is blocked by a big cingulum (10/12; Fig. 7.4A, 4B); it is open in two cases (Fig. 7.4C). The sinus is linguallly open (7/12; Fig. 7.4B) or closed either by a large style (3/12; Fig. 5.4C), or by two crests joined to the hypocone and the protocone (2/12). The protosinus is closed by a big style in 9 out of 14 molars (Fig. 7.4A); it is closed by the lingual anteroloph on the others (Fig. 7.4B, 4C).

M2—Material: 30 specimens. The anterior ectoloph is present in most cases (19/21); it is composed by the anterior ectoloph of the paracone in four molars and it seems to be an enlargement of the enamel at the base of the paracone in the others. (Fig. 7.4D). The posterior ectoloph is always present and composed by the posterior ectoloph of the paracone; it is never connected to the metacone (Fig. 7.4D) but in two cases, is linguallly curved and connected to the mesoloph (Fig. 7.4D). The mesoloph is short (15/27; Fig. 7.4D), incipient (5/27) or absent (7/27). A spur on the posterior wall of the protocone is present in nine out of 12 specimens (Fig. 7.4D). The entomesoloph is only present in one specimen and it is incipient. Labial posteroloph is always present; it is well developed in most specimens (18/23; Fig. 7.4D) and weak in the remaining ones. The lingual posteroloph is short (10/23) or absent in (13/23; Fig. 7.4D). The anterosinus can be closed by the connection of the basis of labial anteroloph and the paracone (25/26). The protosinus is weak in all the molars but one; it is closed by a small style in two out of 24 specimens. The mesosinus is blocked by a cingulum (22/27) or remains open (5/27); the cingulum is low in two specimens (Fig. 7.4D). The sinus is curved forward (9/12; Fig. 7.4D) or straight (3/12). The sinus is open (19/24; Fig. 7.4D) or closed (5/24); it is blocked by a thick cingulum (2/5) or by two crests

attached to the hypocone and to the protocone (3/5).

M3—Material: 27 specimens. It has a rounded outline with a reduced hypocone. The lingual anteroloph is weak although it is clearly distinguishable in most cases (6/7; Fig. 7.4E, 4F). The labial anteroloph is always present and well developed (Fig. 7.4E, 4F). The anterior ectoloph is barely present (5/23); it is composed by the anterior ectoloph of the paracone and one is complete. The posterior ectoloph is always present; it is double (11/24; Fig. 7.4F) or single, formed by the posterior ectoloph of the paracone (13/24; Fig. 7.4E). In six specimens the posterior ectoloph is complete. Some molars (4/22) have a short mesoloph. The enamel coated valley is absent. The entoloph is continuous and the neo-entoloph is present in five out of 23 cases. The posteroloph is absent; several specimens (3/20) have a constriction on the posterior wall of the hypocone which is not considered as a true posteroloph (Fig. 7.4E). The sinus is deep, curved and directed forward in the specimens without neo-entoloph (Fig. 7.4E, 4F); it is small and straight if the neo-entoloph is present.

m1—Material: 36 specimens. The labial anterolophid is well developed and usually does not reach the protoconid (16/25; Fig. 7.4H, 4I, 4J); it connects the anteroconid with the protoconid in the rest of cases (Fig. 7.4H, 4I). A lingual anterolophid is present in one out of 25 specimens and it does not reach the metaconid. Most of the molars (22/24) have a metalophulid I (Fig. 7.4G, 4H, 4I, 4J) and the remaining ones do not have any metalophulid, the metalophulid II being always absent. The longitudinal ridge bears a mesolophid in nearly half of the specimens (11/21; Fig. 7.4G, 4I), and it is clearly distinguishable but short in all cases but one, which is incipient. The ectomesolophid is usually present (16/26); it is long, reaching the labial border (2/16; Fig. 7.4G, 4H, 4J) or ending before the labial edge (7/16), and it is short in seven out of 16 specimens (Fig. 7.4I). The metaconid and the entoconid have, in general, posteriorly directed protuberance near the lingual border (Fig. 7.4G, 4H, 4I). The hypolophulid is transversal (3/27; Fig. 7.4I) or strongly directed forward (24/27; Fig. 7.4G, 4H, 4J). A small labial posterior cingulum is present in three out of 26 cases (Fig. 7.4G). The mesosinusid is blocked by a low cingular ridge (3/11; Fig. 7.4G, 4I); closed by a stylid (3/11) or it is open (5/11; Fig. 7.4H, 4I, 4J). The sinusid is wide and shallow; it is straight and it is blocked by a low cingular ridge (15/19; Fig. 7.4H), by two crests attached to the hypoconid and the protoconid (1/19), or by a stylid with a crest attached to the hypocone (2/19). The sinusid is open in one molar. The posterosinusid is blocked by a cingular ridge (3/26) or it remains open.

m2—Material: 24 specimens. It has two roots; the posterior one is split in the half height in some cases (6/10), but none of the specimens have three complete roots. The labial branch of the anterolophid is well developed and it reaches the protoconid (Fig. 7.4L, 4M) in all cases but one (Fig. 7.4K). The lingual anterolophid is absent (Fig. 7.4K, 4L, 4M). Most specimens have a mesoloph (20/24); it is short (19/20; Fig. 7.4K, 4L, 4M) and the other one present a very reduced

form; it is an enlargement of the enamel instead of a “true” mesolophid. Several teeth possess an ectomesolophid (11/23); it is short (3/11; Fig. 7.4K) or long and reaching the labial border (8/23; Fig. 7.4M). There is no anterosinusid and the mesosinusid is open (7/24; Fig. 7.4K, 4L) or blocked by a low cingulum (17/24; Fig. 7.4M). The sinusid is open (2/23), blocked by a low cingular ridge (14/23; Fig. 7.4K, 4L), by two crests related to the protoconoid and the hypoconid (6/23; Fig. 7.4K), or by a stylid (1/23). A weak labial posterosinusid is formed on the posterior wall of the hypoconid in most specimens (19/21; Fig. 7.4M); it is well distinguishable in five of them. The posterosinusid can be blocked by a cingulid (3/21) or by the connection between the posterolophid and the entoconid (8/21); it is open in the remaining ten specimens (Fig. 7.4K, 4L, 4M).

m3—Material: 24 specimens. The labial anterolophid is long; it reaches the protoconid in 16 out of 19 cases (Fig. 7.4N). A weak lingual anterolophid is present in one case. The mesolophid is present in most specimens (17/19); it is short (Fig. 7.4N) and in four teeth it develops a second branch in the ectolophid (Fig. 7.4O). The ectomesolophid is mostly absent (Fig. 7.4O); but one tooth possess a weak enlargement of the enamel (Fig. 7.4N). The lingual posterolophid is high and extends towards the entoconid. The mesosinus is closed by a low cingulid (11/18; Fig. 7.4N, 4O) or it remains open (7/18). The sinusid is blocked by a cingulid (7/16; Fig. 7.4N), by a stylid (7/16) or it remains open (3/16; Fig. 7.4O). The posterosinusid is closed by a cingulum (6/18) or it is blocked by the posterolophid (7/16; Fig. 7.4O); it is open in the remaining specimens (5/18; Fig. 7.4N).

***Cricetodon nievei* sp. nov. from other sites in the Calatayud-Daroca Basin.**

The studied assemblages of *Cricetodon nievei* sp. nov. from TOR2, 3A, 3B and LP5H (Table 7.1) are morphological and metrically homogeneous (Figs. 7.5–7.6). Most characteristic features present in all fossil sites are: the thick enamel, the presence of styles on the protosinus of the M1, the moderate development of the ectolophs (Fig. 7.5A; 5D; 5F)—being incomplete in most specimens—, the moderately simple pattern of the M3, the presence of metalophulid I on the m1 and absence of metalophulid II (Fig. 7.6A; 6D), the presence of ectomesolophid, and the moderately reduced m3 (Table 7.2).

Although no remarkable differences in morphology are observed in the studied material, there is certain degree of variability in several characters, or state of character, which are absent in the type locality. In the fossils from LP5H, the anterior ectoloph is connected to the basis of the paracone in the M1 (3/6) and M2 (1/9), instead of to the top of the cusp as in the rest. Regarding the M3, LP5H and TOR3B have some specimens with longer mesolophs that can exceed the half of the length of the mesosinus or even reach the labial border (1/3 in LP5H).

On the other hand, *C. nievei* sp. nov. from the type locality do not show significant differences in the metrical variables when compared to other studied samples (Fig. 7.3; Table 7.3).

COMPARISONS AND DISCUSSION

Remarks on the genera *Cricetodon* and *Hispanomys*

Before addressing comprehensive comparisons and discussion, it is crucial to clarify some aspects about the genera *Cricetodon* and *Hispanomys*. Many authors (De Bruijn, 1976; Sen and Ünay, 1979; Ünay, 1980; Ünay and De Bruijn, 1984; De Bruijn et al., 1993; Casanovas-Vilar, 2007; Prieto et al., 2010) pointed out that the differences between some genera of Cricetodontini are diffuse because the limit from one to the other is not well defined. These transitions are especially problematic during the late Aragonian: from *Cricetodon* and *Hispanomys* in Western Europe as well as from *Cricetodon* to *Byzantinia* in the Eastern Mediterranean. Thus, the generic assignment is sometimes very complicated (Prieto et al., 2010).

The genus *Cricetodon* is described by the following features: large, medium or small sized Cricetodontini with low-crowned cheek teeth, ectolophs of the upper molars are absent or weak, the upper M1 has four roots, anterocone of the M1 single or double and majority of the m1 with metalophulid I and II present on the same specimen or metalophulid II only, among others (Mein & Freudenthal, 1971a; De Bruijn et al., 1993). In turn, *Hispanomys* is described as Cricetodontinae of medium size, slight to moderately hypsodont; with either incomplete or complete ectolophs; the anterocone of the M1 with an anterior groove and the M1 with either four or five roots (Van de Weerd, 1976). Regarding the presence of metalophulid II, Prieto et al. (2010) pointed out that it is usually absent in the early *Hispanomys*. However, the same authors did not recommend the use of this structure for the unambiguous separation between *Cricetodon* and *Hispanomys* due to its high morphological variability.

De Bruijn and Ünay (1996) recognized five evolutionary grades of Cricetodontini. Type 1, 2 and 4 are composed exclusively by the species from Anatolia; type 1 and 2 are represented by the basal eastern *Cricetodon* species, whereas type 4 includes mainly representatives of *Byzantinia*. The type 3 comprises most of the species of *Cricetodon* characterized by: M1 and M2 with four roots; a divided anterocone; a posterior ectoloph that varies in length and it may be complete; a short labial posteroloph and an m1 that has usually metalophulid I and II in the same specimen (De Bruijn and Ünay, 1996). Finally, the type 5 contains the Cricetodontini from south-western Europe, assigned by these authors to *Hispanomys* and *Ruscinomys*; the species included here display

the following features: frequent presence of five roots in the M1; check teeth, not “stretched” longitudinally and wear flat, especially the third molar.

Prieto et al. (2010) defined six groups of Cricetodontini as well. Group 1 that comprises the middle Miocene Anatolian species of *Cricetodon*. Group 2, including middle Miocene *Cricetodon hungaricus* from the localities from Hungary. Group 3 is including *Cricetodon klariankae* from the late middle Miocene of Hungary. Prieto et al. (2010) proposed an ancestor-descendant relationship between *C. klariankae* and ‘*Cricetodon*’ *fandli* from Austria, so the latter species is also considered in the group 3. Group 4 that includes the early *Hispanomys* species (middle Miocene to earliest late Miocene) from Spain and France; it is characterized by having less hypsodont and reduced longitudinally “stretched” molars (Prieto et al., 2010). Group 5, which contains brachiodont species of *Cricetodon* from the middle Miocene characterized by ectolophs that are very short or absent (Prieto et al., 2010: 181). And finally, group 6 composed by *Cricetodon caucasicus* from Georgia, which shows non-split anterocone and no ectolophs. However, Pickford et al. (2000) described complete ectolophs in *Cricetodon caucasicus* and its inclusion in group 3 is now taking in consideration by the author (Prieto, pers. comm.).

The proposal of all these classifications that attempt to order the different groups of species of *Cricetodon* and *Hispanomys*, give us an idea about the difficulties that, sometimes, are found when realizing the generic assignation. This problem is particularly noticeable during the late Aragonian, the time interval in which the material studied here is placed.

Comparison with others species of Cricetodontini.

The material of Cricetodontini from TOR2, TOR3A and TOR3B was previously cited and studied by Álvarez Sierra et al. (2003) and López-Guerrero et al. (2009). These authors left the taxonomical assignation as *Cricetodon* spp. and *Cricetodon* aff. *jotae*, respectively. After a new study and descriptions of this material, we have confirmed that the combination of characters is unique and it has not previously been described, allowing us to propose the new taxon.

As we mention before, the generic assignation of the species of Cricetodontini from the late Aragonian is controversial (De Bruijn et al., 1993; Prieto et al., 2010; López-Guerrero et al., 2013) and as noted by Prieto et al. (2010: 425), it ‘depends on whether the morphological or phylogenetic approach is used’. Taking into account these considerations, we discuss, on the first place, the generic ascription of the studied fossils.

The specimens from TOR3A have a combination of primitive and derived characters. The upper molars have the following features: (1) low crowns, (2) sinuous contour, (3) moderate

development of the ectolophs which do not form a continuous wall, (4) presence of mesolophs and entomesolophs, cingular formations and styles on the valleys, (5) Four-rooted M1, and (6) a barely reduced M3. These characteristics are frequently present in *Cricetodon* and they are considered as primitive for the tribe (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; De Bruijn et al., 1993; López-Antoñanzas and Mein, 2009). On the other hand, the lower molars have: (1) short or absent mesolophids, (2) absence of metalophulid II in the m1, and (3) three incipient roots in the m2. These characters are described in many species of *Hispanomys* and are considered as derived within the tribe Cricetodontini (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; De Bruijn et al., 1993; López-Antoñanzas and Mein, 2009, López Antoñanzas et al., 2010). However, the presence of metalophulid I and the absence of metalophulid II in the m1, as well as the three rooted m2 are also found in some species of *Cricetodon* such as: *Cricetodon albanensis*, *C. bolligeri* and *C. jumaensis* studied by Mein and Freudenthal (1971a) and Rummel (1995, 2001), respectively.

In addition, the diagnosis and the descriptions of the different species of *Hispanomys* frequently pointed out the high hypsodonty (Mein and Freudenthal, 1971a; Van de Weerd, 1976); complete and usually double ectolophs forming a continuous wall (Aguilar et al., 1994); the younger species show a short or absent labial anterolophid in the m1 (López-Antoñanzas and Mein, 2009, 2011); and a hypocone highly reduced in the third molars (Agustí, 1982). These characters are absent in the material from TOR3A.

Given that our material presents a large number of characters of *Cricetodon*, as well as the fact that the characters shared with *Hispanomys* are also found in some species assigned to *Cricetodon*, we decided to include the new species of TOR3A in the latter genus; despite of the abovementioned resemblance of its lower molars with *Hispanomys*. Nevertheless, it is essential to compare our new material with the species included in both genera.

In order to make easier the comparisons with the species of *Cricetodon* and *Hispanomys*, these were grouped. Primary order is by age and secondary sort is by size. The species for each group differ from *C. nievei* sp. nov. by at least one of the listed features or a combination of several ones.

Firstly, the material of TOR3A is compared with the older species of *Cricetodon*. Since the Ramblian until the middle Aragonian, the Cricetodontini remains modest in numbers of both specimens and species (De Bruijn and Ünay, 1996). These species are enclosed in the Cricetodontini types 1, 2 and partially 3 defined by De Bruijn and Ünay (1996) as well as in the groups 1, 2 and 6 by Prieto et al. (2010). We include in these groups: *Cricetodon tobieni*, *C. wanhei*, *C. soriae*, *C. aureus*, *C. sansaniensis*, *C. hungaricus*, *C. candirensis*, *C. versteegi*, *C. aliveriensis*, *C. meini*, *C. pasalarensis*, *C. caucasicus*, *C. volkeri* and *C. kasapligili*. Our new species can be easily

differentiated from these species because they possess some of the follow primitive traits: (1) M1 with non-divided anterocone; (2) three roots on the M1; (3) anterior protolophule; and (4) m1 with metalophulid II or both metalophulids, I and II, in the same specimen (Mein and Freudenthal, 1971b; Baudelot, 1972; Klein Hofmeijer and De Bruijn, 1988; De Bruijn et al., 1993; Rummel, 2001; Hernández-Fernández et al., 2006; Hír, 2007; Wu et al., 2009; Qiu, 2010; Maridet and Sen, 2012; López-Guerrero et al., 2013).

In turn, the younger representatives of Cricetodontini, from the Vallesian and the Turolian, belong to the type 5 defined by De Bruijn and Ünay (1996). In this group we include: *Hispanomys adroveri*, *H. freudenthali*, *H. baixasi*, *H. peralensis*, *H. moralesi*, *H. mediterraneus*, *H. thaleri*, *H. aragonensis* and *H. nombrevillae*. These species present complete ectolophs and high degree of hypsodonty. None of these characters are present in the studied material and they confer a clear derivate pattern, according with the trends described by the authors (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas & Mein, 2009).

Our material is compared with the contemporary species from the late Aragonian. Among them, there is a group that is clearly larger than the material from TOR3A (Tables 4–5). This group includes: (A) *C. engesseri*; (B) *C. klariankae* (C) *C. jumaensis*; (D) *C. cariensis*; (E) *C. albanensis*, and (F) *Hispanomys lavocati*. They have also some different features. Each species have, at least, one or more of the following features: (1) long mesolophs and mesolophids (C); (2) complete ectolophs (B, D, F); (3) enamel coated valley in the upper molars (D, E); (4) presence of metalophulid II (B, C, D, E, F); and (5) well-developed posteroloph on the M3 (A) (Sen and Ünay, 1979; Rummel, 2001; Rummel and Kálin, 2003; Hír, 2007; Kálin and Kempf, 2009; López-Guerrero et al., 2008).

The case of *C. albanensis* from La Grive M (MN7/8) (Mein and Freudenthal, 1971a; De Bruijn et al., 1992) is especially interesting. Despite its larger size, the dental proportions are similar to those in TOR3A. Besides, the morphological resemblance with our material is noticeable. Both species share: the low crowned cheek teeth, the degree of development of ectolophs, which are single and not complete; the presence of large styles on the protosinus; the absence of metalophulid II; the presence of two labial bulges on the entoconid and metaconid in the m1 and the presence of a bifid anterolophulid in some m1. However, *C. albanensis* has a number of traits that differs from the studied material, such as: enamel coated valley on the upper molars, and the absence of both mesoloph and ectomesolophid.

There is another set of species from the late Aragonian which has similar size to that of TOR3A (Tables 7.4, 7.5), but displays different morphology. '*Cricetodon*' *fandli*, *C. jotae*, *H. bijugatus* and *H. aguirrei* differ from the *Cricetodon* from TOR3A, because they have metalophulid II or both metalophulids on the same specimen in the m1 (Sesé Benito, 1977; Prieto et al., 2010;

López-Guerrero et al., 2008; López-Antoñanzas and Mein, 2009, 2011). *Cricetodon bolligeri* has anterior protolophule on the M1 and a more developed entomesolophid on the lower molars (Rummel, 1995). *Hispanomys dispectus* possess double and complete ectolophs and a straight contour (Agustí, 1980). *Hispanomys daamsi* has no mesoloph; the anterior and posterior ectolophs are complete and the latter is double (Agustí et al., 2005).

The case of *Hispanomys decedens* needs special attention since it shows quite basal morphology shared with the molars from TOR3A such as: the presence of lingual cingulum closing the valleys, incomplete anterior ectolophs, the presence of mesolophs, four-rooted M1, and low-reduced M3 (López-Antoñanzas and Mein, 2011). Despite these similarities, *H. decedens* shows some differences like the enamel coated valley, complete posterior ectolophs in the upper molars, as well as a well-developed posteroloph on the M3 (López-Antoñanzas and Mein, 2011), which preclude the assignment of our material to *H. decedens*.

Finally, there is one species that is smaller than the molars from TOR3A (Tables 7.4, 7.5), *H. castelnovi*. This species displays also morphological differences with the studied fossils; it has the anterocone slightly divided in the M1 and both ectolophs are complete, forming a continuous wall in the upper molars (Aguilar et al., 1994).

A large number of forms of Cricetodontini have been described from the late Aragonian and early Vallesian in central Europe. In the Swiss Molasse of the North Alpine Foreland Basin (NAFB) *Hispanomys* sp. from Nebelberg TGL II+III (MN9) (Kälin and Kempf, 2009) is similar in size and morphology to the material from TOR3A. Both possess M1 with weak posterior ectoloph and absence of anterior ectoloph, and m1 having mesolophid and metalophulid I, but lacking metalophulid II (Kälin and Kempf, 2009; Rummel and Kälin, 2003). These resemblances indicate that the Swiss specimens could fit the diagnosis of the new species from TOR3A; however, the revision and study of this material is necessary to confirm this statement.

Rummel (2000) reported *Cricetodon* aff. *sansaniensis* from the locality Petersbuch 31 (MN7) in the German Molasse of the NAFB. It displays several similarities with the species from TOR3A: both have weak-developed and not complete ectolophs on the upper molars, short mesolophids on the m1 and, generally, three roots on the m2: the posterior one can be completely or partially split (Rummel, 2000). Furthermore, *Cricetodon* aff. *sansaniensis* is similar in size to the species from TOR3A, although the m2 and m3 are slightly larger and, therefore, its Length m1/Length m3 index is lower (1.08 vs. 1.12, respectively). Besides, it shows metalophulid II in one out of six specimens. All these features preclude the assignation of *Cricetodon* aff. *sansaniensis* to the species from TOR3A.

After the detailed morphometrical study and the comparison with all species above mentioned, it is justified the proposal of the new species *Cricetodon nievei* sp. nov. for the material

from TOR3A, TOR3B, TOR2 and LP5H.




FIGURE 7.3. Length/Width scatter-diagram of the cheek teeth of *Cricetodon nievei* sp. nov. and *Cricetodon jotae* from Manchones.




FIGURE 7.4. *Cricetodon nievei* sp. nov. from the Calatayud-Daroca Basin. Toril 3A. A, inverted right M1, TOR3A-126; B, inverted right M1, TOR3A-127; C, left M1, TOR3A-122; D, inverted right M2, TOR3A-159; E, left M3, TOR3A-219; F, left M3 TOR3A-260. G, left m1, TOR3A-318; H, inverted right m1, TOR3A-322 Holotype; I, inverted right m1, TOR3A-169; J, left m1, TOR3A-176; K, left m2, TOR3A-327; L, inverted right m2, TOR3A-201; M, inverted right m2, TOR3A-207; N, inverted right m3, TOR3A-259; O, inverted right m3, TOR3A-242. Scale bar represent 1 mm.




FIGURE 7.5. *Cricetodon nievei* sp. nov. from the Calatayud-Daroca Basin. Upper molars. Toril 3B: A, left M1, TOR3-143; B, inverted right M2, TOR3-146; C, left M3, TOR3-148; D, left M1, TOR3B-402; E, left M2, TOR3-147. Toril 2: F, left M1, TOR2-258; G, inverted right M2, TOR2-264; H, left M3, TOR2-267. Scale bar represent 1 mm.




FIGURE 7.6. *Cricetodon nievei* sp. nov. from the Calatayud-Daroca Basin. Lower molars. Toril 3B: A, inverted right m1, TOR3B-420; B, inverted right m2, TOR3B-426; C, left m3, TOR3B-433; D, inverted right m1, TOR3B-418; E left m2, TOR3B-431; F, left m3, TOR3B-432. Toril 2: G, left m2, TOR2-271; H, inverted right m3, TOR2-274. Scale bars equals 0.5 mm and 1mm.

FINAL REMARKS & CONCLUSIONS

Relationships with the other *Cricetodontini* present in the Calatayud-Daroca Basin

The first *Cricetodontini* from the Calatayud-Daroca Basin is found in the local biozone E (MN5, middle Aragonian). During biozones F, G1 and G2 (MN6, late Aragonian) the species have a homogeneous basal morphology characteristic of *Cricetodon*. In the biozone G3 (MN7/8, latest late Aragonian) there is an increase of the taxonomical diversity, being registered the highest number of species in the basin. The morphological homogeneity disappears and the intraspecific variability became higher. These species show a combination of primitive and derived characters. Afterwards, the acquisition of derivate characters typical of *Hispanomys* is achieved during the local biozone H (Álvarez-Sierra et al., 2003).

In the scenarios described above, *C. nievei* sp. nov. is registered on the interval of maximum diversification together with *Cricetodon jotae*, *Hispanomys aguirrei* and *H. lavocati*. *Cricetodon jotae* is found in Manchones, Arroyo del Val 6 (local biozone G2, MN6) and TOR 1 (local biozone G3, MN7/8) (Mein and Freudenthal, 1971a; pers. obs.). *Hispanomys aguirrei* and/or *H. lavocati* are recorded on the younger localities from the TOR-Nombrevilla section, Nombrevilla 2, 3 and 4 (López-Guerrero et al., 2008, 2009).

Agustí (1982) propose a phylogeny for the *Cricetodon*-*Ruscinomys* group, in which *C. jotae* was the ancestor of *H. aguirrei*. He pointed out that the evolutionary trends for this lineage were the development of the ectolophs and the acquisition of a slightly modern pattern consisting of: the increase of the hypsodonty and the reduction of the cingular formations (Agustí, 1982: 104, 110). Given that *Cricetodon nievei* sp. nov. is located in the sites of intermediate age between those which have *C. jotae* and *H. aguirrei*, is necessary to analyse the possible relationships that could exist among this three species.

As we remark before, these three species show, each one, a particular combination of basal and derivate traits. In the case of *C. jotae* from Manchones we observe a primitive character as the metalophulid II present in many specimens and, however, it has the following features that are considered derivate for the tribe (Agustí, 1980; López-Antoñanzas and Mein, 2009): absence of mesolophs; presence of double ectolophs and styles in the protosinus of only few M1; low-developed or absent mesolophids; absence of ectomesolophid and weak or absent

labial anterolophid of the m1. In contrast, *Cricetodon nievei* sp. nov. has mesoloph/ids; only one M1 has double ectolophs, styles are more frequent on the protosinus; ectomesolophid present; well-developed labial anterolophid, and metalophulid I in the m1; furthermore, neo-entolophs are present the M3 and it present bigger size (Fig. 7.3).

In the case of *H. aguirrei*, it displays primitive features like: the anterocone on the M1 poorly divided; presence of metalophulids I and II in the same specimen; two roots in the m2 (López Martínez et al., 1977; Sesé Benito, 1977; López-Guerrero et al., 2008), whereas both *C. jotae* and *C. nievei* sp. nov. possess well-divided anterocone and some of their m2 have three roots. But, on the other hand, the ectolophs are complete in a high number of molars (Sesé Benito, 1977; López-Guerrero et al., 2008) and it is a derived feature absent in *C. jotae* as well as in *C. nievei*, sp. nov.

The observed morphology, combining primitive and derivate traits in a particular manner for each species, suggests a mosaic evolution. This consists in different rates of change in the jugal teeth which result in a degree of uncoupling between the upper and lower features (Stebbins, 1983; López Martínez, 1997; Hopkins and Lidgard, 2012). Furthermore, this mosaic evolution is also recognized in other Cricetodontini from the European late Aragonian (De Bruijn et al., 1993; De Bruijn and Ünay, 1996). This fact supposes an additional difficulty for establishing the phylogenetic relationships between the involved taxa *C. jotae*, *C. nievei* sp. nov. and *H. aguirrei*. Therefore, these questions need a separate study, which is beyond the scope of this study, and they should be resolved through the cladistic analysis that is currently in progress. In line with this, a previous cladistics phylogeny, (Sen and Erbajeva, 2011) propose that *Cricetodon* might be paraphyletic, since its species were grouped in different clades. The resulting cladogram allocates *C. albanensis* in a clade with *Byzantinia*, whereas the other European species like *C. jotae* and *C. sansaniensis* are grouped on another clade together with some Asian taxa. Nonetheless, the phylogeny made by Sen and Erbajeva (2011) do not include *Hispanomys* and these results has to be taken carefully.

After making detailed comparisons with the rest of European and Asian species of *Cricetodon* and *Hispanomys* we are allowed to point out a morphological resemblance of *C. nievei* sp. nov. with the species of *Cricetodon* and *Hispanomys* from the Vallès-Penedès, France and central Europe, especially *C. albanensis* and *H. decedens*, that has to be assessed in a complete cladistics analysis.

Having discussed and compared *Cricetodon nievei* sp. nov., *C. jotae* and *Hispanomys aguirrei*, it is important to mention *Hispanomys lavocati* from Nombrevilla2, Nombrevilla 3 and Nombrevilla 4 (López-Guerrero et al., 2008, 2009). It shows a general pattern more derivate than *C. nievei*, sp. nov.; it consists of: ectolophs always present, some of them double and, frequently, complete; the metalophulid I always present, and the metalophulid II absent (pers. obs.); the m2

has always three roots and the posterior one is mostly completely split. The mosaic morphology is also perceived in this species by the presence of short mesoloph-ids in a considerably number of specimens (López-Guerrero et al., 2008). Nevertheless, De Bruijn et al. (1993) remark that *H. lavocati* presents some morphological affinities with the representatives from Anatolia (not specifying which ones) and they suggest that it could be related with a late migration from Asia Minor into Europe during the late Aragonian.

As it is described above, local biozone G3 reports the higher number of species of Cricetodontini. This increase has been described, not only for this tribe, but also for large mammals such as ursids (Abella et al., 2011) and ruminants (Sánchez and Morales, 2006, 2008; DeMiguel et al., 2011). The latter ones also reflect a dietary shift towards less abrasive diets (DeMiguel et al., 2011). Many authors argued that all this changes in the faunal assemblages had been related with an increase of humidity (Van der Meulen and Daams, 1992; Daams et al., 1999c; Alcalá et al., 2000; Álvarez Sierra et al., 2003; DeMiguel et al., 2011) and more diverse vegetation during the G3 (DeMiguel et al., 2011). In Addition, the occurrence of *C. nievei* sp. nov. in TOR3A is coincident with the re-appearance of beavers which supposed the presence of steady water courses (Alcalá et al., 2000; López-Guerrero et al., 2007) agreeing with the proposed increase of humidity. The new environmental conditions established in Calatayud-Daroca Basin seem to be very similar to those from the Vallés-Penedés Basin and Southern France. The implication of these changes in the evolution of Cricetodontini has to be evaluated in the future.

The present work has allowed us to increase the knowledge of the species of Cricetodontini during their maximum diversification interval, as well as to precise their stratigraphical and geographical distribution. Nevertheless, some aspects about the systematics and phylogeny of Cricetodontini remain to be solved; especially regarding those species from the late Aragonian. Indeed, the performance of a cladistic analysis, which is in process, should help us to precise the phylogenetic relationships and evolution of this group.

Locality		Length					Width					L/W
		Min	Mean	Max	SD	N	Min	Mean	Max	SD	N	
M1	TOR3B	3.02	3.18	3.38	0.140	5	1.96	2.10	2.27	0.120	6	1.51
	TOR3A	2.98	3.15	3.28	0.088	11	2.00	2.12	2.20	0.064	12	1.49
	TOR2	3.11	3.19	3.33	0.119	3	2.11	2.15	2.23	0.069	3	1.48
	LP5H	2.97	3.05	3.19	0.085	6	1.99	2.08	2.16	0.071	6	1.47
M2	TOR3B	2.12	2.22	2.34	0.084	6	1.82	1.94	2.15	0.010	8	1.14
	TOR3A	2.13	2.31	2.53	0.097	26	1.76	1.96	2.21	0.119	26	1.18
	TOR2	2.28	2.35	2.50	0.094	5	1.91	2	2.09	0.074	5	1.18
	LP5H	2.15	2.26	2.41	0.077	8	1.77	1.96	2.11	0.103	8	1.15
M3	TOR3B	1.85	1.88	1.94	0.052	3	1.54	1.78	1.90	0.165	4	1.06
	TOR3A	1.63	1.85	2.03	0.104	24	1.62	1.82	1.98	0.097	24	1.02
	TOR2	1.66	1.80	1.93	0.191	2	1.68	1.80	1.92	0.170	2	1.00
	LP5H	1.75	1.88	2.03	0.119	4	1.68	1.78	1.99	0.144	4	1.06
m1	TOR3B	2.58	2.66	2.80	0.074	6	1.65	1.74	1.95	0.109	6	1.53
	TOR3A	2.32	2.64	2.85	0.111	29	1.27	1.74	1.91	0.074	27	1.52
	LP5H	2.49	2.58	2.73	0.088	6	1.66	1.74	1.81	0.055	7	1.48
m2	TOR3B	2.29	2.42	2.53	0.100	4	1.92	1.98	2.14	0.091	5	1.22
	TOR3A	2.43	2.54	2.69	0.084	16	1.89	2.02	2.17	0.088	19	1.26
	TOR2	2.46	2.50	2.53	0.049	2	1.89	1.96	2.03	0.070	3	1.28
	LP5H	2	2.41	2.67	0.102	8	1.58	1.94	2.12	0.081	7	1.24
m3	TOR3B	2.17	2.34	2.49	0.130	5	1.77	1.80	1.88	0.044	5	1.30
	TOR3A	2.14	2.33	2.54	0.113	18	1.63	1.84	1.97	0.007	18	1.27
	TOR2	2.11	2.22	2.32	0.148	2	1.75	1.76	1.76	0.098	2	1.26
	LP5H	2.16	2.25	2.29	0.049	6	1.73	1.77	1.86	0.046	6	1.27

TABLE 7.2. Length and width of the upper and lower molars *Cricetodon nievei* sp. nov. Units are given in mm. Min= minimum values; Max= maximum values; SD= standard deviation; N= number of specimens.

	Levenne's test				ANOVA							
	Length (P-value)		Width (P-value)		Length (P-value)		d.f.1	d.f.2	Width (P-value)		d.f.1	d.f.2
M1	0.321	NS	0.500	NS	0.154	NS	3	21	0.638	NS	3	23
M2	0.593	NS	0.169	NS	0.080	NS	3	41	0.820	NS	3	43
M3	0.715	NS	0.361	NS	0.802	NS	3	29	0.827	NS	3	30
m1	0.477	NS	0.498	NS	0.352	NS	2	39	0.657	NS	2	38
m2	0.774	NS	0.727	NS	0.101	NS	3	25	0.569	NS	3	29
m3	0.290	NS	0.064	NS	0.255	NS	3	27	0.205	NS	3	27

TABLE 7.3. Results of the Levenne's and ANOVA tests comparing length and width variables of *Cricetodon nievei* sp. nov. from the studied assemblages with at least five data. NS= Non-significant differences ($p > 0.05$); d.f.= degree of freedom.

	Species	Length				Width				Source of data
		Min	Mean	Max	N	Min	Mean	Max	N	
M1	<i>H. dispectus</i>	2.79	3.15	3.46	27	1.77	1.98	2.24	27	Agustí, 1980
	<i>H. daamsi</i>	3.01	3.36	3.58	3	1.70	1.92	2.04	5	Agustí et al., 2005
	<i>H. lavocati</i>	3.56	3.69	3.88	6	2.09	2.22	2.34	7	Agustí, 1980
	<i>H. aguirrei</i>	2.68	3.08	3.50	89	1.81	2.10	2.27	89	Sesé Benito, 1977
	<i>H. bijugatus</i>	2.97	3.23	3.50	49	1.92	2.05	2.18	49	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	2.74	3.16	3.59	62	1.76	2.05	2.31	62	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	2.66	2.88	3.06	8	1.80	1.94	2.12	8	Aguilar et al., 1994
	<i>'C'. fandli</i>	2.90	3.04	3.20	4	1.90	1.92	1.97	4	Prieto et al., 2010
	<i>C. bolligeri</i>	2.96	3.20	3.44	89	1.85	2.02	2.20	89	Rummel, 1995
	<i>C. cariensis</i>	3.25	3.33	3.40	3	1.85	2.03	2.15	3	Sen and Ünay, 1979
	<i>C. jotae</i>	2.69	2.96	3.18	19	1.80	2.00	2.18	19	pers. obs.
	<i>C. pasalarensis</i>	2.76	2.94	3.21	19	1.69	1.85	1.99	19	Rummel, 1998
	<i>C. meini</i>	2.58	2.90	3.13	105	1.82	1.96	2.11	105	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	2.85	3.10	3.36	21	1.84	1.99	2.22	21	Rummel, 1998
M2	<i>H. dispectus</i>	2.04	2.35	2.59	23	1.71	1.92	2.14	23	Agustí, 1980
	<i>H. daamsi</i>	2.25	2.61	2.75	6	1.83	1.97	2.20	9	Agustí et al., 2005
	<i>H. lavocati</i>	2.41	2.56	2.75	4	2.05	2.12	2.23	4	Agustí, 1980
	<i>H. aguirrei</i>	2.04	2.35	2.63	93	1.68	1.92	2.27	93	Sesé Benito, 1977
	<i>H. bijugatus</i>	2.30	2.47	2.61	42	1.80	1.92	2.07	42	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	2.14	2.43	2.69	54	1.72	1.92	2.23	54	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	2.06	2.18	2.29	15	1.61	1.74	1.85	15	Aguilar et al., 1994
	<i>'C'. fandli</i>	2.22	2.30	2.43	4	1.78	1.82	1.88	2	Prieto et al., 2010
	<i>C. bolligeri</i>	2.21	2.43	2.65	83	1.79	1.95	2.12	83	Rummel, 1995
	<i>C. cariensis</i>	2.65	2.70	2.75	2	1.80	1.90	2.00	2	Sen and Ünay, 1979
	<i>C. jotae</i>	2.08	2.24	2.41	8	1.76	1.89	2.01	8	pers. obs.
	<i>C. pasalarensis</i>	2.01	2.14	2.35	27	1.63	1.97	1.80	27	Rummel, 1998
	<i>C. meini</i>	2.04	2.24	2.41	106	1.71	1.92	2.12	106	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	2.03	2.26	2.53	17	1.73	1.89	2.04	17	Rummel, 1998
M3	<i>H. dispectus</i>	1.52	1.65	1.75	6	1.52	1.59	1.69	6	Agustí, 1980
	<i>H. daamsi</i>	1.83	1.95	2.01	6	1.73	1.78	2.04	7	Agustí et al., 2005
	<i>H. lavocati</i>	1.78	1.90	2.01	4	1.74	1.86	1.98	4	Agustí, 1980
	<i>H. aguirrei</i>	1.54	1.74	2.04	90	1.52	1.73	2.04	90	Sesé Benito, 1977
	<i>H. bijugatus</i>	1.80	1.97	2.18	41	1.57	1.70	1.93	41	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	1.64	1.82	2.09	26	1.50	1.68	1.90	26	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	1.45	1.57	1.72	25	1.34	1.52	1.65	25	Aguilar et al., 1994
	<i>'C'. fandli</i>	1.73	1.87	1.95	6	1.57	1.69	1.82	6	Prieto et al., 2010
	<i>C. bolligeri</i>	1.79	1.96	2.14	55	1.68	1.83	2.02	55	Rummel, 1995
	<i>C. cariensis</i>	—	2.45	—	1	—	1.95	—	1	Sen and Ünay, 1979
	<i>C. jotae</i>	1.69	1.87	2.00	9	1.60	1.74	1.83	9	pers. obs.
	<i>C. pasalarensis</i>	1.65	1.78	1.97	23	1.51	1.67	1.84	23	Rummel, 1998
	<i>C. meini</i>	1.68	1.88	2.08	133	1.62	1.76	1.91	133	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	1.85	2.00	2.25	14	1.58	1.69	1.79	14	Rummel, 1998

TABLE 7.4. Length and width of the upper molars of medium sized species of *Cricetodon* and the older species of *Hispantomys*. Units are given in mm. Min= minimum values; Max= maximum values; N= number of specimens.

	Species	Length					Width				Source of data
		Min	Mean	Max	N		Min	Mean	Max	N	
m1	<i>H. dispectus</i>	2.29	2.59	2.90	19		1.48	1.70	1.89	19	Agustí, 1980
	<i>H. daamsi</i>	2.45	2.67	2.94	9		1.55	1.72	1.95	10	Agustí et al., 2005
	<i>H. lavocati</i>	2.53	2.82	2.98	8		1.85	1.89	2.01	8	Agustí, 1980
	<i>H. aguirrei</i>	2.36	2.63	2.95	84		1.59	1.78	2.00	84	Sesé Benito, 1977
	<i>H. bijugatus</i>	2.44	2.60	2.80	50		1.54	1.63	1.81	50	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	2.44	2.60	2.80	29		1.60	1.69	1.78	29	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	2.27	2.39	2.60	10		1.47	1.57	1.70	12	Aguilar et al., 1994
	'C': <i>fandli</i>	2.35	2.43	2.55	2		1.55	1.66	1.77	4	Prieto et al., 2010
	<i>C. bolligeri</i>	2.35	2.60	2.79	61		1.53	1.86	1.72	61	Rummel, 1995
	<i>C. cariensis</i>	2.60	2.66	2.75	3		1.72	1.80	1.85	4	Sen and Ünay, 1979
	<i>C. jotae</i>	2.05	2.44	2.68	28		1.50	1.63	1.82	28	pers. obs.
	<i>C. pasalarensis</i>	2.22	2.42	2.57	16		1.46	1.60	1.70	16	Rummel, 1998
	<i>C. meini</i>	2.19	2.41	2.57	90		1.42	1.57	1.73	90	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	2.32	2.47	2.76	23		1.56	1.70	1.82	23	Rummel, 1998
m2	<i>H. dispectus</i>	2.29	2.59	2.90	19		1.48	1.70	1.89	41	Agustí, 1980
	<i>H. daamsi</i>	2.52	2.58	2.70	4		1.87	2.02	1.93	4	Agustí et al., 2005
	<i>H. lavocati</i>	2.46	2.66	2.81	11		1.97	2.07	2.21	11	Agustí, 1980
	<i>H. aguirrei</i>	2.18	2.48	2.77	101		1.77	2.02	2.27	101	Sesé Benito, 1977
	<i>H. bijugatus</i>	2.40	2.54	2.74	43		1.65	1.83	1.93	43	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	2.27	2.48	2.69	34		1.80	1.94	2.08	34	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	2.32	2.35	2.39	12		1.71	1.76	1.84	14	Aguilar et al., 1994
	'C': <i>fandli</i>	2.27	2.43	2.53	5		1.73	1.84	1.87	5	Prieto et al., 2010
	<i>C. bolligeri</i>	2.19	2.52	2.71	59		1.75	1.91	2.01	59	Rummel, 1995
	<i>C. cariensis</i>	2.65	2.68	2.70	4		1.90	2.02	2.10	5	Sen and Ünay, 1979
	<i>C. jotae</i>	2.22	2.39	2.65	27		1.72	1.87	2.07	27	pers. obs.
	<i>C. pasalarensis</i>	2.09	2.26	2.41	29		1.61	1.75	1.92	29	Rummel, 1998
	<i>C. meini</i>	2.14	2.33	2.51	88		1.71	1.83	2.05	88	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	2.21	2.40	2.58	17		1.70	1.82	1.90	17	Rummel, 1998
m3	<i>H. dispectus</i>	1.87	2.13	2.33	27		1.49	1.71	1.89	27	Agustí, 1980
	<i>H. daamsi</i>	2.38	2.47	2.60	4		1.72	1.88	1.98	5	Agustí et al., 2005
	<i>H. lavocati</i>	2.43	2.54	2.74	5		1.88	1.98	2.00	5	Agustí, 1980
	<i>H. aguirrei</i>	1.95	2.18	2.45	78		1.54	1.75	1.97	78	Sesé Benito, 1977
	<i>H. bijugatus</i>	2.32	2.47	2.72	39		1.58	1.68	1.80	39	López-Antoñanzas and Mein, 2009
	<i>H. decedens</i>	1.92	2.23	2.42	25		1.63	1.79	1.93	25	López-Antoñanzas and Mein, 2011
	<i>H. castelnovi</i>	1.90	2.09	2.20	5		1.57	1.64	1.74	5	Aguilar et al., 1994
	'C': <i>fandli</i>	2.08	2.23	2.32	7		1.60	1.69	1.78	7	Prieto et al., 2010
	<i>C. bolligeri</i>	2.16	2.36	2.63	44		1.76	1.88	2.03	44	Rummel, 1995
	<i>C. cariensis</i>	2.08	2.65	2.74	4		1.70	1.87	2.00	5	Sen and Ünay, 1979
	<i>C. jotae</i>	2.20	2.30	2.43	10		1.66	1.83	1.99	10	pers. obs.
	<i>C. pasalarensis</i>	2.10	2.30	2.50	32		1.48	1.66	1.89	32	Rummel, 1998
	<i>C. meini</i>	2.05	2.36	2.77	101		1.56	1.80	2.09	101	Mein and Freudenthal, 1971b
	<i>C. candirensis</i>	2.23	2.34	2.45	12		2.58	1.66	1.75	12	Rummel, 1998

TABLE 7.5. Length and width of the lower molars of medium sized species of *Cricetodon* and the older species of *Hispantomys*. Units are given in mm. Min= minimum values; Max= maximum values; N= number of specimens.

LITERATURE CITED

- Abella, J., P. Montoya, and J. Morales. 2011. A new species of *Agriarctos* (Ailuropodinae, Ursidae, Carnivora) in the locality of Nombrevilla 2 (Zaragoza, Spain). *Estudios Geológicos* 67:187–191.
- Aguilar, J. P. 1982. Contributions à l'étude des Micromammifères du gisement Miocène supérieur de Montredon (Hérault) 2. les rongeurs. *Palaeovertebrata* 12:75–140.
- Aguilar, J. P., M. Calvet, and J. Michaux. 1994. Les rongeurs de Castelnou 6 (Pyrénées-orientales, France) et les corrélations entre faunes ibériques et françaises au Miocène moyen. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 192:109–131.
- Aguilar, J. P., J. Michaux, and V. Lazzari. 2007. Lo Fournas 16-M (Miocène supérieur) et Lo Fournas 16-P (Pliocène moyen), deux nouvelles localités karstiques à Baixas, Sud de la France Partie II – Nouvelles espèces de rongeurs, listes fauniques et remarque sur l'utilisation biochronologique des faunes karstiques. *Géologie de la France* 1:63–81.
- Agustí, J. 1980. La asociación de *Hispanomys* y *Cricetodon* (Rodentia, Mammalia) en el Mioceno superior del Vallès-Penedès (Cataluña, España). *Acta Geologica Hispanica* 15:51–60.
- Agustí, J. 1982. Tendencias evolutivas de la línea *Cricetodon-Ruscinomys* (Rodentia, Mammalia) en la Península Ibérica. *Acta Geologica Hispanica* 17:103–111.
- Agustí, J. 1986. Nouvelles especes de cricetides vicariantes dans le Turolien moyen de Fortuna (Prov. Murcia, Espagne). *Geobios* 19:5–11.
- Agustí, J., I. Casanovas-Vilar, and M. Furió. 2005. Rodents, insectivores and chiropterans (Mammalia) from the late Aragonian of Can Missert (middle Miocene, Vallès-Penedès Basin, Spain). *Geobios* 38:575–583.
- Alcalá, L., A. M. Alonso-Zarza, M. A. Álvarez-Sierra, B. Azanza, J. P. Calvo, J. C. Cañaveras, J. V. Dam, M. Garcés, W. Krijgsman, A. J. Van der Meulen, P. Péláez-Campomanes, A. Pérez-González, S. Sánchez Moral, R. Sancho, and E. Sanz Rubio. 2000. El registro sedimentario y faunístico de las cuencas de Calatayud-Daroca y Teruel: evolución paleoambiental y paleoclimática durante el Neógeno. *Revista de la Sociedad Geológica de España* 13:323–343.

- Álvarez-Sierra, M. A. 1987. Estudio sistemático y bioestratigráfico de los Eomyidae (Rodentia) del Oligoceno superior y Mioceno inferior español. *Scripta Geologica* 86:1–207.
- Álvarez-Sierra, M. A., J. P. Calvo, J. Morales, A. M. Alonso-Zarza, B. Azanza, I. García Paredes, M. Hernández Fernández, A. J. Van der Meulen, P. Pélaez-Campomanes, V. Quiralte, M. J. Salesa, I. M. Sánchez, and D. Soria. 2003. El tránsito Aragoniense-Vallesiense en el área de Daroca-Nombrevilla (Zaragoza, España). *Coloquios de Paleontología Volumen Extraordinario* 1:25–33.
- Argyropulo, A. I. 1938. On the fauna of Tertiary Cricetidae of the USSR. *Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS* 20:223–226.
- Azanza, B., A. M. Alonso-Zarza, M. A. Álvarez-Sierra, J. P. Calvo, S. Fraile, I. García Paredes, E. Gómez, M. Hernández Fernández, A. J. Van der Meulen, D. De Miguel, P. Montoya, J. Morales, X. Murelaga, P. Pélaez-Campomanes, B. Pérez, B., Quiralte, M. J. Salesa, A. Sánchez Marco, and D. Soria. 2004. Los yacimientos de vertebrados continentales del Aragoniense superior (Mioceno medio) de Toril, Cuenca de Calatayud-Daroca. *Geotemas* 6:271–274.
- Baudelot, S. 1972. Etude des Chiroptères, Insectivores et Rongeurs du Miocène de Sansan (Gers). 364 pp. Ph.D. dissertation, Université Paul Sabatier, Toulouse, France.
- Casanovas-Vilar, I., and J. Agustí. 2007. Ecogeographical stability and climate forcing in the Late Miocene (Vallesian) rodent record of Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 248:169–189.
- Daams, R., and M. Freudenthal. 1981. Aragonian: the stage concept versus Neogene mammal zones. *Scripta Geologica* 62:1–17.
- Daams, R., and M. Freudenthal. 1988. Synopsis of the Dutch-Spanish collaboration program in the Neogene of the Calatayud-Teruel basin. *Scripta Geologica Special Issue* 1:3–18.
- Daams, R., and A. J. Van der Meulen. 1984. Paleoenvironmental and paleoclimatic interpretation of micromammal faunal successions in the upper Oligocene and Miocene of north central Spain; pp. 241–257 in Meulenkamp, J. (ed.), *Paleoenvironnements continentaux en Méditerranée au Néogène et évolution paléoclimatique*. *Paleobiologie Continentale*, Montpellier.
- Daams, R., A. J. Van der Meulen, P. Peláez-Campomanes, M. A. Álvarez-Sierra. 1999a. Trends in rodent assemblages from the Aragonian (Early Middle Miocene) of the Calatayud-Daroca Basin (Aragón, Spain); pp. 127–139 in J. Agustí, L. Rook, P. Andrews (eds.), *Hominoid evolution and climate change in Europe. 1 The evolution of Neogene terres-*

- trial ecosystems in Europe. Cambridge University Press, Cambridge.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, and W. Krijgsman. 1999b. Aragonian stratigraphy reconsidered and a re-evaluation of the middle Miocene mammal biochronology in Europe. *Earth and Planetary Science Letters* 165:287–294.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999c. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.
- Daams, R., L. Alcalá, M. A. Álvarez-Sierra, B. Azanza, J. A. Van Dam, A. J. Van der Meulen, J. Morales, M. Nieto, P. Peláez-Campomanes, and D. Soria. 1998. A stratigraphical framework for Miocene (MN4–MN13) continental sediments of Central Spain. *Comptes rendus de l'Académie des sciences, Séries IIA* 327:625–631.
- De Bruijn, H. 1976. Vallesian and Turolian rodents from Biotia, Attica and Rhodes (Greece). *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen* 79:361–384.
- De Bruijn, H., and E. Ünay. 1996. On the evolutionary history of the Cricetodontini from Europe and Asia Minor and its bearing on the reconstruction of migrations and the continental biotope during the Neogene; pp. 227–234 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Saraç, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a Discussion of the Evolutionary History of the Cricetodontini. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 96:151–216.
- De Bruijn, H., R. Daams, V. Fahlbusch, G. Daxner-Höck, L. Ginsburg, P. Mein, and J. Morales. 1992. Report of the RCMNS working group on fossil mammals, Reimsburg 1990. *Newsletters of Stratigraphy* 26:65–118.
- DeMiguel, D., B. Azanza, and J. Morales. 2011. Paleoenvironments and paleoclimate of the Middle Miocene of central Spain: A reconstruction from dental wear of ruminants. *Palaeogeography, Palaeoclimatology, Palaeoecology* 302:452–463.
- Fischer, G. 1817. *Adversaria zoologica*, Mémoires de la Société Impériale des Naturalistes de Moscou 5:357–472.
- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mitteleuropas und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.

- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B 69:298–317.
- Garcés, M., W. Krijgsman, P. Peláez-Campomanes, M. A. Álvarez-Sierra, and R. Daams. 2003. *Hipparion* dispersal in Europe: magnetostratigraphic constraints from the Daroca area (Spain). *Coloquios de Paleontología Volumen Extraordinario* 1:171–178.
- García-Paredes, I. 2006. Patrones evolutivos de los Gliridae (Rodentia, Mammalia) del Mioceno inferior y medio del área tipo del Aragoniense (Cuenca de Calatayud–Montalbán). Ph.D. dissertation, Universidad Complutense de Madrid, Madrid, 676 pp.
- García-Paredes, I., P. Peláez-Campomanes, and M. A. Álvarez-Sierra. 2009. Gliridae (Rodentia, Mammalia) with a simple dental pattern: a new genus and new species from the European Early and Middle Miocene. *Zoological Journal of the Linnean Society* 157:622–652.
- García-Paredes, I., P. Peláez-Campomanes, and M. A. Álvarez-Sierra. 2010. *Microdyromys remerti*, sp. nov., a new Gliridae (Rodentia, Mammalia) from the Aragonian Type Area (Miocene, Calatayud-Montalbán Basin, Spain). *Journal of Vertebrate Paleontology* 30:1594–1609.
- Hartenberger, J. L. 1965. Les Cricetidae (Rod.) de Can Llobateres (Néogène d'Espagne). *Bulletin de la Société Géologique de France* 7:487–498.
- Hernández Fernández, M., J. A. Cárdena, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Higgen, F. J., L. J. Lourens, and J. Van Dam. 2012. The Neogene Period; pp. 923–978 in Gradstein, F. M., Ogg, J. G. and Schmitz, J. G. (eds.), *Geologic Time Scale 2012*. Cambridge University Press, Cambridge.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőtárkány-Felnémet (Northern Hungary). *Fragmenta Paleontologica Hungarica* 24–25:16–24.
- Hopkins, M. J., and S. Lidgard. 2012. Evolutionary mode routinely varies among morphological traits within fossil species lineages. *Proceedings of the National Academy of Sciences of the United States of America* 109:20520–20525.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für*

- Geologie und Paläntologie, Abhandlungen 254:177–235.
- Kordos, L. 1986. Upper miocene hamsters (Cricetidae, Mammalia) of Hasznos and Szentendre. Magyar Allami Földtani Intézet Jelentése Az 1984:523–553.
- Klein Hofmeijer, G., and H. de Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece) Part 8: The Cricetidae. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B 91:185–204.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d’une récapitulation des diverses espèces d’animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d’autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. J. A. Portes, Auch, 41 pp.
- López-Antoñanzas, R., and P. Mein. 2009. First detailed description of *Hispanomys bijugatus* Mein & Freudenthal, 1971 (Rodentia, Cricetodontinae) from the Upper Aragonian of La Grive-Saint Alban (France): Biostratigraphical implications. Geobios 42:783–796.
- López-Antoñanzas, R., and P. Mein. 2011. First detailed description of *Hispanomys decedens* (Rodentia) from the Middle Miocene of La Grive-Saint Alban (France). Swiss Journal of Geosciences 104:345–353.
- López-Antoñanzas, R., P. Peláez-Campomanes., M. A. Álvarez-Sierra, I. García Paredes. 2010. New species of *Hispanomys* (Rodentia, Cricetodontinae) from the Upper Miocene of Batallones (Madrid, Spain). Zoological Journal of the Linnean Society 160:725–747.
- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). Journal of Vertebrate Paleontology 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, R. López-Antoñanzas, A. Oliver Pérez, and P. Peláez-Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). Palaeontologica Nova 8:259–271.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra, R. López-Antoñanzas, and A. Oliver. 2009. Cricetodontini (Rodentia, Mammalia) from the upper Aragonian and lower Vallesian of the Toril-Nombrevilla section (Middle and Upper Miocene, Calatayud-Daroca basin, Zaragoza, Spain). Journal of Vertebrate Paleontology 29(3, supplement):161A.
- López-Guerrero, P., A. Oliver Pérez, M. A. Álvarez-Sierra, I. García Paredes, and P. Peláez-Campomanes. 2007. El registro de los castoridae (Rodentia, Mammalia) del Aragoniense y Vallesiense (Mioceno Medio y Superior) de las cuencas centrales españolas; pp. 118–119

- in J.C. Braga, A. Checa, and M. Company (eds.), XXIII Jornadas de la Sociedad Española de Paleontología (Caravaca de la Cruz 3–6 de Octubre de 2007). Libro de resúmenes. Instituto Geológico y Minero de España & Universidad de Granada, Granada.
- López-Guerrero, P., I., García-Paredes, L. W., Van den Hoek Ostende, J., Van Dam, M. A., Álvarez-Sierra, V., Hernández-Ballarín, A. J., Van der Meulen, A., Oliver, A., and P. Peláez-Campomanes. 2011. Cañada: a new micromammal succession from the lower Vallesian and Turolian of the Daroca area (Calatayud-Montalbán basin, Spain). *Estudios Geológicos* 67:443–453.
- López-Martínez, N. 1997. Lignées évolutives, formes intermédiaires et évolution en mosaïque: exemple des lagomorphes européens; pp. 105–120 in J. P., Aguilar, S. Legendre, and J. Michaux (eds.), *Congrès BiochroM '97. Mémoires et Travaux EPHE*, Institut Montpellier, Montpellier.
- López-Martínez, N., C., Sesé Benito, and J. L. Sánchez-García. 1977. La microfauna (Rod., Insect., Lagomorph y Reptilia) de las fisuras del Mioceno medio de Escobosa de Calatañazor (Soria, España). *Acta Geologica Hispanica* 12:47–73.
- Maridet, O., and S. Sen. 2012. Les Cricetidae du gisement de Sansan; pp. 29–64 in S. Peigné and S. Sen (eds.), *Les mammifères du gisement de Sansan. Mémoires du Muséum national d'Histoire naturelle*, Paris.
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Oliver, A., and P. Peláez-Campomanes. 2013. *Megacricetodon vandermeuleni*, sp. nov. (Rodentia, Mammalia), from the Spanish Miocene: a new evolutionary framework for *Megacricetodon*. *Journal of Vertebrate Paleontology* 33:943–955.
- Oliver, A., P. López-Guerrero, I. García-Paredes, M. A. Álvarez-Sierra, and P. Peláez-Campomanes. 2009. Evolution of *Megacricetodon* tooth pattern through geometric morphometric analysis. *Journal of Vertebrate Paleontology* 29(3, supplement):158A.
- Peláez-Campomanes, P. 2001. Revision of the Aragonian (Miocene) *Atlantoxerus* (Sciuridae). *Journal of Paleontology* 75:418–426.
- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.

- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata Palasiatica* 48:27–47.
- Rummel, M. 1995. *Cricetodon bolligeri* n. sp. ein neuer Cricetide aus dem Obermiozän von Petersbuch bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 35:109–123.
- Rummel, M. 1998. Die Cricetiden aus dem Mittel- und Obermiozän der Türkei. *Documenta naturae* 123:1–300.
- Rummel, M. 2000. Miocene Cricetodontini from Petersbuch near Eichstaett. The genus *Cricetodon* Lartet 1851. *Senckenbergiana Lethaea* 80:149–171.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde B* 311:1–6.
- Rummel, M., and D. Kälin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Mittelmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Sánchez I. M., and J. Morales. 2006. Distribución biocronológica de los Moschidae (Mammalia, Ruminantia) en España. *Estudios Geológicos* 62:533–543.
- Sánchez I. M., and J. Morales. 2008. *Micromeryx azanzae* sp. nov. (Ruminantia: Moschidae) from the middle-upper Miocene of Spain, and the first description of the cranium of *Micromeryx*. *Journal of Vertebrate Paleontology* 28:873–885.
- Schaub S. 1925. Die hamsterartigen Nagetiere des Tertiärs und ihre lebenden Verwandten. *Abhandlungen der Schweizerischen paläontologischen Gesellschaft* 45:1–114.
- Sen, S., and M. A. Erbajeva. 2011. A new species of *Gobicricetodon* Qiu, 1996 (Mammalia, Rodentia, Cricetidae) from the Middle Miocene of Aya Cave, Lake Baikal. *Vertebrata Palasiatica*, 49:257–274.
- Sen, S., and E. Ünay. 1979. Sur quelques Cricetodontini (Rod.) du Miocène moyen d'Anatolie. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* 82:293–301.
- Sesé Benito, C. 1977. Los cricétidos (Rodentia, Mammalia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno Cuaternario* 8:127–80.
- Simpson, G. C. 1945. The principles of classification and a classification of mammals. *Bulletin of the American Museum of Natural History* 85:1–350.
- Stebbins G. L. 1983. Mosaic evolution: An integrating principle for the modern synthesis. *Experientia* 39:823–834.

- Stehlin, H. G., and S. Schaub. 1951. Die Trigonodontie der simplicidentaten Nager. Schweizerische paläontologische Abhandlungen 67:1–385.
- Steininger, F. 1999. Chronostratigraphy, Geochronology and Biochronology of the Miocene ‘European Land Mammals Mega-Zones’ (ELMMZ) and the Miocene ‘Mammal-Zones (MN-Zones)’; pp. 9–24 in G. Rößner and K. Heißig (eds.), *The Miocene Land Mammals of Europe*. Verlag Dr. Friedrich Pfeil, München.
- Tobien, H. 1978. New species of Cricetodontini (Rod. Mam.) from the Miocene of Turkey. *Mainzer Geowissenschaftliche Mitteilungen Geologisches Landesamt Rheinland-Pfalz* 6:209–219.
- Ünay, E. 1980. The Cricetodontini (Rod. Mam.) from the Bayraktepe section (Çanakkale, Turkey). *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen, Series B* 83:399–418.
- Ünay, E., and H. De Bruijn. 1984. On some Neogene Rodent assemblages from both sides of the Dardanelles, Turkey. *Newsletter of Stratigraphy* 13:119–132.
- Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. Van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodent faunas of the Mio–Pliocene continental sediments of the Teruel-Alfambra region, Spain. *Utrecht Micropaleontological Bulletin, Special Publication* 2:1–217.
- Van der Meulen, A. J., and R. Daams. 1992. Evolution of Early–Middle Miocene rodent faunas in relation to long-term palaeoenvironment changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 93:227–253.
- Van der Meulen, A. J., P. Peláez-Campomanes, and R. Daams. 2003. Revision of medium-sized Cricetidae from the Miocene of the Daroca-Villafeliche area in the Calatayud-Teruel basin (Zaragoza, Spain). *Coloquios de Paleontología Volumen extraordinario* 1:385–441.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, and P. Peláez-Campomanes. 2012. Updated Aragonian biostratigraphy: Small Mammal distribution and its implications for the Miocene European Chronology. *Geologica Acta* 10:1–24.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, P. López-Guerrero, V. Hernández-Ballarín, and P. Peláez-Campomanes. 2011. Biostratigraphy or Biochronology? Lessons from the Early and Middle

Miocene small Mammal events in Europe. *Geobios* 44:309–321.

Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.

Chapter 8

Hispanomys



SPECIES OF *HISPANOMYS* FROM THE LATE ARAGONIAN AND EARLY VALLESIAN (MIDDLE-LATE MIOCENE) OF THE CALATAYUD-DAROCA BASIN, ZARAGOZA, SPAIN

“The prolonged difference (MN 7/8-12) between an Asiatic/Eastern European group of species, type 4, classified in Byzantinia and a Southwest European group of species, type 5, classified in Hispanomys and Ruscinomys, suggests that there must have been an effective barrier between these two areas during the late Miocene.”

Hans de Bruijn & Engin Ünay

Resumen

8

En este capítulo se estudia la presencia de uno de los géneros de Cricetodontini en la Cuenca de Calatayud-Daroca. El género *Hispanomys* se ha encontrado en muchas localidades de la Península Ibérica, Francia, Europa central y occidental; su distribución estratigráfica está restringida entre el final del Mioceno medio y el comienzo del Mioceno superior. Las especies que se encuentran en la Cuenca de Calatayud-Daroca son *Hispanomys aguirrei* (biozona local G3, Aragoniense superior), *H. lavocati* (biozona local G3, Aragoniense superior), *H. nombrevillae* (biozona local H, Aragoniense superior-Vallesiense inferior), e *H. aragonensis* (biozona local I, Vallesiense inferior). Morfológicamente, estas cuatro especies se pueden dividir en dos grupos: *Hispanomys aguirrei* e *H. lavocati* presentan características típicas de las especies basales de *Hispanomys*. El otro grupo está formado por *H. nombrevillae* e *H. aragonensis*. Estas últimas especies son morfológicamente más homogéneas que las otras dos y debido a que todos sus caracteres son típicos de *Hispanomys*, son más fáciles de asignar a nivel genérico. El estadio evolutivo de estas especies es fácil de reconocer; *Hispanomys nombrevillae* tiene caracteres descritos como basales dentro de la tribu e *H. aragonensis* los tiene derivados. Los dos grupos de especies se originaron en el suroeste de Europa pero no parecen estar filogenéticamente relacionados entre sí. Se observa la tendencia propuesta previamente en la que se reduce el tamaño de la especies con el tiempo; *Hispanomys aguirrei* e *H. lavocati* son más antiguas y más grandes que *H. nombrevillae* e *H. aragonensis*.



Abstract

The presence of a genus of Cricetodontini in the Calatayud-Daroca Basin is reported. *Hispanomys* has been recovered from many localities in the Iberian Peninsula, France, central and western Europe; its stratigraphic distribution is restricted to the middle and the beginning of the late Miocene. The species found in the Calatayud-Daroca Basin are *Hispanomys aguirrei* (local biozone G3, late Aragonian), *H. lavocati* (local biozone G3, late Aragonian), *H. nombrevillae* (local biozone H, late Aragonian-early Vallesian) and *H. aragonensis* (local biozone I, early Vallesian). Morphologically, these four species can be split in two groups: *Hispanomys aguirrei* and *H. lavocati* display characteristics typical of a basal *Hispanomys* and their generic assignation have required a detailed study of the material. The other group is formed by *H. nombrevillae* and *H. aragonensis*. They present more homogeneous morphology than the above-mentioned species and their taxonomical assignation is clearer because all their characters are typical of *Hispanomys*. The evolutionary stage of these species is easy to recognize; *Hispanomys nombrevillae* have characters described as basal within the tribe and *H. aragonensis* is more derived. Both groups of species were originated in southwestern Europe but they do not seem to be phylogenetically related. The trend previously stated of size decreasing in time is observed; *Hispanomys aguirrei* and *H. lavocati* are older and larger than *H. nombrevillae* and *H. aragonensis*.

INTRODUCTION

The genus *Hispanomys* Mein and Freudenthal, 1971a is a large-sized rodent of the family Cricetidae. It is characterized by its high hypsodonty; its well-developed ectolophs and its poorly-developed structures closing the valleys. *Hispanomys* contains 16 fossil species; four of them are present in the Calatayud-Daroca Basin (Freudenthal, 1966; Mein and Freudenthal, 1971a; Álvarez-Sierra et al., 2003; López-Guerrero et al., 2008, 2009) (Table 8.1). The oldest occurrence of the genus in Europe is quite controversial. *Hispanomys castelnovi* Aguilar et al., 1994 is described in the fissure filling of Castelnou 6. These authors placed the locality at the top of the MN6 (late Aragonian) and correlated it with the local biozone G2 from Calatayud-Daroca (Aguilar et al., 1994). However, the material was recovered from a karstic fissure filling and the association described by Aguilar et al., (1994) could include fossils from a wide time range, so this date has to be carefully taken. *Hispanomys decedens* (Schaub, 1925) and *H. bijugatus* Mein and Freudenthal, 1971a were recovered from La Grive M and L3 respectively. This locality is also a fissure filling. The relative position of La Grive M and La Grive L3 is dubious. López-Antoñanzas and Mein, (2011) considered La Grive M the oldest fissure filling of La Grive-Saint Alban and is dated around 13 Ma (Maridet, 2002). However, Maridet (2003), considered La Grive L3 older than La Grive M based in the evolutionary stages of several rodent genera. Moreover, La Grive M can be biostratigraphically correlated with the middle part of the local biozone G3 (upper Aragonian) from the Calatayud-Daroca Basin (Peláez-Campomanes, pers., comm.). In any case, the oldest occurrence of the genera seems to be in La Grive-Saint Alban sites. The youngest occurrence is *H. adroveri* Agustí, 1986 from the locality JUN-2B (Canteras del Jun section, Granada Basin, Spain) which is placed in the middle Turolian (MN12; García-Alix et al., 2008).

As explained in chapter 5, Mein and Freudenthal (1971a), described new genera, subgenera and species when emended the original diagnosis of *Cricetodon*. One of the subgenera proposed was *Hispanomys*. This first diagnosis of *Hispanomys* was very general and partially referred to the mandible and incisors, elements that usually are absent in the fossil assemblages. Later on, Van de Weerd (1976) gave generic rank to *Hispanomys* and *Ruscinomys*. He synonymized some taxa and emended the diagnosis of *Hispanomys*. Since then, the generic status of *Hispanomys* and *Ruscinomys* were no longer discussed and is widely accepted. Focusing on the Calatayud-Daroca Basin, Freudenthal (1966) described *H. nombrevillae* and *H. aragonensis* in the lower Vallesian of Nombrevilla and Daroca areas. Álvarez-Sierra et al. (2003) presented the new localities of the

Toril-Nombrevilla section and they reported the presence of *Cricetodon* in the Aragonian and *Hispanomys* in the Vallesian, but they did not determinate which species were found. Therefore, the presence of *Hispanomys* in the late Aragonian from Nombrevilla section was not noticed until the revision of the fossil site of Nombrevilla 2 by López-Guerrero et al. in 2008.

The species of the Calatayud-Daroca Basin provide very valuable information about the origins of *Hispanomys* and its relationship with *Cricetodon*.

Species	Locality	Age	Original reference	Reference for the age
<i>H. adroveri</i>	Casa del Acero	MN12	Agustí, 1986	Agustí, 1986
<i>H. freudenthali</i>	Masada del Valle 2	MN12	Van de Weerd, 1976	Van de Weerd, 1976
<i>H. baixasi</i>	Lo Fournas 16-M	MN11	Aguilar et al., 2007	Aguilar et al., 2007
<i>H. peralensis</i>	Perales C	MN10	Van de Weerd, 1976	Van de Weerd, 1976
<i>H. moralesi</i>	Bataillon 10	MN10	López-Antoñanzas et al., 2010	López-Antoñanzas et al., 2010
<i>H. mediterraneus</i>	Montredon	MN10	Aguilar 1982	Aguilar, 1982
<i>H. thaleri</i>	Can Llobateres	MN9	Hartenberger, 1965	Hartenberger, 1965
<i>H. aragonensis</i>	Pedregueras 2C	MN9	Freudenthal, 1966	Daams et al., 1999
<i>H. nombrevillae</i>	Nombrevilla 1	MN9	Freudenthal, 1966	Daams et al., 1999
<i>H. dispectus</i>	Hostalets de Pierola	MN7/8-MN9	Agustí, 1980	Agustí, 1980
<i>H. daamsi</i>	Can Missert	MN9	Agustí et al., 2005	Gómez Cano et al., 2011
<i>H. lavocati</i>	Hostalets de Pierola	MN7/8	Freudenthal, 1966	López-Antoñanzas and Mein, 2009
<i>H. aguirrei</i>	Escobosa de Calatañazor	MN7/8	Sesé in López-Martínez et al., 1977	Sesé Benito, 1977
<i>H. bijugatus</i>	La Grive-Saint Alban, L3	MN7/8	Mein and Freudenthal, 1971a	Mein and Freudenthal, 1971a
<i>H. decedens</i>	La Grive-Saint Alban, L5'	MN7/MN8	Schaub, 1925	López-Antoñanzas and Mein, 2009
<i>H. castelnovi</i>	Castelnou 6	MN6?	Aguilar et al., 1994	Aguilar et al., 1994

TABLE 8.1. List of the described species of *Hispanomys*, type locality and their ages. The references for the age (MN zones) of the type localities are given in the fourth column.

MATERIAL & METHODS

The studied material includes 2,359 upper and lower molars of *Hispanomys* collected at 19 fossil sites from the Calatayud-Daroca Basin (Fig. 8.1). The fossils are housed at the Museo Nacional de Ciencias Naturales-CSIC (Madrid, Spain), the Naturalis Biodiversity Center (Leiden, The Netherlands), and the Instituut voor Aardwetenschappen, Utrecht Universiteit (The Netherlands).

Table 8.2 lists the studied localities, including the number of examined molars, the age of the localities, and their abbreviations. In order to complete the available information of the different species of *Hispanomys*, the type material of several species was studied personally (Table 8.3). The nomenclature used for the check teeth is the one described by Mein and Freudenthal (1971b) and modified by López-Guerrero et al. (2013). Length and width measurements for each specimen, given in mm, were taken using a Nikon Measuroscope 10 microscope with digital micrometer. Maximum length and width were measured following the procedure described in section II. (Material y Métodos). Descriptive statistics of metrical variables from all the studied localities and from the type localities of *H. aguirrei* (Escobosa de Calatañazor) and *H. lavocati* (Hostalets de Pierola) are given in tables 8.4 to 8.7. We built length/width scatter-diagrams for each element (Fig. 8.2-8.5) including also Cricetodontini with size similar to the studied species. Different statistical analyses were carried out in those localities that contain at least five data per dental element. These are the following: ANOVA test among the localities with *H. aguirrei*, including the data for the type locality, Escobosa de Calatañazor; Student's *t* test for *H. nombrevillae* as well as for *H. aragonensis*, and finally, another Student's *t* to compare the size of *H. nombrevillae* and *H. aragonensis*. The results of the different analyses are given in tables 8.8-8.10. The photographs were taken with an environmental Scanning Electron




FIGURE 8.1. Geological map of the Calatayud-Daroca Basin. The square marks the situation of Daroca area (Modified after García-Paredes et al., 2010).

Microscope FEI Quanta 200 at the Museo Nacional de Ciencias Naturales-CSIC (Madrid, Spain). The image processing was made using Adobe Photoshop software.

	Species	Locality	Abb.	Collections	M1	M2	M3	m1	m2	m3	N
<i>Hispanomys aragonensis</i> / <i>Hispanomys</i> cf. <i>aragonensis</i>											
9.98	<i>Hispanomys</i> cf. <i>aragonensis</i>	Nombrevilla 19	NOM19	MNCN	—	—	—	—	—	1	1
10.38	<i>Hispanomys aragonensis</i>	Pedregueras 2C	PED2C	RGM	19	30	40	49	30	39	207
10.62	<i>Hispanomys aragonensis</i>	Pedregueras 2A	PED2A	IAUU	67	75	53	71	74	31	371
	<i>Hispanomys aragonensis</i>	Pedregueras 1A	PED1A	IAUU	1	4	2	3	—	2	12
10.68	<i>Hispanomys</i> cf. <i>aragonensis</i>	Nombrevilla 14	NOM14	MNCN	2	1	2	1	—	1	7
10.72	<i>Hispanomys</i> cf. <i>aragonensis</i>	Nombrevilla 13	NOM13	MNCN	1	—	—	1	1	—	3
	<i>Hispanomys</i> cf. <i>aragonensis</i>	Nombrevilla 22	NOM22	MNCN	—	2	2	1	—	—	5
				Total	90	112	99	126	105	73	605
<i>Hispanomys nombrevillae</i> / <i>Hispanomys</i> cf. <i>nombrevillae</i>											
	<i>Hispanomys</i> cf. <i>nombrevillae</i>	Cañada 11	CAÑ9	MNCN	—	—	1	—	—	—	1
	<i>Hispanomys nombrevillae</i>	Cañada 10	CAÑ10	MNCN	—	2	1	2	—	—	5
	<i>Hispanomys</i> cf. <i>nombrevillae</i>	Cañada 9	CAÑ9	MNCN	—	2	—	1	1	—	4
	<i>Hispanomys</i> cf. <i>nombrevillae</i>	Cañada 8	CAÑ8	MNCN	—	2	—	—	2	—	4
10.77	<i>Hispanomys nombrevillae</i>	Nombrevilla 1	NOM1	IAUU/MNCN	19	19	15	25	23	23	124
11.18	<i>Hispanomys</i> cf. <i>nombrevillae</i>	Nombrevilla 10	NOM10	MNCN	2	3	5	2	2	2	16
11.21	<i>Hispanomys nombrevillae</i>	Nombrevilla 9	NOM9	MNCN	5	5	6	8	9	13	46
				Total	26	33	28	38	37	38	200
<i>Hispanomys aguirrei</i> / <i>Hispanomys</i> cf. <i>aguirrei</i>											
11.33	<i>Hispanomys aguirrei</i>	Carrilanga 1	CARR1	RGM	25	27	27	27	31	24	161
11.80	<i>Hispanomys</i> cf. <i>aguirrei</i>	Nombrevilla 4	NOM4	MNCN	1	1	—	—	—	1	3
11.84	<i>Hispanomys</i> cf. <i>aguirrei</i>	Nombrevilla 3	NOM3	MNCN	3	3	—	—	2	3	11
11.87	<i>Hispanomys aguirrei</i>	Nombrevilla 2	NOM2	MNCN	79	95	93	91	75	115	548
12.01	<i>Hispanomys aguirrei</i>	Solera	SOL	RGM	61	64	38	60	65	64	352
12.39	<i>Hispanomys</i> cf. <i>aguirrei</i>	Paje 2	PJE2	MNCN	6	1	4	6	5	4	26
12.26	<i>Hispanomys</i> cf. <i>aguirrei</i>	Paje 1	PJE1	MNCN	1	2	—	4	1	—	8
				Total	176	193	162	188	179	211	1109
<i>Hispanomys lavocati</i> / <i>Hispanomys</i> cf. <i>lavocati</i>											
11.80	<i>Hispanomys</i> cf. <i>lavocati</i>	Nombrevilla 4	NOM4	MNCN	3	2	—	2	2	—	9
11.84	<i>Hispanomys</i> cf. <i>lavocati</i>	Nombrevilla 3	NOM3	MNCN	1	—	2	—	—	1	4
11.87	<i>Hispanomys lavocati</i>	Nombrevilla 2	NOM2	MNCN	76	71	85	57	57	86	432
				Total	80	73	87	59	59	87	445

TABLE 8.2. Material of *Hispanomys* from the Calatayud-Daroca Basin, studied localities, abbreviations for the localities and institutions where the fossils are stored (see institutional abbreviations). The localities are dated following Van Dam et al. (2006) and Van Dam et al. (in press.). Abbreviations: Abb, abbreviations for the localities; N, number of specimens.

Institutional Abbreviations—**BSPG**, Bayerische Staatssammlung für Paläontologie und Geologie, Munich (Germany); **IAUU**, Instituut voor Aardwetenschappen, Utrecht University (The Netherlands); **MNCN**, Museo Nacional de Ciencias Naturales-CSIC, Madrid (Spain); **NHNB**, Muséum d'Histoire Naturelle de Bâle (Switzerland); **PM**, Páztó Múzeum, Páztó (Hungary); **RGM**, Naturalis Biodiversity Center, former Rijksmuseum van Geologie en Mineralogie, Leiden (The Netherlands); **UCM**, Facultad de Geología, Universidad Complutense de Madrid (Spain).

Species	Locality	Age	Institution
<i>H. aragonensis</i>	Pedregueras 2C	MN9, biozone I	IAUU
<i>H. nombrevillae</i>	Nombrevilla 1	MN9, biozone H	IAUU
<i>'C'. fandli</i>	Gratkorn	MN9	BSPG
<i>C. bolligeri</i>	Petersbuch 10	MN7/8	BSPG
<i>H. lavocati</i>	Hostalets de Pierola	MN7/8	ICP
<i>H. aguirrei</i>	Escobosa de Calatañazor	MN7/8	MNCN
<i>'C'. klariankae</i>	Felsőtárkány-Felnémet	MN7/8	PM
<i>H. bijugatus</i>	La Grive-Saint Alban, L3	MN7/8	RGM
<i>H. decedens</i>	La Grive-Saint Alban, L5'	MN7/8	RGM
<i>C. albanensis</i>	La Grive-Saint Alban	MN7/8	RGM
<i>C. sansaniensis</i>	Sansan	MN6	MHNB
<i>C. jotae</i>	Manchones	MN6, biozone G2	IAUU
<i>C. soriae</i>	SomosaguasN	MN5, biozone E	UCM
<i>C. aureus</i>	Vieux-Collonges	MN5	RGM
<i>C. meini</i>	Vieux-Collonges	MN5	RGM

TABLE 8.3. List of species personally studied for the comparison of *Hispanomys* from the Calatayud-Daroca Basin. They belong either to *Hispanomys* or *Cricetodon*. Species, type locality and institutions where the fossils are stored (see institutional abbreviations).

SYSTEMATIC PALEONTOLOGY

Order RODENTIA Bowdich, 1821

Order RODENTIA Bowdich, 1821

Family CRICETIDAE Fischer, 1817

Subfamily CRICETODONTINAE Stehlin and Schaub, 1951

Tribe CRICETODONTINI Simpson, 1945

Genus *HISPANOMYS* Mein and Freudenthal, 1971a

Diagnosis

Original Diagnosis (Mein and Freudenthal, 1971a:19, translated from French)—“[Medium-sized Cricetodontini, the upper incisors are provided with a weak longitudinal groove, and the upper molars have partial ectolophes; the M1 has a divided anterocone; four roots and often a trace of a fifth one which originates at the base of the large anterior root; the molars are slightly higher than in the subgenus *Cricetodon* but the hypsodonty value is never as high as it can be in *Ruscinomys* and *Pseudoruscinomys*. The size is smaller than that of *Deperetomys* and *Pseudoruscinomys*.]”

Emended Diagnosis (Van de Weerd, 1976: 105)—“Cricetodontinae of medium size, with either incomplete or complete ectolophs. The M1 has either four or five roots. The anterocone of the M1 has an anterior groove. The anterolophid of the lower molars may be absent. The molars are slightly to moderately hypsodont. The depth of the groove in the upper incisor is variable”.

Differential Diagnosis (Van de Weerd, 1976: 105)—“Most species of *Hispanomys* are larger than those of *Cricetodon*, the ectolophs of *Hispanomys* are better developed than those of *Cricetodon* and the M3 of *Hispanomys* are relatively smaller. *Hispanomys* has smaller and less hypsodont molars than *Ruscinomys*. The M3 of *Hispanomys* is less reduced than that of *Ruscinomys*. In contrast to *Hispanomys*, the teeth of *Ruscinomys* have always five roots and never show an anterolophid.”

Type species—*Hispanomys aragonensis* (Freudenthal, 1966)

Other species included—See table 8.1.

HISPANOMYS AGUIRREI (Sesé Benito, 1977 in López Martínez et al., 1977)

(Figs. 8.2; 8.6-8.8; Tables 8.2, 8.4; Appendix 8.1)

Material

Original Type Locality—Escobosa de Calatañazor, Duero Basin, Spain (MN7/8, upper Aragonian, middle Miocene).

Stratigraphic Range in the Calatayud-Daroca Basin—Local biozone G3 (MN7/8, upper Aragonian, middle Miocene).

Geographical Range—Southwestern Europe.

Measurements—See table 8.4 and figure 8.2.

Diagnosis

Original Diagnosis (López Martínez et al., 1977:56, translated from Spanish)—“[Medium-sized *Cricetodon* [sic]. The upper and lower M3 are more reduced than in *C. jotae* from Manchones and less reduced than in *C. decedens nombrevillae* [sic] from Nombrevilla. The metaconid is joined to the anteroconid through an anterior metalophulid on the m1. The m2 develops three roots in most of the molars. The presence of well-developed ectolophs is very common in the upper cheek teeth. The posterior ectoloph is always present on the M1 and it is connected with the metacone in the 70% of the teeth, and the anterior ectoloph is present in the 90%. It is connected with the paracone in the 50%. The posterior ectoloph is always present in the M2 and it is complete, approximately, in the half of the sample. Both mesoloph and mesolophid are reduced or absent. The upper incisors have a slight groove.]”

HISPANOMYS AGUIRREI FROM THE CALATAYUD-DAROCA BASIN

Description of the Material from Nombrevilla 2

M1—This molar has four roots. The anterocone is large and divided in two parts by a shallow groove on the anterior wall of the tooth. This groove does not reach the base of the anterocone and this cusp looks like single in worn teeth. A labial anteroloph is found in 20 out of 53 cases, extending through the protosinus. The anterolophule connects the protocone and the lingual part of the anterocone (Fig. 8.6A; 8.6D). A posterior protolophule connects the paracone with the anterior branch of the hypocone (Fig. 8.6A; 8.6D). The anterior protolophule and the labial spine of the anterolophule are absent (Fig. 8.6A; 8.6D). Almost all teeth have anterior ectoloph (62/65); it is composed by the posterior ectoloph of the anterocone (52/62; Fig. 8.6A) or it is double (10/62; Fig. 8.6D), composed also by the anterior ectoloph of the paracone. The anterior ectoloph forms a continuous connection between the anterocone and the paracone (is complete) in 30 out of 63 specimens, although this connection is lower than the labial cusps in eight cases (Fig. 8.6D). The posterior ectoloph is always present; it is composed by the posterior ectoloph of the paracone (43/61; Fig. 8.6A; 8.6D) or it is double, composed by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone (18/61). The posterior ectoloph can be complete (32/58) or not (26/58; Fig. 8.6A; 8.6D). Eighteen out of 63 cases have mesoloph; it is short (5/18), long, but not exceeding the half of the mesosinus (3/18), or it can be an enlargement of the enamel (incipient) (10/18; Fig. 8.6A; 8.6D). The mesoloph is connected to the ectoloph in three out of 72 specimens. The anterior wall of the protocone sometimes shows a spine (6/19; Fig. 8.6A). Three specimens possess entomesoloph, and it is disconnected to the longitudinal ridge. The metalophule is posterior and connected to the labial posteroloph (Fig. 8.6A; 8.6D). Both labial and lingual posterolophs are well developed (Fig. 8.6A; 8.6D). The anterosinus is closed by a low cingulum in 25 out of 57, by a cingulum (29/57; Fig. 8.6A; 8.6D), or it remains open. The mesosinus is always closed by a cingulum (Fig. 8.6A; 8.6D), which is low in 11 out of 57. The sinus is closed by a cingulum (8/62), by a style (12/62) or it remains open (42/62; Fig. 8.6A; 8.6D). The protosinus is closed by the lingual anteroloph (20/53), by a cingulum (17/53; Fig. 8.6A), by a style (14/53) or it remains open (2/53; Fig. 8.6D).

M2—The anterior ectoloph is present in 28 out of 78 cases; it is always composed by the posterior ectoloph of the anterocone. The anterior ectoloph is complete in 20 specimens, although in one of them this connection is lower than the labial cusps. All teeth have posterior ectoloph; it is simple, composed by the posterior ectoloph of the paracone (55/79; Fig. 8.6E), or by both anterior ectoloph of the metacone and posterior ectoloph of the paracone (24/79; Fig. 8.6E). The posterior ectoloph is complete in 17 out of 79 specimens. Only eight cases possess mesoloph, being incipient in four of them. One out of 80 molars displays a short entomesoloph. The anterior wall of the protocone develops a spine in 19/24 cases (Fig. 8.6B; 8.6E) and five teeth lack of it (Fig. 8.6A). Most of the M2 show labial posteroloph (55/57; Fig. 8.6B; 8.6E), whereas the

lingual posteroloph is less frequent (19/48; Fig. 8.6B; 8.6E). The protosinus is weak in 14 out of 26 specimens (Fig. 8.6B; 8.6E) and it is closed by a low cingulum in one of them. The mesosinus is closed in 39 out of 44, by a low cingulum in 21 (Fig. 8.6E), by a cingulum in 16 (Fig. 8.6B) and by two crests in the remaining two teeth. The sinus can be open (48/67; Fig. 8.6E), or closed by a cingulum (12/67; Fig. 8.6B), by two crests (2/67), or by a style (5/67).

M3—It is reduced and the hypocone is barely distinguishable (Fig. 8.6C; 8.6F). The lingual anteroloph is weak (6/75; Fig. 8.6C; 8.6F). All specimens have a well-developed labial anteroloph (Fig. 8.6C; 8.6F). The anterior ectoloph is always absent but one case (1/75), where it is simple, composed by the anterior ectoloph of the paracone, and complete. The posterior ectoloph is always present; it is simple, composed by the posterior ectoloph of the paracone, in 38 out of 70 (Fig. 8.6C; 8.6F); it is double, composed by both anterior ectoloph of the metacone and posterior ectoloph of the paracone, in 32 out of 70. In all specimens, the ectolophs start from the labial part of the cusps (Fig. 8.6C), but one, which starts from the protolophule (Fig. 8.6F). The posterior ectoloph can be complete (47/68) or not (21/68; Fig. 8.6C; 8.6F). The mesoloph, entomesoloph, enamel coated valley and neo-entoloph are absent (Fig. 8.6C; 8.6F). The axioloph is always continuous (Fig. 8.6C; 8.6F). Twenty out of 71 teeth possess posteroloph; it is well developed in eight of them (Fig. 8.6C) and incipient in 12 (Fig. 8.6F). The sinus is long, curved and directed forward in 28 out of 40 cases (Fig. 8.6C; 8.6F); it is transversal in 10 molars.

m1—The anteroconid is rounded and situated on the longitudinal axis of the occlusal surface. The labial anterolophid is present in all cases but one, and it can reach the protoconid (22/56) or not (33/56; Fig. 8.7A; 8.7D). The lingual anterolophid is absent. All specimens but three have metalophulid I (Fig. 8.7A; 8.7D), and 34 out of 56 possess also the metalophulid II (Fig. 8.7D). Six out of 86 teeth show mesolophid (Fig. 8.7A; 8.7D); it is incipient in three and short in the other three. The ectomesolophid is absent. The posterolophid is an oval cusp rather than a ridge, due to a constriction on the posterior wall (Fig. 8.7A; 8.7D). The mesosinusid is blocked by a low cingulid in 19 out of 78 specimens; it is open in the remaining ones (Fig. 8.7A; 8.7D). The sinusid is narrow, long and oblique (Fig. 8.7A; 8.7D); it is closed in almost all teeth (64/65), by a stylid in one, by a cingulid in 59 (Fig. 8.7A) and by two crests attached to the hypoconid and the protoconid in the remaining four specimens (Fig. 8.7D). The posterosinusid is blocked by a cingular ridge in 18 out of 44 cases; it is open in the rest (Fig. 8.7A; 8.7D).

m2—The posterior roots of the molar is split in 28 out of 29 cases; it is partially split in 16 out of 28 and is completely split in 12. The labial branch of the anterolophid is always present (Fig. 8.7B; 8.7E), but it does not reach the protoconid in three out of 44 teeth. Two out of 39 molars have a weak lingual anterolophid. Twenty-eight out of 72 specimens show mesolophid; it is short (23/28; Fig. 8.7B) or incipient (2/28); it is curved and connected with the metaconid in three out of 28 (Fig. 8.7E). Only two out of 75 teeth have ectomesolophid. The lingual posterolophid

is constricted behind the hypoconid and its lingual part is cuspidate (Fig. 8.7B; 8.7E). There is no lingual anterosinusid (Fig. 8.7B; 8.7E) and the mesosinusid is closed by a low cingulid in two out of 58 fossils; it remains open in the rest (Fig. 8.7B; 8.7E). The sinusid can be blocked by a cingulid in 43 out of 51 cases (Fig. 8.7B; 8.7E) or by two crests attached to the protoconid and the hypoconid in six out of 51; it is open in the rest. A weak labial posterosinusid is formed on the posterior wall of the hypoconid in 23 out of 27 teeth (Fig. 8.7E). The lingual posterolophid reaches the entoconid almost always (72/75; Fig. 8.7B; 8.7E).

m3—The labial branch of the anterolophid is always present and it reaches the protoconid in 74 out of 80 cases (Fig. 8.7C; 8.7F). Two specimens have lingual anterolophid. Thirty-four out of 101 teeth have mesolophid (Fig. 8.7F) and it is short (28/34) or long, but it does not exceed the half length of the mesosinusid (6/34). The ectomesolophid is absent. One tooth shows enamel coated valley. The lingual posterolophid is high and extends towards the entoconid. The hypolophulid is short and connected to the anterior branch of the hypoconid. The sinusid is closed in 59 out of 80 molars, by a cingulid in 54 (Fig. 8.7C), by a stylid in one, and by two crests attached to the protoconid and the hypoconid in four cases. The mesosinusid is closed by a low cingulid in three out of 101 specimens; it is open in the remaining ones (Fig. 8.7C; 8.7F). The lingual posterosinusid is closed in 43 out of 66 cases; it is blocked by a cingulid (4/43) or by the lingual posterolophid (39/43).

***Hispanomys aguirrei* from other sites in the Calatayud-Daroca Basin**

The most characteristic features of the studied upper molars are: anterior ectoloph frequently double and complete, but also simple and incomplete; posterior ectoloph, frequently complete, composed by both anterior ectoloph of the metacone and posterior ectoloph of the paracone; cingula in the labial valleys on the M1 and poor-developed mesoloph, which is absent in the M3; posteroloph commonly present in the M3, and well developed in some specimens. For the lower molars: the presence of metalophulid I and II in the same specimen, or metalophulid I, in the m1; mesolophid and ectomesolophid highly reduced or absent, and three-rooted m2 (Appendix 8.1).

Remarkable morphological differences among the material from the different studied sites are not observed. Hence, we consider that the whole studied sample belongs to the same species.

However, there is certain degree of variability in several characters, or states of character, among the fossils from SOL, NOM2, and CARR. In general, these specimens possess differences in the frequencies of presence for a number of states of character. The specimens from SOL show more basal features than those from NOM2, whereas the material from CARR is more derived. For instance, from SOL to CARR, there is an increase of the complexity of the ectolophs. This

is shown by the percentage of double and complete ectolophs (Figs. 8.6, 8.8; Appendix 8.1). Almost half of the M1 and M2 from SOL display short mesoloph, whereas it is absent in most upper molars from CARR (Figs. 8.6, 8.8; Appendix 8.1). The sinus is open in fewer specimens from SOL than from CARR (Figs. 8.6, 8.8; Appendix 8.1). In CARR the M3 have posteroloph more frequently than in the rest of the sites (Figs. 8.6, 8.8; Appendix 8.1). The m1 from SOL still have metalophulid II, whereas most m1 from CARR have metalophulid I or both metalophulids (I+II) in the same specimen (Figs. 8.7, 8.8; Appendix 8.1). The mesolophid of the m2 and m3 is connected with the metacone in SOL and NOM2 but not in CARR (Figs. 8.7, 8.8; Appendix 8.1). To sum up, features such as the short mesoloph, the metalophulid II in the m1, and the mesolophid connected with the metacone, are indicative of the basal morphology of the specimens from SOL. By contrast, traits such as the absence of mesoloph, the presence of metalophulid I in the m1, and the absence of mesolophid, indicate the more derived morphology of the material from CARR.

Discussion

The studied specimens fit within the size range of *Hispanomys aguirrei* from its type locality Escobosa de Calatañazor (Duero Basin, Spain). The fossils from NOM2, were previously assigned to *H. aguirrei* by López-Guerrero et al. (2008), and those from NOM3 and NOM4 were already assigned to *Hispanomys* cf. *aguirrei* by López-Guerrero et al. (2009); here we maintain this adscription. Similarly, we consider that the fossils from PJE1 and PJE2 also belong to *Hispanomys* cf. *aguirrei* due to the scarce material available. The M1 from Paje localities are similar to those from Escobosa, but they present medium-length mesoloph and open protosinus in some M1 which are not present in the type locality.

Hispanomys aguirrei was described by Sesé Benito (1977) in Escobosa de Calatañazor I, (Duero Basin, middle Miocene, MN7/8). This species was also recovered in Escobosa M and Escobosa G-2, (MN7/8). Sesé Benito diagnosed the new species in López-Martínez et al., 1977. However, detailed descriptions and discussion were presented in Sesé Benito (1977). She described the new species under the genus *Cricetodon*, and compared it with the other representatives of the genus. Later on, Sesé Benito (1980) re-assigned the species to *Hispanomys* following Mein and Freudenthal (1971a) and Van de Weerd (1976). The reasons that the author gave are: “[Mein and Freudenthal 1971 created the subgenus *Hispanomys* within the genus *Cricetodon* which Weerd 1976 erected to genus category. The meaning of the genus *Hispanomys*, in the original diagnosis of Mein and Freudenthal, as well as in the emended diagnosis of Weerd is very wide. We include the species of Escobosa in this broad sense of the genus *Hispanomys*.]” (Sesé Benito, 1980:261; translated from Spanish). Since then, this change is widely accepted (e.g., Agustí, 1982;

López-Guerrero et al., 2008, 2009; López-Antoñanzas et al., 2010, 2011).

Apart from the type locality, *H. aguirrei* was found in Nombrevilla 2, 3 and 4 (López-Guerrero et al., 2008, 2009). *Hispanomys* cf. *aguirrei* was also reported from the Abocador de Can Mata series in the Vallès-Penedès Basin, which is dated between 12.5 and 10.6 Ma (Alba et al., 2011).

The most characteristic traits of *H. aguirrei* from the Calatayud-Daroca Basin are: the slightly divided anterocone; the degree of development of the ectolophs; the absence of mesoloph on the M3; the presence of cingula on the labial valleys from the M1; the presence of styles on the sinus and protosinus of the M1; the absence of metalophulid II solely in the m1; and the frequencies of presence of metalophulid I, both metalophulids (I+II), and short mesolophids in the m1. Some of these are basal features for the tribe (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas and Mein, 2009). By contrast, the ectolophs are double and frequently complete; and the labial anterolophid does not reach the protoconid in some m1 which are advanced traits.

Some characters are more frequent in the material from Escobosa de Calatañazor than in the Calatayud-Daroca Basin such as: the double ectoloph of the M1 and the mesoloph on the M2. Besides, the posterior ectoloph of the M3 is always complete in the material from the Duero Basin, but it remains uncompleted in some specimens from the Calatayud-Daroca Basin; moreover, the mesoloph is less frequent in the M3 from Escobosa.

The number of specimens from Escobosa and NOM2 are enough to consider that the differences in the relative frequencies of some states of character between both localities are not due to the sample size. According to the trends described for the tribe (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas and Mein, 2009), these differences may reflect a more derived morphological pattern of the molars from Escobosa de Calatañazor. Therefore, the variations between both assemblages could be the result of changes through time according to the referred trends. The scatter-diagram (Fig. 8.2) indicates that the material from the Calatayud-Daroca Basin is very similar to that from the type locality. However, the ANOVA test reveals particular significant differences between localities (Fig. 8.1; Table 8.8a), which have been evaluated by means of Post-Hoc tests (Table 8.8b). It can be observed that there are differences in length and width of the M1, M2, and m2, and SOL is the locality with higher number of significant differences. Figure 8.2 shows that SOL and PJE have the smallest size. These localities are also the youngest. This could reflect a trend towards size increase explaining the significant differences.

Hispanomys aguirrei was assigned either to *Cricetodon* or to *Hispanomys* (e.g., Sesé Benito, 1977, 1980; Agustí et al., 2005; López-Guerrero et al., 2008). The species could fit in any of these genera; it depends on the importance given to a particular character. Thus, the generic assignment of *H. aguirrei* has to consider carefully every morphometrical character. Therefore, the study of *H.*

aguirrei from Calatayud-Daroca Basin is very important, providing valuable information about the intraspecific variability and the morphological trends of this species. This information will be incorporated in the cladistic analysis (chapter 10) and the posterior taxonomical discussion of the species *H. aguirrei*.

Similarly, chapter 10 treated the proposed phylogenetic relationships regarding *H. aguirrei*. As explained in chapters 6 and 7, Agustí (1982) proposed that *C. jotae* was the ancestor of *H. aguirrei*. He defined an evolutionary trend in *C. jotae*-*H. aguirrei* towards the development of the ectolophs, the increase of the hypsodonty and the reduction of the cingular formations. Another phylogenetic hypothesis involves *Hispanomys lavocati* (Freudenthal, 1966). Both *H. aguirrei* and *H. lavocati* co-occur in some localities such as NOM2, NOM3, and NOM4 (López-Guerrero et al., 2008, 2009). *Hispanomys lavocati* shows a general pattern (ectolophs always present in the upper molars, some of them double and frequently complete; absence of metalophulid II on the m1; and a three-rooted m2, being the posterior root completely split in most of them) which is more derived than *H. aguirrei* following the evolutionary trends described (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas and Mein, 2009) and both species could be phylogenetically related.

The main questions about *H. aguirrei* (its relationship with *C. jotae* and *H. lavocati* and its generic assignation) will be evaluated and discussed in chapter 10 through a cladistics analysis.




FIGURE 8.2. Length/Width scatter-diagrams of the cheek teeth of *Hispanomys aguirrei* and *Hispanomys cf. aguirrei* from the Calatayud-Daroca Basin. The data from the type locality, Escobosa de Calatañazor (Duero Basin, Spain), are also included. Abbreviations: See table 8.2.




FIGURE 8.6. Upper molars of *Hispanomys aguirrei* and *Hispanomys* cf. *aguirrei* from the Calatayud-Daroca Basin. *Hispanomys aguirrei*: Nombrevilla 2: A, left M1, NOM2-1673 ; B, left M2, NOM2-1722; C, left M3, NOM2-1854; D, inverted right M1, NOM2-1601; E, left M2, NOM2-1785; F, inverted right M3, NOM2-1821. *Hispanomys* cf. *aguirrei*: Nombrevilla 3: G, inverted right M1, NOM3-01; H, inverted right M2, NOM3-11; Nombrevilla 4: I, left M1, NOM4-14. Paje 1 and 2: J, left M1, PJE2-95; K, left M2, PJE1-9; L, left M3, PJE2-100; M, inverted right M1, PJE2-97; N, left M2, PJE2-52; O, left M3, PJE2-98. Scale bar equals 1 mm.




FIGURE 8.7. Lower molars of *Hispanomys aguirrei* and *Hispanomys* cf. *aguirrei* from the Calatayud-Daroca Basin. *Hispanomys aguirrei*: Nombrevilla 2: A, left m1 NOM2-1949, B, inverted right m2, NOM2-2006; C, inverted right m3, NOM2-2116; D, inverted right m1, NOM2-1922; E, left m2, NOM2-2037; F, m3, NOM2-2177. *Hispanomys* cf. *aguirrei*: Paje1: G, left m1, PJE1-14. Nombrevilla 3: H, inverted right m2, NOM3-21; I, inverted right m3, NOM3-46. Paje2: J, inverted right m1, PJE2-104; K, inverted right m2, PJE2-105; L, inverted right m3, PJE2-109; M, inverted right m1, PJE2-55; N, left m2, PJE2-57; O, left m1, PJE2-53. Scale bar equals 1 mm.




FIGURE 8.8. *Hispanomys aguirrei* from the Calatayud-Daroca Basin. Carrilanga. Upper molars: A, left M1, RGM568253; B, inverted right M2, RGM568301; C, left M3, RGM568383; D, inverted right M1, RGM568273; E, left M2, RGM568319; F, left M3, RGM568378; G, inverted right M1, RGM568260; H, inverted right M3, RGM568366. Lower Molars: I, left m1, RGM568286; J, inverted right m1, RGM568281; K, left m2, RGM568353; L, left m3, RGM568400; M, left m1, RGM568291; N, inverted right m2, RGM568333; O, inverted right m3, RGM568386. Scale bar equals 1 mm.

HISPANOMYS LAVOCATI (Freudenthal, 1966)

(Figs. 8.3; 8.9-8.10; Tables 8.2, 8.5)

Original Type locality—Hostalets de Pierola, Vallès-Penedès Basin, Spain (MN7/8, upper Aragonian, middle Miocene).

Stratigraphic Range in the Calatayud-Daroca Basin—local biozone G3, (MN7/8, upper Aragonian, middle Miocene).

Geographical range—Southwestern Europe.

Measurements—See Table 8.5.

Diagnosis

Original Diagnosis (Freudenthal, 1966:311)—“Larger than *Cricetodon meini*, *decedens*, etc. Comparable in size with *Cricetodon sansaniensis*. The typical *decedens*-pattern, the labial lophs connecting anteroloph, paracone and metacone, is [sic] well developed.”

HISPANOMYS LAVOCATI FROM THE CALATAYUD-DAROCA BASIN

Description of the Material from Nombrevilla 2

M1—This molar has four roots and a sinuous outline. The anterocone is large; it is divided in two parts by a shallow groove. The connection between protocone and anterocone is through the anterolophule, which is connected to the lingual part of the anterocone (Figs. 8.9A; 8.9D). A posterior protolophule connects the paracone with the posterior branch of the protocone (Figs. 8.9A; 8.9D). The anterior protolophule and the labial spine of the anterolophule are absent (Figs. 8.9A; 8.9D). The anterior ectoloph is always present; it is composed by the posterior ectoloph of the anterocone (46/6; Figs. 8.9A; 8.9D) or it is double, composed by the posterior ectoloph of the anterocone and the anterior ectoloph of the paracone (20/66). The anterior ectoloph is complete in 42 out of 68 fossils. The posterior ectoloph is always present, composed by the posterior ectoloph of the paracone (15/60; Figs. 8.9A; 8.9D); it is double, composed by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone, in 45 out of 60 specimens (Figs. 8.9A; 8.9D). The posterior ectoloph is complete in 54 out of 60 cases (Figs. 8.9A; 8.9D). Nine out of

61 teeth have mesoloph, which is incipient in four; it is clearly short in three, and long, but not reaching the labial border, in two. The enamel coated valley and the entomesoloph are absent (Figs. 8.9A; 8.9D). One specimen possess a crest in the mesosinus that connects the ectoloph with the longitudinal crest (Fig. 8.9G). All teeth display labial posteroloph (Figs. 8.9A; 8.9D), whereas the lingual posteroloph is absent in nine out of 25 (Figs. 8.9A; 8.9D). The anterosinus is always closed, by a cingulum (28/53; Fig. 8.9D), by a low cingulum (4/53; Fig. 8.9A), or by a style (21/53). The mesosinus is blocked by a cingulum (62/63; Figs. 8.9A; 8.9D) or by a style (1/63). The sinus is open in 21 out of 62 molars (Figs. 8.9A; 8.9D); it can be blocked by a cingulum (24/63), by a style (14/63), or by two crests attached to the hypocone and protocone respectively (3/63). The protosinus is mostly closed (57/59); it can be blocked by a cingulum (32/57; Fig. 8.9D), by a crest attached to the anterocone (20/57; Fig. 8.9A), or by a style (5/57).

M2—The anterior ectoloph is present in 15 out of 56 cases; it is always complete and composed by the anterior ectoloph of the paracone. All teeth have posterior ectoloph, which is composed by a posterior ectoloph of the paracone (29/59; Fig. 8.9E) or by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone (30/59; Fig. 8.9E). The posterior ectoloph is complete in 28 out of 59 (Fig. 8.9B). Only three out of 58 fossils have mesoloph (Figs. 8.9B, 8.9E); it is short in two cases, and incipient in one tooth. The entomesoloph and the enamel-coated valley are absent (Figs. 8.9B, 8.9E). There is a labial posteroloph (Fig. 8.9E) in 37 out of 49 specimens. The lingual posteroloph is absent in 45 out of 52 molars and it is short in the rest (Figs. 8.9B, 8.9E). The protosinus is weak but well distinguished (Figs. 8.9B, 8.9E) in all cases but one. The mesosinus is always blocked by a cingulum (Figs. 8.9B, 8.9E); it is low in four specimens. The sinus is open (15/56; Fig. 6.9B); it is closed by a cingulum (33/56; Fig. 8.9E) or by a style (8/56).

M3—The lingual anteroloph is weak but clearly distinguishable in 22 out of 83 cases (Figs. 8.9C, 8.9F). The labial anteroloph is always present and well developed (Figs. 8.9C, 8.9F). The anterior ectoloph is mostly absent (75/76; Figs. 8.9C, 8.9F); it is composed by the posterior ectoloph of the paracone in one tooth. Most of the M3 possess posterior ectoloph (75/77); it can be simple (43/ 75; Figs. 8.9C, 8.9F) or double (32/75). The posterior ectoloph of the paracone can be attached to the paracone (47/75) or to the protolophule (28/75; Figs. 8.9C, 8.9F). In 56 specimens the posterior ectoloph is complete. The longitudinal ridge is posteriorly interrupted, being the hypocone isolated (Figs. 8.9C, 8.9F). Only five out of 77 teeth have mesoloph; it is long and reaches the lingual border in 3 of them (Figs. 8.9C, 8.9F). One tooth displays enamel-coated valley. The axioloph is interrupted in 13 out of 77 molars (Figs. 8.9C, 8.9F) and the neo-entoloph is absent (Figs. 8.9C, 8.9F). In 56 out of 77 cases there is a well-developed posteroloph (Figs. 8.9C, 8.9F). The sinus is long, narrow, and straight (Figs. 8.9C, 8.9F).

m1—The labial anterolophid do not reach the protoconid in some specimens (15/49; Figs.

8.10A, 8.10D). The labial anterolophid connects the anteroconid with the protoconid in 34 out of 49 teeth. The lingual anterolophid is absent (Figs. 8.10A, 8.9D). The metalophulid I is always present (Figs. 8.10A, 8.10D), whereas the metalophulid II is always absent. In 21 out of 53 cases there is a mesolophid; it is short in 16 and incipient in five. The ectomesolophid is absent (Figs. 8.10A, 8.10D). The mesosinus is blocked by a low cingulid (7/51; Fig. 8.10D) or by a stylid (1/51); it is open in the rest (Fig. 8.10A). The sinusid is narrow and long. It is backwards directed (Figs. 8.10A, 8.10D); it can be blocked by a low cingular ridge (24/45; Fig. 8.10D), by two crests attached to the hypoconid and the protoconid (17/45), or open (4/45; Fig. 8.10A). The posterosinusid is blocked by a cingular ridge in three out of 52 fossils (Fig. 8.10A); it is open in the remaining ones (Fig. 8.10D).

m2—This molar has always three roots; the posterior one can be completely (6/16) or partially split (10/16). The labial branch of the anterolophid is well developed and always reaches the protoconid (Figs. 8.10B, 8.10E). The lingual anterolophid is absent (Figs. 8.10B, 8.10E). Twenty-nine out of 51 teeth have mesolophid; it can be short (26/51; Fig. 8.10E), or incipient (3/51). The ectomesolophid is absent (Figs. 8.10B, 8.10E). There is no anterosinusid and the mesosinusid is blocked by a low cingular ridge in 15 out of 47 cases (Figs. 8.10B, 8.10E), and is open in the remaining ones. The sinusid is closed by a cingulid (27/46) or by two crests attached to the hypoconid and protoconid respectively (19/46; Figs. 8.10B, 8.10E). A weak labial posterosinusid is formed on the posterior wall of the hypoconid in most specimens (55/57; Figs. 8.10B, 8.10E). The lingual posterolophid always reaches the entoconid (Figs. 8.10B, 8.10E).

m3—The labial branch of the anterolophid is long and reaches the protoconid (Figs. 8.10C, 8.10I). The lingual branch of the anterolophid is absent (Figs. 8.10C, 8.10I). The mesolophid is present in 13/72 specimens; it is short and develops a second branch in seven cases. The ectomesolophid is absent (Figs. 8.10C, 8.10I). The hypolophulid, which is short, is connected to the anterior branch of the hypoconid (Figs. 8.10C, 8.10I). The mesosinusid is open in 24 out of 68 teeth (Figs. 8.10C, 8.10I) or closed by a low cingulid (44/68). The sinusid is blocked in 59 out of 71 fossils; it can be closed by a cingulid (52/59; Figs. 8.10C, 8.10I), by two crests attached to the hypoconid and protoconid respectively (5/59), or by a stylid (2/59). The lingual posterolophid is high and reaches the entoconid (Figs. 8.10C, 8.10I).

Discussion

The material of *H. lavocati* found in NOM2, NOM3, and NOM4 was already studied by López-Guerrero et al. (2008, 2009). These assemblages are morphometrically homogeneous (Figs. 8.3, 8.9, 8.10; Table 8.5). There are only three differences between NOM2, NOM3, and NOM4. The m3 from NOM3 shows lingual anterolophid; the anterior ectoloph of the M2 from NOM4

is formed by the anterior ectoloph of the paracone and the posterior ectoloph of the anterocone and the sinus of one m2 from NOM4 is closed by a stylid. López-Guerrero et al. (2008, 2009) compared *H. lavocati* with the other large sized species of Cricetodontini and with *H. aguirrei*. We maintain here the assignation of the fossils from NOM2 to *H. lavocati* and the material from NOM3, and NOM4 to *Hispanomys* cf. *lavocati*

Hispanomys lavocati (Freudenthal, 1966) is one of the first described Cricetodontini in the Iberian Peninsula. The holotype comes from the lower levels of Hostalets de Pierola (MN7/8) and is also found in Sant Quirze from the Vallès-Penedès Basin. The taxonomy of this species has been very controversial and its generic assignation differs from one study to another. *Hispanomys lavocati* was firstly included within the genus *Cricetodon* by Freudenthal (1966). Due to its high hypsodonty, well-developed ectolophs and big size, Mein and Freudenthal (1971a) placed it in the subgenus *Pseudoruscinomys*. Later, *Pseudoruscinomys* is synonymized with *Ruscinomys* by Van de Weerd (1976) because its “diagnostic features” are present in both *Hispanomys* and *Ruscinomys*. Some of the species of *Pseudoruscinomys* were included in *Hispanomys* and part in *Ruscinomys*. Agustí (1977) ascribed this species to *Hispanomys* but later on (Agustí, 1980) studied the fossils from Hostalets de Pierola concluding that the material belonged to two species that differ in size. The small species was described as *Hispanomys dispectus*, and the large-sized species referred as ‘*Cricetodon*’ *lavocati*. He pointed out that ‘*Cricetodon*’ *lavocati* could not be included in *Cricetodon* due to its well-developed ectolophs but neither in *Hispanomys* because of its large size and non-reduced M3. Therefore, Agustí (1980) defined a new subgenus, *Pararuscinomys*, to allocate the “large species” from Hostalets de Pierola. However, this change was not followed by other authors and recently, Casanovas-Vilar and Agustí (2007) named the large species from Hostalets de Pierola as *H. lavocati* (Table 8.11). This question will be addressed in chapter 10 taking into account the results of the cladistic analysis.

In the present work *Hispanomys lavocati* is considered a basal member of *Hispanomys* because it has some features in common with other species of this genus. These are: the sinuous contour; the poor development of cingular formations; the ectolophs well developed, specially the anterior ectoloph of the metacone; the anterocone completely split on the M1; the poorly developed labial anterolophid; the absent metalophulid II and the three-rooted m2. Similarly, it is considered as a basal species of the genus because it shows some of the ectolophs not complete; mesolophs in the upper molars; the non-reduced M3; a weak posteroloph on some M3 and some stylids in the lower molars.

There are two main hypotheses about the phylogenetic relationships of *H. lavocati*. The first one assumes an endemic origin from the Iberian Peninsula (Agustí, 1982) and the other hypothesis states that it probably has an Asiatic origin and represents a latest middle Miocene immigrant from Asia Minor (De Bruijn et al., 1993). *Hispanomys lavocati* from the Calatayud-

Daroca Basin possess some features which have been found in the representatives of *Cricetodon* and *Byzantinia* from Anatolia such as: a large anterocone and a short cingulum between the lobes in some M1. That supports the possible Asian origin of *H. lavocati* as well as the existence of a possible common ancestor between *Byzantinia* and *Hispanomys*. These questions will be discussed in the chapter 10. A cladistic analysis and a posterior paleobiogeographical discussion could confirm these hypotheses.




FIGURE 8.3. Length/Width scatter-diagrams of the cheek teeth of *Hispanomys lavocati* and *Hispanomys cf. lavocati* from the Calatayud-Daroca Basin. Abbreviations: See table 8.2.




FIGURE 8.9. Upper molars of *Hispanomys lavocati* and *Hispanomys* cf. *lavocati* from the Calatayud-Daroca Basin. *Hispanomys lavocati*: Nombrevilla 2: A, left M1, NOM2-1073; B, inverted right M2, NOM2-1114; C, inverted right M3, NOM2-1200; D, left M1, NOM2-1035; E, left M2, NOM2-1140; F, left M3, NOM2-1275; G, left M1, NOM2-1067; H, left M2, NOM2-1156; I, left M3, NOM2-1255; J, left M1, NOM2-1044. *Hispanomys* cf. *lavocati*: Nombrevilla 4: K, left M2, NOM4-21; Nombrevilla 3: L, left M3, NOM3-36; M, inverted right M1, NOM4-12. Scale bar equals 1 mm.




FIGURE 8.10. Lower molars of *Hispanomys lavocati* and *Hispanomys* cf. *lavocati* from the Calatayud-Daroca Basin. *Hispanomys lavocati* Nombrevilla 2: A, left m1, NOM2-1329; B, inverted right m2, NOM2-1415; C, inverted right m3, NOM2-1526; D, inverted right m1, NOM2-1318; E, left m2, NOM2-1445; F, left m3, NOM2-1569; G, left m1, NOM2-1343; H, left m2, NOM2-1424; I, left m3, NOM2-1574; J, inverted right m3, NOM2-1523; K, left m2, 1437. *Hispanomys* cf. *lavocati*: Nombrevilla 4: L left m3, NOM4-40; M, left m1, NOM4-04. Scale bar equals 1 mm.

HISPANOMYS NOMBREVILLAE (Freudenthal, 1966)

(Figs. 8.4; 8.11–8.12; Tables 8.2, 8.6, Appendix 8.2)

Original Type locality—Nombrevilla 1, Calatayud-Daroca Basin, Spain, local biozone H (MN9, lower Vallesian, upper Miocene).

Stratigraphical Range in the Calatayud-Daroca Basin—local biozone H (MN7/8–MN9, upper Aragonian–lower Vallesian, middle–upper Miocene).

Geographical range—Southwestern Europe.

Measurements—See Table 8.6.

Diagnosis

Original Diagnosis (Freudenthal, 1966:3)—“Resembling *Cricetodon decedens decedens* [sic] Schaub, 1925, from La Grive in size and dental pattern. Mesolophs are absent, mesolophids are either absent or short. The lophs connecting the anteroloph, paracone and metacone are well developed. Smaller than *C. sansaniensis* Lartet, 1851 and *C. meini* Freudenthal, 1963. The width of the upper molars is smaller than in *C. decedens decedens* [sic]. Length/width ratio of upper molars smaller than in *C. decedens aragonensis*, nov. subsp., [sic].”

Emended diagnosis—Small-sized Cricetodontini. The M1 has four roots. The anterior ectoloph is always present on the M1 and M2 and it can be absent on the M3; it is mainly composed by the posterior ectoloph of the anterocone and almost always complete. The anterior ectoloph can be also composed by both anterior ectoloph of the paracone and posterior ectoloph of the anterocone. The posterior ectoloph is always present in the upper cheek teeth; it is composed mainly by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone; it is almost always complete. The mesoloph is absent in the M1 and M2 and in some M3. The main valleys are open or closed. The M3 is highly reduced and the hypocone and metacone are barely distinguishable. Half of the lower molars possess mesolophid; it is frequently poorly developed and can be connected with the metacone. The metalophulid I is always present, some specimens have also metalophulid II. Most m2 display three roots totally split.

Differential diagnosis—Differs from ‘*Cricetodon*’ *klariankae* Hír, 2007, by its smaller size; absence of a lingual spine on the protosinus of the M1 and complete anterior ectoloph in all molars. Differs from ‘*Cricetodon*’ *fandli* Prieto, Böhme and Gross, 2010 by its smaller size and highly reduced M3. Differs from *Hispanomys dispectus* Agustí, 1980 by its less developed ectolophs in the

M1 and the presence of metalophulid II in the m1. Differs from *H. daamsi* Agustí, Casanovas-Vilar and Furió, 2005 by its lower size; its more reduced M3 and the presence of metalophulid II in the m1. Differs from *H. lavocati* (Freudenthal, 1966) by its smaller size; more developed and complete ectolophs in the upper molars and presence of metalophulid II in the m1. Differs from *H. aguirrei* (Sesé Benito, 1977 in López Martínez et al., 1977) by its smaller size; more developed ectolophs and absence of posteroloph in the M3. Differs from *H. bijugatus* Mein and Freudenthal, 1971a by its smaller size; presence of cingulum in the anterosinus and mesosinus and highly reduced M3. Differs from *H. decedens* (Schaub, 1925) by its absence of enamel-coated valley; more developed ectolophs and highly reduced M3 with posteroloph. Differs from *H. castelnovi* Aguilar, Calvet and Michaux, 1994 by its absence of metalophulid I in the m1; more developed ectolophs and deeply split anterocone. Differs from *H. adroveri* Agustí, 1986 by its smaller size; presence of four roots in the M1; presence of labial anterolophid in the m1 and less hypsodonty. Differs from *H. baixasi* Aguilar, Michaux and Lazzari, 2007 by its smaller size; presence of labial anterolophid in the m1; absence of metalophulid I in the m1 and more developed ectolophs. Differs from *H. peralensis* Van de Weerd, 1976 by its smaller size; presence of four roots in the M1 and absence of metalophulid I in the m1. Differs from *H. moralesi* López-Antoñanzas, Peláez-Campomanes, Álvarez-Sierra and García-Paredes, 2010 by its presence of labial anterolophid and absence of metalophulid I in the m1. Differs from *H. mediterraneus*, Aguilar 1982 by its lower size and presence of posteroloph in a highly reduced M3. Differs from *H. thaleri* (Hartenberger, 1965) by its smaller size; more developed posterior ectoloph; presence of enamel-coated valley and absence of metalophulid I in the m1. Differs from *H. aragonensis* (Freudenthal, 1966) by its less developed ectolophs; absence of enamel-coated valley; higher reduced M3 and presence of cingula surrounding the upper molars.

HISPANOMYS NOMBREVILLAE FROM THE CALATAYUD-DAROCA BASIN

Description of the Material from Nombrevilla 1.

M1—This molar has four roots. The anterocone is large and divided in two parts by a shallow groove in the anterior wall. These two parts are fused when the tooth is worn; however, a constriction on the enamel of the anterocone is always found. One tooth shows a labial anteroloph extended through the protosinus. A posterior protolophule connects the paracone with the posterior branch of the protocone. In eight out of 15 cases, the protolophule is transversal and connected to the longitudinal crest. The anterior protolophule and the labial spine of the anterolophule are absent. All the specimens have anterior ectoloph; it is composed by the anterior ectoloph of the paracone (6/15) or by the posterior ectoloph of the anterocone (2/15). In seven teeth the anterior ectoloph is double, composed by both anterior ectoloph of the paracone and

posterior ectoloph of the anterocone. The anterior ectoloph always forms a continuous connection between the anterocone and the paracone, although in one tooth this connection is lower than the labial cusps. The posterior ectoloph is always present; it is composed by the posterior ectoloph of the paracone (4/14) or by both anterior ectoloph of the paracone and posterior ectoloph of the metacone (10/14). The posterior ectoloph is always complete. There is no mesoloph, nor enamel-coated valley. The anterior wall of the protocone bears a ridge-like structure in eight out of 14 specimens. Two teeth possess entomesoloph. The metalophule is posterior and connected to the labial posteroloph. Both labial and lingual posteroloph are present. The anterosinus is closed by a cingulum (2/16) or by a low cingulum (13/16), and it is open in one case. The mesosinus is closed by a cingulum (5/16) or by a low cingulum (10/16), and it is open in one tooth. The sinus is mostly closed (8/17), by a cingulum (6/8), by a style (1/8) or by two crests attached to the hypocone and protocone respectively (1/8). The protosinus is closed by a cingulum (6/12) or by a style (5/12).

M2—The anterior ectoloph is always present and complete (Fig. 8.11H); it is composed by the posterior ectoloph of the anterocone (3/18), by the anterior ectoloph of the paracone (9/18) or by both (6/18; Fig. 8.11H). All teeth have posterior ectoloph and it is composed by the posterior ectoloph of the paracone (2/19) or by both posterior ectoloph of the paracone and anterior ectoloph of the metacone (17/19; Fig. 8.11H). The posterior ectoloph is always complete (Fig. 8.11H). The mesoloph, entomesoloph and enamel-coated valley are absent (Fig. 8.11H). The protocone always bears a ridge-like structure (Fig. 8.11H). Both labial and lingual posterolophs are found in all specimens (Fig. 8.11H). Eighteen out of 19 cases have lingual anterosinus; it is open (11/18) or closed by a low cingulum (7/18). The protosinus is weak but clearly distinguishable in all the molars but one; it is open (12/19) or closed by a style (7/19). The mesosinus is open (13/19; Fig. 8.11H) or blocked by a low cingulum (6/19). The sinus is open (5/17; Fig. 8.11H) or closed by a cingulum (12/17). In one tooth the sinus is blocked by two crests attached to the hypocone and to the protocone, respectively (Fig. 8.11H).

M3—The hypocone is barely distinguishable (Fig. 8.11I). The lingual anteroloph is absent (Fig. 8.11I). The labial anteroloph is always present (Fig. 8.11I); it is weak (3/12) or well developed (9/12; Fig. 8.11I). Nine out of 12 fossils possess anterior ectoloph; it is composed by the posterior ectoloph of the anteroloph (6/9) or by the anterior ectoloph of the paracone (3/9). The anterior ectoloph is always complete (Fig. 8.11I). All teeth have complete posterior ectoloph (Fig. 8.11I); it is composed by the posterior ectoloph of the paracone (1/12; Fig. 8.11I) or by both posterior ectoloph of the paracone and anterior ectoloph of the metacone (11/12; Fig. 8.11I). Only one tooth has mesoloph and it is short. The entomesoloph and the enamel-coated valley are absent (Fig. 8.11I). The axioloph is continuous (Fig. 8.11I) and the neo-entoloph is found in two out of 12 cases. The posteroloph is absent (Fig. 8.11I). The sinus is long, curved and forward directed in the specimens without neo-entoloph (Fig. 8.11I) and is small and straight when the neo-entoloph is missing.

m1—This molar has two roots. The labial anterolophid is a well-developed ridge that reaches the protoconid in 11 out of 17 specimens. The lingual anterolophid is usually absent (13/16); it is displayed in three and it is short. The metalophulid I is found in 16 out of 20 specimens. Both metalophulids, I and II, are present in the same tooth in four cases. Some teeth (9/20) have mesolophid; it can be an enlargement of the enamel (5/9) or short (4/9). The ectomesolophid is absent. The anterosinusid is open (12/15) or closed by the lingual anterolophid (3/15). The mesosinusid is open (5/21) or closed by a cingulum (16/21), which can be low (3/16). The sinusid is narrow and short; it is oblique and closed by a cingular ridge (17/19), by two crests attached to the hypoconid and the protoconid (1/19), or by a stylid (1/19). The posterosinusid is open (13/20) or blocked by a cingular ridge (7/20).

m2—This molar has three roots. The posterior root is totally (6/7) or partially split (1/7). The labial branch of the anterolophid is well developed and always reaches the protoconid. The lingual anterolophid is absent. The mesolophid is present in 13 out of 20 cases; it is short (6/13); curved and connected to the metaconid (6/13) or double (1/13). The ectomesolophid is absent. There is no lingual anterosinusid and the mesosinusid is open (4/19) or closed (15/19), either by a low cingulid (14/15) or by a stylid (1/15). The sinusid is always closed; it is blocked by a cingulid (10/13), by a stylid (2/13), or by two crests attached to the protocone and hypocone. A labial posterosinusid is formed on the posterior wall of the hypoconid in five out of 14 specimens, being well developed in one of them. The lingual posterosinusid is open (9/17), blocked by a cingular ridge (7/17), or blocked by the lingual posterolophid (1/17).

m3—The labial branch of the anterolophid is found in 18 out of 20 cases; it reaches the protoconid in 15 out of 18. The lingual branch of the anterolophid is barely present (1/20) and it does not reach the metaconid. Five out of 20 teeth possess a short mesolophid; it can develop a second branch (2/5). The ectomesolophid is absent. The lingual posterolophid is high and extends towards the entoconid. A short hypolophulid is connected to the anterior branch of the hypoconid. The sinusid is open (1/20) or closed (19/20); it is blocked by a cingulid (17/19) or by a stylid (2/19). The mesosinusid is open (16/20) or closed by a low cingulid (4/20). The lingual posterosinusid is closed (13/16); it is blocked by a cingulid (12/13) or by a stylid (1/13).

***Hispanomys nombrevillae* from other sites in the Calatayud-Daroca Basin**

The assemblages of *H. nombrevillae* found in the Calatayud-Daroca Basin do not display important changes in morphology or size. However, there is certain degree of variability in several characters, or states of character, that are considered as part of the variability of the species. A complete summary of the states of character found in the studied molars is presented in the Appendix 8.2.

Some differences are found in the fossils from NOM 9, mainly in the M1, which have not the anterior ectoloph complete. The mesosinus of the M2 is closed by a large cingulum in three out of seven specimens; the sinus is closed by a style in one case and there is an entomesoloph. The posteroloph is distinguishable in two out of four M3 (Fig. 8.11). One out of eight m1 has ectomesolophid. One m3 has long mesolophid and another one has the sinusid closed by two crests (Fig. 8.12).

In the rest of the localities, only four differences are found: in NOM10, one m1 has a weak ectomesolophid; one m1 from CAÑ9 has a double mesolophid (Fig. 8.12); one M3 from CAÑ10 has not complete posterior ectoloph (Fig. 8.11) and one m1 from CAÑ10 lacks the labial anterolophid (Fig. 8.11).

The statistical analysis, Student's *t* test, reveals that there are significant differences between the size of NOM1 and NOM9 in the length of the M2 and the width of the m1 (Fig. 8.4; Table 8.13). Both samples show a large overlap in size (Fig. 8.4) but the M2 from NOM9 are longer than those from NOM1. Similarly, the m1 from NOM9 is slightly narrower than the fossils from NOM1. Given that the number of localities and the short stratigraphical distribution, it is difficult to assess whether these differences follow any evolutionary trend. Therefore, we maintain the specific assignation of the studied material and recommend taking this into account in further works.

Discussion

The fossils from Nombrevilla and the Cañada sections were already assigned to *H. nombrevillae* and *Hispanomys* cf. *nombrevillae* by López-Guerrero et al. (2009, 2011). But detailed morphological descriptions and measurements are provided in the present work. The fossils from NOM10, NOM9, CAÑ8, CAÑ9 and CAÑ10 share a number of features with those from the type locality of *H. nombrevillae* (NOM1) such as: the small size; the degree of development of the ectoloph; the poor-developed or absent mesoloph/ids; the reduction of the cingular formations and, more important, the reduction of the M3. Therefore, we decided to maintain the previous assignations of NOM9 and CAÑ10 as *H. nombrevillae*. The scarcity of the material and the absence of the M1 (highly diagnostic element) in NOM10, CAÑ8, CAÑ9 and CAÑ11 led us to leave the assignation in open nomenclature as *Hispanomys* cf. *nombrevillae*.

Hispanomys nombrevillae was firstly described by Freudenthal (1966) as a subspecies of *Cricetodon decedens*, but it was later allocated in *Hispanomys* by Mein and Freudenthal (1971a) being widely accepted (Agustí, 1982; Álvarez Sierra et al., 2003; López-Guerrero et al., 2009; López-Antoñanzas and Mein, 2009). Apart from the studied localities from the Calatayud-

Daroca Basin, *Hispanomys nombrevillae* is also registered in the Vallesian deposits of the fissure filling from Molina de Aragón (Lacomba Andueza, 1988) but is unknown elsewhere.

The morphology displayed by *H. nombrevillae* is clearly derived among the Cricetodontini: moderately hypsodonty; low size; poor-developed cingula/cingulids; trilobe contour; ectolophs always complete; short or absent mesolophs; highly reduced M3 without posteroloph; absence of metalophulid II in the m1 and three-rooted m2 (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas and Mein, 2009; Table 8.12). However, the same characteristics are considered basal within the genus *Hispanomys*. Therefore, *Hispanomys nombrevillae* is considered a basal member of the genus *Hispanomys* and it is more derived than any species of *Cricetodon*.

Several phylogenies explained the phylogenetic relationships of *H. nombrevillae* (Freudenthal, 1966; Mein and Freudenthal, 1971a; Van de Weerd, 1976; Agustí, 1982). Two species were proposed as the ancestor of *H. nombrevillae*, the French *H. bijugatus* and the Iberian *H. aguirrei*. *Hispanomys bijugatus* possesses enamel-coated valley (Freudenthal, 1966; Agustí, 1980; López-Antoñanzas and Mein, 2009). This trait is used to establish many relationships (Freudenthal, 1966) since it were described only in French-related species. Thus, the absence of enamel-coated valley in *H. nombrevillae* theoretically precludes any relationship with *H. bijugatus*. *Hispanomys nombrevillae* was related with *H. aguirrei* because it possesses more basal characters and it is present in the immediately older localities from the Calatayud-Daroca Basin. Nevertheless, as described in chapter 7, the mosaic morphology of *H. aguirrei* supposes an additional difficulty for establishing the phylogenetic relationships among Cricetodontini. Therefore, we need a cladistic analysis to fully confirm or reject these hypotheses, hence, this question will be addressed in chapter 10.




FIGURE 8.4. Length/Width scatter-diagrams of the cheek teeth of *Hispanomys nombrevillae* and *Hispanomys cf. nombrevillae* from the Calatayud-Daroca Basin. The data from the type locality of *Hispanomys aragonensis*, Pedregueras 2C from Calatayud-Daroca Basin, are also included. Abbreviations: See table 6.2.




FIGURE 8.11. Upper molars of *Hispanomys nombrevillae* and *Hispanomys* cf. *nombrevillae* from the Calatayud-Daroca Basin. *Hispanomys nombrevillae*: Nombrevilla 9: A, left M1, NOM9-01; B, left M2, NOM9-13; C, inverted right M3, NOM9-43; D, inverted right M3, NOM9-38; E, inverted right M2, NOM9-18; F, inverted right M3, NOM9-47. G, left M3, NOM9-48. Nombrevilla 1: H, left M2, NOM1-05; I, inverted right M3, NOM1-04. Cañada 10: J, inverted right M2, CAÑ10-03 K, inverted right M3, CAÑ10-05; *Hispanomys* cf. *nombrevillae*: Nombrevilla 10: L, left M3, NOM10-21; M, inverted right M3, NOM10-23; N, inverted right M3, NOM10-22. Scale bar equals 1 mm.




FIGURE 8.12. Lower molars of *Hispanomys nombrevillae* and *Hispanomys cf. nombrevillae* from the Calatayud-Daroca Basin. *Hispanomys nombrevillae*: Nombrevilla 9: A, inverted right m1, NOM9-22; B, left m2, NOM9-32; C, left m3, NOM9-37; D, inverted right m1, NOM9-23; E, inverted right m2, NOM9-36; F, inverted right m3, NOM9-49; G, left m1, NOM9-25; H, inverted right m1, NOM9-21. Nombrevilla 1: J, inverted right m1, NOM1-02; K, left m3, NOM1-03. Cañada 10: I, left m1, CAÑ10-20. *Hispanomys cf. nombrevillae*: Cañada 8: L, left m2, CAÑ8-23. Cañada 9: M, inverted right m1, CAÑ9-10. Nombrevilla 10: N, left m1, NOM10-31; O, left m1, NOM10-06. Scale bar equals 1 mm.

HISPANOMYS ARAGONENSIS (Freudenthal, 1966)

(Figs. 8.5; 8.13-8.14; Tables 8.2, 8.7, Appendix 8.3)

Material

Original Type locality—Pedregueras 2C, Calatayud-Daroca Basin, Spain, local biozone I (MN9, lower Vallesian, upper Miocene).

Stratigraphical Range in the Calatayud-Daroca Basin—local biozone H-local biozone I, (MN9, lower Vallesian, upper Miocene).

Geographical range—Southwestern Europe.

Measurements—See Table 8.7

Digagnosis

Original Diagnosis (Freudenthal, 1966:303)—“Closely related to the former subspecies. Length/width ratio of the upper molars considerably larger. The lophs connecting anteroloph, paracone and metacone are well developed. Mesolophs may be present. Mesolophids are better developed than in *C. d. nombrevillae* [*sic*]. Ectostylids are present in most m1 and in many m2.”

Emended diagnosis—Small-sized Cricetodontini. The upper molars have well-developed ectolophs. These ectolophs are mainly double and always complete. The valleys are mainly blocked by low cingula, and the protosinus has styles. The M1 has four roots. The posterior ectoloph of the M2 is oblique and connects the metacone to the point where the posterior protolophule reaches the longitudinal crest. Half of the M1 possess lingual anterolophid but it is absent in the M2 and M3. The mesoloph and the enamel-coated valley are frequent in the M3. The M3 often shows a weak posteroloph. The lower molars have well developed labial anterolophid, but only some specimens display lingual anterolophid. The mesolophid is present in most of the lower molars and it can be double or connected with the metacone in some m2 and m3. The m1 frequently shows metalophulid I; some m1 have metalophulid I and II in the same specimen. The m2 has the posterior root completely split.

Differential diagnosis—*Hispanomys aragonensis* differs from ‘*Cricetodon*’ *klariankae* Hír, 2007 by its smaller size and complete anterior ectoloph in all molars. Differs from ‘*C.*’ *fandli* Prieto, Böhme and Gross, 2010 by its smaller size. Differs from *Hispanomys dispectus* Agustí, 1980 by its

less developed ectolophs in the M1 and the presence of metalophulid II in the m1. Differs from *H. daamsi* Agustí, Casanovas-Vilar and Furió, 2005 by its smaller size and presence of metalophulid II in the m1. Differs from *H. lavocati* (Freudenthal, 1966) by its smaller size; more developed and complete ectolophs in the upper molars and by the presence of metalophulid II in the m1. Differs from *H. aguirrei* (Sesé Benito, 1977 in López Martínez et al., 1977) by its smaller size and more developed ectolophs. Differs from *H. bijugatus* Mein and Freudenthal, 1971a by its smaller size; presence of metalophulid I without metalophulid II in the m1 and cingulum in the anterosinus and mesosinus. Differs from *H. decedens* (Schaub, 1925) by its absence of enamel-coated valley; more developed ectolophs and highly reduced M3 with posteroloph. Differs from *H. castelnovi* Aguilar, Calvet and Michaux, 1994 by its more developed ectolophs and deeply split anterocone. Differs from *H. adroveri* Agustí, 1986 by its smaller size; presence of four roots in the M1; presence of labial anterolophid in the m1 and less hypsodonty. Differs from *H. baixasi* Aguilar, Michaux and Lazzari, 2007 by its smaller size; presence of labial anterolophid in the m1 and more developed ectolophs. Differs from *H. peralensis* Van de Weerd, 1976 by its smaller size and the presence of four roots in the M1. Differs from *H. moralesi* López-Antoñanzas, Peláez-Campomanes, Álvarez-Sierra and García-Paredes, 2010 by the presence of labial anterolophid in the m1; presence of metalophulid II and less hypsodonty. Differs from *H. mediterraneus* Aguilar, 1982 by its smaller size and the presence of enamel-coated valley. Differs from *H. thaleri* (Hartenberger, 1965) by its smaller size; the more developed posterior ectoloph and the presence of enamel-coated valley. Differs from *H. nombrevillae* (Freudenthal, 1966) by its more developed ectolophs; presence of enamel-coated valley; less reduced M3 and absence of cingula surrounding the upper molars.

HISPANOMYS ARAGONENSIS FROM THE CALATAYUD-DAROCA BASIN

Description of the Material from Pedregueras 2C.

M1—This molar has four roots. The anterocone is large and divided in two parts. Eight out of 17 cases have lingual anteroloph extended through the protosinus. The anterior protolophule and the labial spine of the anterolophule are absent. The anterior ectoloph is always present; it is composed by the posterior ectoloph of the anterocone (10/18) or by both posterior ectoloph of the anterocone and anterior ectoloph of the paracone (8/18). The anterior ectoloph is always complete, although in one specimen this connection is lower than the labial cusps. All fossils have posterior ectoloph; it is composed by the posterior ectoloph of the paracone (3/19) or by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone (16/19). The posterior ectoloph is always complete. Some molars (3/16) display a long mesoloph which does not reach the labial border. The enamel-coated valley is present in four cases. The anterior wall of the protocone bears

a ridge-like structure in 13 out of 19 specimens. The entomesoloph is absent. The metalophule is posterior and connected to the labial posteroloph. Both labial and lingual posterolophs are well developed. The anterosinus is open (2/16) or closed by a low cingulum (14/16). The mesosinus is open (3/16) or closed by a low cingulum (13/16). The sinus is open (15/17), closed by a style (1/17) or by two crests attached to the hypocone and protocone, respectively (1/17). The protosinus is always closed; either by the lingual anteroloph (8/17), by a cingulum (4/17) or by a style (5/17).

M2—The anterior ectoloph is almost always present (28/29); it is composed by the anterior ectoloph of the paracone (5/28), by the posterior ectoloph of the anterocone (6/28), or by both ectolophs (17/28). The anterior ectoloph is always complete, although in four specimens it is lower than the labial cusps. The posterior ectoloph is always present; it is composed by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone (20/29) or only by the posterior ectoloph of the paracone (9/29). The posterior ectoloph is always complete and it has an oblique position, connecting the metacone to the point where the posterior protolophule reaches the longitudinal crest. Only one tooth displays mesoloph, being part of the enamel-coated valley. Two teeth possess entomesoloph and it is short (1/2) or highly reduced (1/2). The anterior wall of the protocone bears a ridge-like structure in 19 out of 24 specimens. Both labial and lingual posterolophs are always present. The protosinus is weak in 14 out of 26 cases; it is closed by a low cingulum (1/29) or remains open (28/29). The mesosinus is open (13/29) or closed by a low cingulum (12/29), by a cingulum (3/29) or by a style (1/29). The sinus is open (25/27) or closed by a cingulum (2/27).

M3—It is reduced and the hypocone is barely distinguishable. The lingual anteroloph is absent; six specimens have a constriction of the anterior wall of the protocone which is not considered as a true anteroloph. The labial anteroloph is always well developed. The anterior ectoloph is present in 12 out of 40 cases; it is composed by the anterior ectoloph of the paracone (6/12) or by the posterior ectoloph of the anterocone (6/12). The anterior ectoloph is complete in all fossils but one. The posterior ectoloph is present in most of the teeth (40/41). It is composed by the posterior ectoloph of the paracone (5/40) or by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone (35/40). The posterior ectoloph is complete in all specimens but one. Some cases (6/40) show a mesoloph which can be short (4/6) or long (2/6), reaching the labial border in one case. The entomesoloph is absent. Only one out of 40 teeth have enamel-coated valley. The axioloph is continuous in 28 out of 37 and it is interrupted in nine. Three out of 40 specimens bear a neo-entoloph. The posteroloph is frequent (20/35); it can be well developed (8/20) or incipient (12/20). The sinus is long, curved and directed forwards in 28 out of 40 cases; it is transversal in 10, and it is small and straight in two specimens.

m1—The labial anterolophid is always present and reaches the protoconid in 20 out of 46 specimens. The lingual anterolophid is absent. Fourteen out of 46 cases display metalophulid

I and four out of 46 have both metalophulids I and II together. The metalophulid I is joined to the anterolophule in some teeth (12/24), whereas it is joined to the anteroconid in the rest. Twenty-nine out of 46 fossils have mesolophid which is short (18/29) or reduced (11/29). The ectomesolophid is absent. The anterosinusid is mainly open (40/46); it is blocked by a stylid (2/46) or by a cingulid (4/46). The mesosinusid is open (34/47) or blocked by a cingulid (9/47), by a low cingulid (3/47) or by a stylid (1/47). The sinusid is narrow, long, and oblique; it is open (4/44) or closed by a stylid (21/44), by a cingulid (12/44), by two stylids (5/44) or by two crests attached to the hypoconid and the protoconid (2/44). The posterosinusid is open (26/44) or blocked by a cingular ridge (18/44).

m2—This molar has three roots. The posterior root is always totally split. The labial branch of the anterolophid is always present, but it does not reach the protoconid in three out of 25 cases. Ten out of 22 specimens have a weak lingual anterolophid. Twenty-seven out of 30 cases show a short mesolophid. Five out of 27 teeth develop a second branch of the mesolophid; it is curved and connected with the metaconid in 12 out of 27 fossils. The ectomesolophid is absent. There is no lingual anterosinusid. The mesosinusid is open (11/28) or closed by a low cingulid (17/28). The sinusid is closed in all molars but one; it is blocked by a cingulid (13/26), by a stylid (11/26) or by two crests attached to the protoconid and the hypoconid (2/26). A weak labial posterosinusid is formed on the posterior wall of the hypoconid in 23 out of 27 specimens. The lingual posterosinusid is open (9/28), blocked by a cingular ridge (12/28) or blocked by the lingual posterolophid (5/28).

m3—The labial branch of the anterolophid is always present; it reaches the protoconid in 33 out of 36 cases. The lingual anterolophid is displayed by 19 out of 37 fossils. It reaches the metaconid (4/19) or shows a constriction of the anterior wall (15/19). The mesolophid is present in 20 out of 35 teeth; it is short (15/20) or connected to the metaconid (5/20). The ectomesolophid is absent. The lingual posterolophid is high and extends towards the entoconid. The short hypolophulid is connected to the anterior branch of the hypoconid. The sinusid is open (11/35) or closed (24/35); it is blocked by a cingulid (10/24), by a stylid (13/24) or by two crests attached to the protoconid and the hypoconid (1/24). The mesosinusid is open (27/37) or closed by a low cingulid (10/37). The lingual posterosinusid is open (14/44) or closed (30/44); by a cingulid (11/30) or by the lingual posterolophid (19/30).

***Hispanomys aragonensis* from other sites in the Calatayud-Daroca Basin**

The material of the localities from Pedregueras and Nombrevilla possess many morphological similarities (Figs. 8.13, 8.14; Appendix 8.3), there are differences in some structures, although the important features such as the degree of development of the ectolophs are similar in each

assemblage.

In general, the material from the older localities (NOM13, NOM19, NOM22) has more frequently closed valleys and larger structures than the fossils from PED2C. Besides, the upper molars from PED2A have higher frequency of mesoloph on the M1 and a style in the mesosinus of some M2 (Appendix 8.3). The m1 shows lingual anterolophid (11/53 in PED2A); a medium length mesolophid is present in the m1 (8/56 in PED2A; Fig. 8.14A) and m2 (14/44 in PED2A). Additionally, the m3 from PED2A has ectomesolophid. These differences point out that the fossils from the type locality (PED2C) show more derived morphological pattern, according to the trends described for the tribe (Mein and Freudenthal, 1971a; Agustí, 1982; Hír, 2007; López-Antoñanzas and Mein, 2009).

The statistical analysis performed here included the fossil sites PED2A and PED2C. The Student's *t* test reveals significant differences in the length and the width for all elements but the M3 (Table 8.10). The scatter-diagram (Fig. 8.5) indicates that the larger elements from PED2A are absent in PED2C. This absence could produce these results in the statistical analysis.

Discussion

The material from the Pedregueras section assigned to *H. aragonensis* was already described in Freudenthal (1966) and López-Guerrero et al. (2009). The fossils from Nombrevilla are morphometrically very similar to those from Pedregueras. In particular, the material from NOM13 and NOM14 displays double and complete ectolophs in the M1; both labial and lingual posterolophs in the M1 and metalophulid I in the m1. The M2 from NOM22 has oblique ectoloph and the M3 is not much reduced. The m3 from NOM19 possesses a short hypolophulid which is connected to the anterior branch of the hypoconid. All these traits are diagnostic of *H. aragonensis*. However, some elements are still missing in NOM22, NOM13, NOM14 and NOM22. Because of that, we decided to name the specimens from Nombrevilla as *Hispanomys* cf. *aragonensis*.

Freudenthal (1966) established two new subspecies of *Cricetodon decedens*; *Cricetodon decedens aragonensis* was one of them. The generic adscription was later modified by Mein and Freudenthal (1971a), who described the new genus *Hispanomys* and designated *H. aragonensis* as the type species. Apart from the type locality PED2C, Freudenthal (1966) described PED1A and PED2A which contain *H. aragonensis* also and are stratigraphically situated in lower levels (Freudenthal and Sondaar, 1964).

Hispanomys aragonensis has characters which are considered derived for the Cricetodontini, such as: the high development of the ectolophs; reduced M3 and high frequency of metalophulid

I in the m1. By contrast, it shows: four-rooted M1; high frequency of mesolophs on the M1; labial anterolophid in the m1 and structures closing the valleys. The latter traits are considered basal according to the trends described for the genus *Hispanomys* (Agustí, 1982; Agustí et al., 2005). In addition, *H. aragonensis* shows the mesosinus and anterosinus markedly reduced and a sinuous outline. This sinuous contour was proposed as a derivative feature by Aguilar et al. (2007).

Hispanomys aragonensis is endemic from the Iberian Peninsula (Daams et al., 1998; De Bruijn et al., 1992) and its distribution in the Calatayud-Daroca Basin is limited to the lower Vallesian (local biozones H and I, MN9). A younger assemblage of *Hispanomys* cf. *aragonensis* is present in Torremormojón 1 (TM1) from the Duero Basin (local biozone *Progonomys hispanicus*, MN10, late Vallesian) (Álvarez Sierra, 1983; García-Moreno, 1987; Álvarez Sierra et al., 1990). The fossils from this site have shorter molars than those from PED2C, except for the M2. Besides, they lack mesolophids, and cingulids and stylids on the sinusids of the m3. Nevertheless, the differences in the relative frequency of this character cannot be fully discussed due to the small sample size of the assemblage from TM1 in comparison with those from the Calatayud-Daroca Basin.

Some authors (Mein and Freudenthal, 1971a; Agustí, 1982; López-Antoñanzas and Mein, 2011) related *H. aragonensis* to *H. decedens* and stated that *H. decedens* could be the ancestor of *H. aragonensis*. Besides, Van de Weerd (1976) noted that *H. peralensis* could be the descendant of *H.*




FIGURE 8.5. Length/Width scatter-diagrams of the cheek teeth of *Hispanomys aragonensis* and *Hispanomys cf. aragonensis* from the Calatayud-Daroca Basin. Abbreviations: See table 6.2.




FIGURE 8.13. Upper molars of *Hispanomys aragonensis* and *Hispanomys cf. aragonensis* from the Calatayud-Daroca Basin. *Hispanomys aragonensis*: Pedregueras 2A: A, left M1, RGM-567801; B, left M2, RGM-567984; C, inverted right M3, RGM-568041; D, left M1, RGM-567795; E, inverted right M2, RGM-567937; F, left M3, RGM-568063; G, inverted right M1, RGM-567763; H, inverted right M1, RGM-567756. *Hispanomys cf. aragonensis*: Nombrevilla 14: I, inverted right M1, NOM14-01; J, inverted right M2, NOM14-05; K, inverted right M3, NOM14-10; L, inverted right M3, NOM14-09. Nombrevilla 22: M, inverted right M2, NOM22-01; N, inverted right M3, NOM22-12. Scale bar equals 1 mm.




FIGURE 8.14. Lower molars of *Hispanomys aragonensis* and *Hispanomys* cf. *aragonensis* from the Calatayud-Daroca Basin. *Hispanomys aragonensis*: Pedregueras 2A: A, left m1, RGM-567859; B, inverted right m2, RGM-567884; C, left m3, RGM-568030; D, inverted right m1, RGM-567824; E, inverted right m2, RGM-567890; F, left m3, RGM-568031; G, inverted right m1, RGM-567851; H, left m2, RGM-567920; I, inverted right m3, RGM-567007; J, left m2, RGM-567293. *Hispanomys* cf. *aragonensis*: Nombrevilla 19: K, left m3, NOM19-01. Nombrevilla 14: L, left m1, NOM14-04; M, inverted right m3, NOM14-15; N, left m3, NOM14-14. Scale bar equals 1 mm.

	Species	Locality	Length						Width				
			Min	Mean	Max	SD	N		Min	Mean	Max	SD	N
M1	<i>Hispanomys aguirrei</i>	CARR	2.81	2.96	3.17	0.076	15		1.96	2.03	2.15	0.053	16
	<i>Hispanomys aguirrei</i>	ESC	2.62	3.08	3.54	0.146	77		1.64	1.97	2.27	0.108	79
	<i>Hispanomys cf. aguirrei</i>	NOM4	3.01	3.01	3.01		1		2.02	2.02	2.02		1
	<i>Hispanomys cf. aguirrei</i>	NOM3	2.88	2.96	3.03		2		2	2.04	2.08		2
	<i>Hispanomys aguirrei</i>	NOM2	2.76	3.08	3.40	0.135	56		1.88	2.05	2.15	0.072	56
	<i>Hispanomys aguirrei</i>	SOL	2.59	2.91	3.29	0.158	31		1.83	1.94	2.18	0.087	32
M2	<i>Hispanomys aguirrei</i>	CARR	2.08	2.24	2.40	0.083	18		1.53	1.80	2.01	0.117	18
	<i>Hispanomys aguirrei</i>	ESC	2.07	2.29	2.56	0.115	41		1.08	1.77	2.01	0.134	41
	<i>Hispanomys cf. aguirrei</i>	NOM4	2.23	2.23	2.23		1		1.82	1.82	1.82		1
	<i>Hispanomys cf. aguirrei</i>	NOM3	2.16	2.31	2.46		2		1.65	1.85	2.07		3
	<i>Hispanomys aguirrei</i>	NOM2	2.14	2.37	2.65	0.116	72		1.74	1.92	2.06	0.075	72
	<i>Hispanomys aguirrei</i>	SOL	1.97	2.17	2.42	0.106	38		1.69	1.88	2.17	0.121	41
M3	<i>Hispanomys aguirrei</i>	CARR	1.57	1.70	1.80	0.078	23		1.44	1.60	1.78	0.088	23
	<i>Hispanomys aguirrei</i>	ESC	1.51	1.73	1.95	0.096	35		1	1.57	1.84	0.086	35
	<i>Hispanomys aguirrei</i>	NOM2	1.48	1.70	1.86	0.076	60		1.38	1.60	1.73	0.070	61
	<i>Hispanomys aguirrei</i>	SOL	1.58	1.70	1.89	0.075	28		1.49	1.66	1.86	0.085	28
m1	<i>Hispanomys aguirrei</i>	CARR	2.19	2.53	2.70	0.124	20		1.48	1.71	1.94	0.117	21
	<i>Hispanomys aguirrei</i>	ESC	2.36	2.63	2.95	0.136	71		1.59	1.78	2	0.093	71
	<i>Hispanomys aguirrei</i>	NOM2	2.36	2.57	2.77	0.103	72		1.55	1.73	1.93	0.087	74
	<i>Hispanomys aguirrei</i>	SOL	2.22	2.48	2.82	0.143	40		1.53	1.66	1.96	0.101	43
m2	<i>Hispanomys aguirrei</i>	CARR	2.22	2.38	2.60	0.109	18		1.67	1.84	2.02	0.097	22
	<i>Hispanomys aguirrei</i>	ESC	2.38	2.55	2.72	0.132	43		1.83	1.94	2	0.134	43
	<i>Hispanomys cf. aguirrei</i>	NOM3	2.39	2.39	2.39		1		1.80	1.86	1.91	0.077	2
	<i>Hispanomys aguirrei</i>	NOM2	2.20	2.42	2.67	0.114	46		1.76	1.92	2.07	0.087	46
	<i>Hispanomys aguirrei</i>	SOL	2.17	2.36	2.55	0.098	35		1.58	1.85	2.13	0.117	42
m3	<i>Hispanomys aguirrei</i>	CARR	2.05	2.23	2.41	0.100	18		1.44	1.71	1.91	0.118	19
	<i>Hispanomys aguirrei</i>	ESC	2.05	2.20	2.42	0.098	28		1.91	1.65	1.30	0.111	28
	<i>Hispanomys cf. aguirrei</i>	NOM4	2.21	2.21	2.21		1		1.69	1.69	1.69		1
	<i>Hispanomys cf. aguirrei</i>	NOM3	2.20	2.00	2.26	0.042	2		1.67	1.73	1.79	0.084	2
	<i>Hispanomys aguirrei</i>	NOM2	2.00	2.23	2.50	0.112	76		1.55	1.77	1.91	0.084	76
	<i>Hispanomys aguirrei</i>	SOL	1.97	2.20	2.46	0.119	43		1.47	1.74	1.89	0.082	43

TABLE 8.4. Length and width of the upper and lower molars of *Hispanomys aguirrei* and *Hispanomys cf. aguirrei*. Units are given in mm. Abbreviations: Min, minimum; Max, maximum; SD, standar deviation; N, number of specimens.

	Species	Locality	Length						Width				
			Min	Mean	Max	SD	N		Min	Mean	Max	SD	N
M1	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM4	3.28	3.39	3.46	0.100	3		2.21	2.28	2.35	0.070	3
	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM3		3.38			1			2.21			1
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	3.18	3.58	4.06	0.190	37		2.11	2.29	2.56	0.090	38
	<i>Hispanomys</i> <i>lavocati</i>	SQ	3.38		3.69		2		2.12	2.21	2.31	0.082	4
	<i>Hispanomys</i> <i>lavocati</i>	HP	3.56	3.69	3.88		6		2.09	2.22	2.34		7
M2	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM4	2.52	2.58	2.63	0.080	2		2.08	2.11	2.14	0.040	2
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	2.40	2.64	2.88	0.090	53		2.12	2.22	2.45	0.070	54
	<i>Hispanomys</i> <i>lavocati</i>	SQ	2.54	2.71	2.92	0.146	8		2.02	2.21	2.34	0.101	8
	<i>Hispanomys</i> <i>lavocati</i>	HP	2.41	2.56	2.75		4		2.05	2.12	2.23		4
M3	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM3		1.88			1			1.83			1
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	1.65	1.96	2.14	0.090	75		1.71	1.85	2.00	0.080	75
	<i>Hispanomys</i> <i>lavocati</i>	SQ	1.93	2.04	2.14		2		1.73	1.85	1.96		2
	<i>Hispanomys</i> <i>lavocati</i>	HP	1.78	1.9	2.01		4		1.74	1.86	1.98		4
m1	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM4	2.87	2.96	3.04	0.120	2		1.87	1.89	1.90	0.020	2
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	2.47	2.85	3.05	0.130	56		1.8	1.95	2.12	0.090	60
	<i>Hispanomys</i> <i>lavocati</i>	SQ	2.02	2.72	3.21	0.391	6		1.74	1.85	2.07	0.135	6
	<i>Hispanomys</i> <i>lavocati</i>	HP	2.53	2.82	2.98		8		1.85	1.89	2.01		8
m2	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM4	2.60	2.67	2.73	0.090	2						0
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	2.55	2.74	2.9	0.090	50		2.11	2.22	2.35	0.060	50
	<i>Hispanomys</i> <i>lavocati</i>	SQ	2.52	2.66	2.79	0.126	6		1.8	2	2.79	0.165	6
	<i>Hispanomys</i> <i>lavocati</i>	HP	2.46	2.66	2.81		11		1.97	2.07	2.21		11
m3	<i>Hispanomys</i> cf. <i>lavocati</i>	NOM3	2.41	2.41	2.41		1			2.05			1
	<i>Hispanomys</i> <i>lavocati</i>	NOM2	2.26	2.44	2.64	0.090	71		1.93	2.04	2.17	0.060	71
	<i>Hispanomys</i> <i>lavocati</i>	SQ	2.45	2.53	2.65	0.073	5		1.82	1.87	1.94	0.044	6
	<i>Hispanomys</i> <i>lavocati</i>	HP	2.43	2.54	2.74		5		1.88	1.98	2		5

TABLE 8.5. Length and width of the upper and lower molars of *Hispanomys lavocati* and *Hispanomys* cf. *lavocati*. The measurements of the material from Sant Quirze and Hostalets de Pierola (Vallès-Penedès Basin) are from Freudenthal (1966) and Agustí (1980). Units are given in mm. Abbreviations: Min, minimum values; Max, maximum values; SD, standard deviation; N, number of specimens; SQ: Sant Quirze; HP: Hostalets de Pierola.

	Species	Locality	Length						Width				
			Min	Mean	Max	SD	N		Min	Mean	Max	SD	N
M1	<i>Hispanomys nombrevillae</i>	NOM1	2.65	2.82	3.00	0.110	10		1.83	1.91	2.02	0.060	10
	<i>Hispanomys nombrevillae</i>	NOM9	2.58	2.71	2.83	0.180	2		1.80	1.84	1.91	0.060	3
M2	<i>Hispanomys cf. nombrevillae</i>	CAÑ9		1.90			1			1.60			1
	<i>Hispanomys cf. nombrevillae</i>	CAÑ8		2.06			1			1.64			1
	<i>Hispanomys nombrevillae</i>	NOM1	1.69	2.02	2.12	0.100	16		1.57	1.70	1.85	0.080	14
	<i>Hispanomys nombrevillae</i>	NOM9	1.99	2.16	2.28	0.100	7		1.56	1.74	1.89	0.110	8
M3	<i>Hispanomys cf. nombrevillae</i>	CAÑ11		1.49			1			1.26			1
	<i>Hispanomys nombrevillae</i>	CAÑ10		1.52			1			1.43			1
	<i>Hispanomys cf. nombrevillae</i>	NOM1	1.25	1.47	1.77	0.150	12		1.27	1.44	1.64	0.100	12
	<i>Hispanomys cf. nombrevillae</i>	NOM10	1.50	1.51	1.52	0.010	2		1.48	1.50	1.51	0.020	2
	<i>Hispanomys nombrevillae</i>	NOM9	1.42	1.48	1.56	0.050	6		1.30	1.41	1.55	0.090	6
	<i>Hispanomys nombrevillae</i>	CAÑ10		2.16			1			1.45			1
m1	<i>Hispanomys cf. nombrevillae</i>	CAÑ9		2.60			1			1.59			1
	<i>Hispanomys nombrevillae</i>	NOM1	2.28	2.41	2.66	0.110	13		1.48	1.57	1.73	0.060	18
	<i>Hispanomys cf. nombrevillae</i>	NOM10		2.64			1			1.63			1
	<i>Hispanomys cf. nombrevillae</i>	NOM9	2.30	2.38	2.55	0.090	7		1.40	1.50	1.58	0.060	7
	<i>Hispanomys cf. nombrevillae</i>	CAÑ9		2.00			1		1.65	1.65	1.65		1
m2	<i>Hispanomys cf. nombrevillae</i>	CAÑ8		2.17			1		1.61	1.61	1.61		1
	<i>Hispanomys nombrevillae</i>	NOM1	1.98	2.21	2.38	0.100	14		1.64	1.73	1.78	0.040	15
	<i>Hispanomys nombrevillae</i>	NOM9	2.13	2.26	2.41	0.100	8		1.57	1.74	1.88	0.110	7
	<i>Hispanomys cf. nombrevillae</i>	CAÑ8		1.64			1			1.28			1
m3	<i>Hispanomys nombrevillae</i>	NOM1	1.61	1.88	2.21	0.130	21		1.41	1.54	1.74	0.080	22
	<i>Hispanomys nombrevillae</i>	NOM9	1.37	1.86	2.12	0.230	10		1.30	1.53	1.88	0.180	11

TABLE 8.6. Length and width of the upper and lower molars of *Hispanomys nombrevillae* and *Hispanomys cf. nombrevillae*. Units are given in mm. Abbreviations: Min, minimum; Max, maximum; SD, standar deviation; N, number of specimens.

	Species	Locality	Length						Width				
			Min	Mean	Max	SD	N		Min	Mean	Max	SD	N
M1	<i>Hispanomys aragonensis</i>	PED2C	2.69	2.82	3.02	0.113	13		1.73	1.82	1.93	0.061	14
	<i>Hispanomys aragonensis</i>	PED2A	2.79	3.01	3.25	0.130	34		1.75	1.90	2.01	0.066	35
	<i>Hispanomys aragonensis</i>	PED1A		2.60			1			1.67			1
	<i>Hispanomys cf. aragonensis</i>	NOM14		2.82			1		1.85	1.89	1.92	0.049	2
	<i>Hispanomys cf. aragonensis</i>	NOM13		1.79			1						0
M2	<i>Hispanomys aragonensis</i>	PED2C	1.99	2.17	2.31	0.083	25		1.58	1.70	1.86	0.074	25
	<i>Hispanomys aragonensis</i>	PED2A	1.98	2.26	2.46	0.112	44		1.56	1.74	1.91	0.082	46
	<i>Hispanomys aragonensis</i>	PED1A	2.09	2.12	2.14	0.035	2		1.63	1.65	1.67	0.018	4
	<i>Hispanomys cf. aragonensis</i>	NOM14					0			1.71			1
	<i>Hispanomys cf. aragonensis</i>	NOM22		2.18			1			1.58			1
M3	<i>Hispanomys aragonensis</i>	PED2C	1.39	1.64	1.81	0.105	37		1.31	1.49	1.66	0.086	38
	<i>Hispanomys aragonensis</i>	PED2A	1.44	1.66	2.06	0.123	38		1.37	1.53	1.77	0.091	38
	<i>Hispanomys aragonensis</i>	PED1A	1.54	1.59	1.63	0.064	2		1.55	1.56	1.57	0.014	2
	<i>Hispanomys cf. aragonensis</i>	NOM14	1.49	1.53	1.58	0.064	2		1.46	1.47	1.47	0.007	2
	<i>Hispanomys cf. aragonensis</i>	NOM22		1.53			1			1.40			1
m1	<i>Hispanomys aragonensis</i>	PED2C	2.09	2.45	2.76	0.120	43		1.36	1.53	1.66	0.07	42
	<i>Hispanomys aragonensis</i>	PED2A	2.30	2.51	2.72	0.098	47		1.40	1.58	1.70	0.069	48
	<i>Hispanomys aragonensis</i>	PED1A		2.35			1			1.54			1
	<i>Hispanomys cf. aragonensis</i>	NOM13		2.41			1			1.49			1
m2	<i>Hispanomys aragonensis</i>	PED2C	2.15	2.29	2.43	0.070	21		1.61	1.76	1.92	0.076	21
	<i>Hispanomys aragonensis</i>	PED2A	2.22	2.37	2.50	0.081	31		1.58	1.78	1.92	0.072	32
m3	<i>Hispanomys cf. aragonensis</i>	NOM19		2.15			1			1.69			1
	<i>Hispanomys aragonensis</i>	PED2C	1.92	2.13	2.30	0.100	35		1.40	1.60	1.76	0.087	34
	<i>Hispanomys aragonensis</i>	PED2A	2.04	2.16	2.32	0.087	24		1.50	1.63	1.75	0.057	23
	<i>Hispanomys aragonensis</i>	PED1A	1.86	2.04	2.22	0.255	2		1.40	1.56	1.72	0.226	2
	<i>Hispanomys cf. aragonensis</i>	NOM14	1.95	1.97	1.99	0.028	2		1.58	1.59	1.61	0.015	3

TABLE 8.7. Length and width of the upper and lower molars of *Hispanomys aragonensis* and *Hispanomys cf. aragonensis*. Units are given in mm. Abbreviations: Min, minimum; Max, maximum; SD, standar deviation; N, number of specimens.

<i>H. aguirrei</i>				
	Levenne's test		ANOVA	
	Length	Width	Length	Width
M1	0.003	0.031	0	0
M2	0.050	0.004	0	0
M3	0.551	0.070	0.497	0.018
m1	0.178	0.368	0.001	0
m2	0.370	0.652	0.036	0.003
m3	0.491	0.550	0.965	0.006

(a)

			Localities			
			CARR	ESC	NOM2	SOL
Localities				Width (p value)		
M1	CARR	Length (p value)		0.040	0.789	0.001
	ESC		0		0.004	0.917
	NOM2		0.001	0.719		0
	SOL		0.473	0	0	
M2	CARR	Length (p value)		0.938	0.002	0.100
	ESC		0.704		0	0.032
	NOM2		0	0.001		0.144
	SOL		0.073	0	0	
M3	CARR	Length (p value)		0.834	1	0.220
	ESC		0.916		0.604	0.004
	NOM2		1	0.598		0.078
	SOL		1	0.847	1,000	
m1	CARR	Length (p value)		0.951	0.963	0.313
	ESC		0.003		1	0.002
	NOM2		0.883	0.001		0.002
	SOL		0.465	0	0.002	
m2	CARR	Length (p value)		0.953	0.008	0.979
	ESC		0		0.219	0.995
	NOM2		0.567	0		0.009
	SOL		0.887	0	0.049	
m3	CARR	Length (p value)		0.223	0.078	0.660
	ESC		1		0	0
	NOM2		1	1		0.690
	SOL		0.974	0.982	0.802	

(b)

TABLE 8.8. Results for the statistical test for the length and width of the molars of *Hispanomys aguirrei* (SOL, NOM2, ESC and CARR). (a). Results for the Levenne's and ANOVA tests. Numbers in red: Significant differences ($p < 0.05$), homogeneous variances. (b). Results of the Post-Hoc tests. In these case we have used the Games-Howel contrasts given that the non-homogeneity of the variances. Numbers in black: No significant Differences; blue: Significant differences ($p < 0.05$); red: highly significant differences ($p < 0.01$).

NOM1 Vs NOM9								
	Levenne's test				Student's T			
	Length		Width		Length		Width	
M1	0.404	H	0.511	H	0.249	NS	0.121	NS
M2	0.361	H	0.605	H	0.001	HS	0.162	NS
M3	0.080	H	0.848	H	0.847	NS	0.600	NS
m1	0.533	H	0.788	H	0.622	NS	0.017	S
m2	0.934	H	0.040	NH	0.330	NS	0.780	NS
m3	0.048	NH	0.005	NH	0.748	NS	0.811	NS

(a) TABLE 8.9. Results for the Levenne's and Student's t test for the length and width of the molars of (a) *Hispanomys nombrevillae* (NOM1 and NOM9) (b) for *H. nombrevillae* and *H. aragonensis* (NOM1 and PED2C). Abbreviations: H: No Significant, equal variances; NH: unequal variances; NS: No significant Differences; S: Significant differences ($p < 0.05$); HS: highly significant differences ($p < 0.01$).

<i>H. nombrevillae</i> Vs <i>H. aragonensis</i>								
	Levenne's test				Student's T			
	Length		Width		Length		Width	
M1	0.778	H	0.849	H	0.769	NS	0.001	HS
M2	0.973	H	0.719	H	0	HS	0.886	NS
M3	0.252	H	0.790	H	0	HS	0.097	NS
m1	0.693	H	0.304	H	0.346	NS	0.031	S
m2	0.234	H	0.172	H	0.014	S	0.202	NS
m3	0.138	H	0.587	H	0	HS	0.014	S

(b)

<i>H. aragonensis</i>								
	Levenne's test				Student's T			
	Length		Width		Length		Width	
M1	0.404	H	0.678	H	0	HS	0.001	HS
M2	0.140	H	0.426	H	0	HS	0.018	S
M3	0.833	H	0.577	H	0.320	NS	0.022	S
m1	0.419	H	0.918	H	0.011	S	0.005	HS
m2	0.400	H	0.948	H	0.001	HS	0.281	NS
m3	0.850	H	0.010	NH	0.138	NS	0.101	NS*

TABLE 8.10. Results of the Levenne's and Student's t tests for the length and width of the molars of *Hispanomys aragonensis* from Pedregueras 2A and Pedregueras 2C. Abbreviations: H: No Significant, equal variances; NH: unequal variances; NS: No significant Differences; *result for unequal variances; S: Significant differences ($p < 0.05$); HS: highly significant differences ($p < 0.01$).

Reference	Genus	Subgenus
Freudenthal, 1966	<i>Cricetodon</i>	—
Mein and Freudenthal, 1971a	<i>Cricetodon</i>	<i>Pseudoruscinomys</i>
Van de Weerd, 1976	<i>Hispanomys</i>	—
Agustí, 1977a	<i>Cricetodon</i>	—
Agustí, 1977b	<i>Hispanomys</i>	—
Agustí, 1978	<i>Cricetodon</i>	—
Agustí and Gilbert, 1979	<i>Cricetodon</i>	—
Agustí, 1980	<i>Cricetodon</i>	—
Sesé, 1980	<i>Hispanomys</i>	—
Agustí, 1981	<i>Cricetodon</i>	<i>Pararuscinomys</i>
Agustí, 1982	<i>Cricetodon</i>	<i>Pararuscinomys</i>
Aguilar, 1982	<i>Hispanomys</i>	—
Agustí, 1997	<i>Cricetodon</i>	—
Agustí, 2005	<i>Cricetodon</i>	<i>Pararuscinomys</i>
Casanovas-Vilar et al., 2006	<i>Cricetodon</i>	—
Casanovas-Vilar and Agustí, 2007	<i>Hispanomys</i>	—
López-Guerrero et al., 2008	<i>Hispanomys</i>	—
López-Antoñanzas et al., 2010	<i>Hispanomys</i>	—
López-Guerrero et al., 2014	<i>Hispanomys</i>	—

TABLE 8.11. List of the genera and subgenera where *Hispanomys lavocati* has been ascribed to.

Basal for the tribe	Derived for the tribe	Basal for the genus <i>Hispanomys</i>
Poor development/absence of the ectolophs	Complete Ectolophs	Presence of labial anteroloph
Cingular formations well developed	Poor development of the cinguls/ids	Moderately hypsodonty
Metalophulid II or I+II	Absence of metalophulid II without I	Presence of both metalophulids in the same m1
Anterocone slightly split	Low presence/absence of M3 posteroloph	Small size
Two rooted m2	Three rooted m2	

TABL 8.12. Features present in *Hispanomys nombrevillae* classified by its derivate/primitive character according to the trends within tribe Cricetodontini (left and middle columns) or within genus *Hispanomys* (right column).

FINAL REMARKS & CONCLUSIONS

aragonensis. Both hypotheses seem plausible, but they will be fully studied and discussed in chapter 10.

The fossils from the Calatayud-Daroca Basin ascribed to *Hispanomys* have been assigned to four different species. Two of them, *H. aguirrei* and *H. lavocati* are recovered from the late Aragonian, another, *H. nombrevillae*, is present in the late Aragonian and the early Vallesian, and the last one, *H. aragonensis*, is recovered from the early Vallesian.

These species can be clustered in two groups. The first one includes *H. aguirrei* and *H. lavocati* which display a combination of basal and derived features. This condition complicated their taxonomical study and their generic assignation was very difficult. After present work, we conclude that their morphology corresponds to a basal *Hispanomys*. The morphological observations and comparisons made here cannot confirm their phylogenetic relationships in the context of the Calatayud-Daroca Basin. There are several possibilities equally plausible for both species. Moreover, their origin, especially in the case of *H. lavocati* is still ambiguous. Therefore, these taxa need a supplemental cladistic analysis to establish the phylogeny and to better understand their evolutionary history.

The other group is formed by *H. nombrevillae* and *H. aragonensis*. These species are less controversial than the other two because their features are characteristic of *Hispanomys*. The evolutionary stage of each one is easily recognizable; *Hispanomys nombrevillae* is a basal representative of the genus, whereas *H. aragonensis* has more derived features. Both species are clearly originated in southwestern Europe (Spain and France), but both clusters do not seem to be related. *Hispanomys aguirrei* and *H. lavocati* may have a different origin than *H. nombrevillae* and *H. aragonensis*. These questions will be treated in chapter 10.

Hispanomys aguirrei and *H. lavocati* (late Aragonian) are bigger than *H. nombrevillae* and *H. aragonensis* from the Vallesian. This follows the trend indicated by Agustí (1982) towards size decreasing among the species of *Hispanomys* for the Calatayud-Daroca Basin. Besides, there is a variation in size within the assemblages of *H. aguirrei*: the older fossils are smaller than the younger ones. This could indicate a trend towards intraspecific size increase, apart from the abovementioned interspecific trend.

LITERATURE CITED

- Aguilar, J. P. 1982. Contributions à l'étude des Micromammifères du gisement Miocène supérieur de Montredon (Hérault) 2. les rongeurs. *Palaeovertebrata* 12:75–140.
- Aguilar, J. P., M. Calvet, and J. Michaux. 1994. Les rongeurs de Castelnou 6 (Pyrénées-orientales, France) et les corrélations entre faunes ibériques et françaises au Miocène moyen. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 192:109–131.
- Aguilar, J. P., V. Lazzari, J. Michaux, M. Sabatier, and M. Calvet. 2007. Lo Fournas 16-M (Miocène supérieur) et Lo Fournas 16-P (Pliocène moyen), nouvelles localités karstiques à Baixas, Sud de la France. Partie I Description et implications géodynamiques. *Géologie de la France* 1:55–62.
- Agustí, J. 1977. Contribución al conocimiento del género *Hispanomys* (Cricetidae, Rod., Mam.) En la cuenca Miocénica del Vallés-Penedés. *Butlletí informatiu Institut provincial de paleontologia de Sabadell. Boletín informativo*, (julio-diciembre, 1977) año IX, 2:30–32.
- Agustí, J. 1980. La asociación de *Hispanomys* y *Cricetodon* (Rodentia, Mammalia) en el Mioceno superior del Vallés-Penedés (Cataluña, España). *Acta Geologica hispanica* 15:51–60.
- Agustí, J. 1982. Tendencias evolutivas de la línea *Cricetodon-Ruscinomys* (Rodentia, Mammalia) en la Península Ibérica. *Acta Geologica hispanica* 17:103–111.
- Agustí, J. 1986. Nouvelles espèces de Cricétidés vicariantes dans le Turolien moyen de Fortuna (Province de Murcia, Espagne). *Geobios* 19:5–11.
- Agustí, J., and J. Gibert. 1979. Nuevos datos sobre los Roedores e Insectívoros (Mammalia) del mioceno superior del Vallés (Cataluña, España). *Paleontología i Evolució* 14:17–20.
- Agustí, J., I. Casanovas-Vilar, and M. Furió. 2005. Rodents, insectivores and chiropterans (Mammalia) from the late Aragonian of Can Missert (middle Miocene, Vallés-Penedés Basin, Spain). *Geobios* 38:575–583.
- Alba, D. M., I. Casanovas-Vilar, J. M. Robles, and S. Moyà-Solà. 2011. Parada 3. El Aragoniense superior y la transición con el Vallesiense: Can Mata y la exposición paleontológica de els Hostalets de Pierola; pp. 95–109 in M. Pérez de los Ríos, J. Marigó, R. Minwer-Barakat, A. Bolet, D. DeMiguel (eds.), *Guía de Campo de las XXVII Jornadas de la Sociedad Es-*

- pañola de Paleontología. Paleontologia i Evolució, memoria especial número 6. Sabadell.
- Álvarez-Sierra, M. A. 1983. Paleontología y bioestratigrafía del Mioceno superior del sector central de la Cuenca del Duero. Estudio de los micromamíferos de la serie de Torremormojón (Palencia). Memoria de grado de licenciatura. Universidad Complutense de Madrid, Madrid, 188 pp.
- Álvarez Sierra, M. A., J. Civis, A. Corrochano, R. Daams, C. J. Dabrio, E. García, A. González, N. López-Martínez, R. Mediavilla, R. Rivas Carballo, and M. F. Valle. 1990. Un estratotipo del límite Aragoniense-Vallesiense (Mioceno medio-Mioceno superior) en la sección de Torremormojón (Cuenca del Duero, Provincia de Palencia). *Studia Geologica Salmanticensia, Actas de Paleontología*:57–64.
- Álvarez Sierra, M. A., J. P. Calvo, J. Morales, A. Alonso-Zarza, B. Azanza, I. García Paredes, M. Hernández Fernández, A. J. Van der Meulen, P. Peláez-Campomanes, V. Quiralte, M. J. Salesa, I. M. Sánchez, and D. Soria. 2003. El tránsito Aragoniense-Vallesiense en el área de Daroca-Nombrevilla (Zaragoza, España). *Coloquios de Paleontología Volumen extraordinario* 1:25–33.
- Bowdich, T. E. 1821. *An Analysis of the Natural Classification of Mammals for the Use of Students and Travelers*. J. Smith, Paris, 115 pp.
- Casanovas-Vilar, I., and J. Agustí. 2007. Ecogeographical stability and climate forcing in the Late Miocene (Vallesian) rodent record of Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 248:169–189.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.
- Daams, R., L. Alcalá, M. A. Álvarez-Sierra, B. Azanza, J. A. Van Dam, A. J. Van der Meulen, J. Morales, M. Nieto, P. Peláez-Campomanes, and D. Soria. 1998. A stratigraphical framework for Miocene (MN4-MN13) continental sediments of Central Spain. *Comptes rendus de l'Académie des sciences, Séries IIA* 327:625–631.
- De Bruijn, H., V. Fahlbusch, G. Saraç, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Critetodon* with a Discussion of the Evolutionary History of the Cricetodontini. *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen-Biological Chemical Geological Physical and Medical Sciences* 96:151–216.

- De Bruijn, H., R. Daams, V. Fahlbusch, G. Daxner-Höck, L. Ginsburg, P. Mein, and J. Morales. 1992. Report of the RCMNS working group on fossil mammals, Reisensburg 1990. *Newsletters of Stratigraphy* 26:65–118.
- Fischer, G. 1817. *Adversaria zoologica*, Mémoires de la Société Impériale des Naturalistes de Moscou 1917:357–472.
- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mittelspaniens und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) Ia. Genera *Cricetodon* and *Ruscinomys* (Rodentia). Koninklijke Nederlandse Akademie Van Wetenschappen-Proceedings Series B-Physical Sciences 69:298–317.
- Freudenthal, M., and Sondaar, P.Y. 1964. Les faunes à Hipparion des environs de Daroca (Espagne) et leur valeur pour la stratigraphie du Néogène de l'Europe. *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen*, sér. B 67:473–490.
- García-Alix, A., Minwer-Barakat, R., Martín-Suárez, E., Freudenthal, M., 2008. Muridae from the Mio-Pliocene boundary in the Granada Basin (southern Spain). *Biostratigraphic and phylogenetic implications*. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 248, 183–215.
- García-Moreno, E. 1987. *Roedores y Lagomorphos del Mioceno de la Zona central de la Cuenca del Duero. Sistemática, Bioestratigrafía y Paleoeología*. Ph.D. dissertation, Universidad Complutense, Madrid, Madrid, 258 pp.
- García-Paredes, I., P. Peláez-Campomanes, and M. A. Álvarez-Sierra. 2009. Gliridae (Rodentia, Mammalia) with a simple dental pattern: a new genus and new species from the European Early and Middle Miocene. *Zoological Journal of the Linnean Society* 157:622–652.
- Gómez Cano, A. R., M. Hernández Fernández, and M. A. Álvarez-Sierra. 2011. Biogeographic provincialism in rodent faunas from the Iberocitanian Region (southwestern Europe) generates severe diachrony within the Mammalian Neogene (MN) biochronologic scale during the Late Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 307:193–204.
- Hartenberger, J. L. 1965. Les Cricetidae (Rod.) de Can Llobateres (Néogène d'Espagne). *Bulletin De La Societe Geologique De France* 7:487–498.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőltárkány-Fenémet (Northern Hungary). *Fragmenta Palaeontologica Hungarica* 24-25:15–24.

- Lacomba Andueza, J. I. 1988. Rodents and lagomorphs from a Lower Vallesian fissure filling near Molina de Aragon (Prov. de Guadalajara, Spain). *Scripta Geologica Special Issue* 1:19–38.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d'une récapitulation des diverses espèces d'animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d'autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. J. A. Portes, Auch, 41 pp.
- López-Antoñanzas, R., and P. Mein. 2009. First detailed description of *Hispanomys bijugatus* Mein and Freudenthal, 1971 (Rodentia, Cricetodontinae) from the Upper Aragonian of La Grive-Saint Alban (France): Biostratigraphical implications. *Geobios* 42:783–796.
- López-Antoñanzas, R., and P. Mein. 2011. First detailed description of *Hispanomys decedens* (Rodentia) from the Middle Miocene of La Grive-Saint Alban (France). *Swiss Journal of Geosciences* 104:345–353.
- López-Antoñanzas, R., P. Peláez-Campomanes, M. A. Álvarez-Sierra, and I. García-Paredes. 2010. New species of *Hispanomys* (Rodentia, Cricetodontinae) from the Upper Miocene of Batallones (Madrid, Spain). *Zoological Journal of the Linnean Society* 160:725–747.
- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89:573–592.
- López-Guerrero, P., M. Álvarez-Sierra, I. García-Paredes, R. López-Antoñanzas, and A. Oliver. 2009. Cricetodontini (Rodentia, Mammalia) from the Upper Aragonian and Lower Vallesian of the Toril-Nombrevilla Section (Middle and Upper Miocene, Calatayud-Daroca Basin, Zaragoza, Spain). *Journal of Vertebrate Paleontology* 29:161A–161A.
- López-Guerrero, P., M. A. Álvarez-Sierra, R. López-Antoñanzas, A. Oliver Pérez, and P. Peláez-Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). *Palaeontologica Nova* 8:259–271.
- Lopez-Guerrero, P., I. García-Paredes, L. W. V. Ostende, J. A. Van Dam, M. A. Álvarez-Sierra, V. Hernández-Ballarín, A. J. Van der Meulen, A. Oliver, and P. Peláez-Campomanes. 2011. Cañada: a new micromammal succession from the lower Vallesian and Turolian of the


- Daroca area (Calatayud-Montalbán basin, Spain). *Estudios Geológicos* 67:443–453.
- López-Martínez, N., C. Sesé Benito, and J. L. Sanz García. 1977. La microfauna (Rodentia, Insectivora, Lagomorpha y Reptilia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno/Cuaternario* 8:47–73.
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam.Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.
- Schaub, S. 1925. Die Hamsterartige Nagetiere des Tertiärs und ihre lebenden Verwandten. *Abhandlungen des Schweizerischen paläontologische Gesellschaft = Mémoires de la Société paléontologique suisse* 45:1–114.
- Sesé Benito, C. 1977. Los cricétidos (Rodentia, Mammalia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno Cuaternario* 8:127–180.
- Sesé Benito, C. 1980. Mamíferos del Mioceno medio de Escobosa de Calatañazor (Soria). Ph. D dissertation. Universidad Complutense de Madrid, Madrid, 406 pp.
- Simpson, G. C. 1945. The principles of classification and a classification of mammals. *Bulletin of the American Museum of Natural History* 85:1–350.
- Stehlin, H. G., and S. Schaub. 1951. Die Trigonodontie der simplicidentaten Nager. *Schweizerische paläontologische Abhandlungen* 67:1–385.
- Van Dam, J., W. Krijgsman, H. A. Abels, M. A. Álvarez-Sierra, I. García-Paredes, P. López-Guerrero, P. Peláez-Campomanes, and D. Ventura. In press. Updated chronology for middle to late Miocene mammal sites of the Daroca area (Calatayud-Montalbán Basin), Spain. DOI: <http://dx.doi.org/10.1016/j.geobios.2014.07.002>
- Van Dam, J. A., H. Abdul-Aziz, M. A. Álvarez Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. Lourens, P. Mein, A. J. Van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodents fauna of the Mio-Pliocene continental sediments of the Teruel-Alfambra region, Spain. *Utrecht micropaleontological bulletins. Special publication. Utrecht* 2:185.

Appendix 8.1

Appendix 8.1.

Morphological variability of *Hispanomys aguirrei* (Carrilanga, Solera and Nombrevilla 2) and *Hispanomys* cf. *aguirrei* (Paje 1, Paje 2, Nombrevilla 4, and Nombrevilla 3) in the Calatayud-Daroca Basin; number of specimens that have each state of character and the percentage (in brackets) over the total number of specimens (N).

M1





	Open	Low Cingul	Cingul	N
Anterosinus				
Carrilanga	2 (10.53%)	7 (36.84%)	10 (52.63%)	19
Nombrevilla 4	—	—	2	2
Nombrevilla 3	—	—	1	1
Nombrevilla 2	3 (5.26%)	25 (43.86 %)	29 (50.88%)	57
Solera	2 (5.41%)	14 (37.84%)	21 (56.76%)	37
Paje 2	—	2	1	3


	Absent	Posterior Anterocone	Anterior Paracone	Double	N
Anterior ectoloph					
Carrilanga	2 (10.53%)	2 (10.53%)	10 (52.63%)	5 (26.32%)	19
Nombrevilla 4	—	—	3	—	3
Nombrevilla 3	—	—	—	1	1
Nombrevilla 2	3 (4.62%)	52 (80%)	—	10 (15.38%)	65
Solera	2 (5%)	35 (87.5%)	—	3 (7.5%)	40
Paje 2	—	6	—	—	6
Paje 1	1	—	—	—	1

	Not Complete	Complete	N
Anterior ectoloph			
Carrilanga	6 (35.29%)	11 (64.71%)	17
Nombrevilla 4	—	2	2
Nombrevilla 3	1	—	1
Nombrevilla 2	33 (52.38%)	30 (47.62%)	63
Solera	24 (63.16%)	14 (36.84%)	38
Paje 2	4	1	5







M1






	Absent	Simple	Double	N
				
Posterior ectoloph				
Carrilanga	—	4 (20%)	16 (80%)	20
Nombrevilla 4	—	—	3	3
Nombrevilla 3	—	—	1	1
Nombrevilla 2	—	43 (70.49%)	18 (29.51%)	61
Solera	1 (2.7%)	34 (91.89%)	2 (5.41%)	37
Paje 2	—	4	—	4
Paje 1	—	—	1	1





	Not Complete	Complete	N
			
Posterior ectoloph			
Carrilanga	2 (10%)	18 (90%)	20
Nombrevilla 4	—	3	3
Nombrevilla 3	—	1	1
Nombrevilla 2	26 (44.83%)	32 (55.17%)	58
Solera	38 (95%)	2 (5%)	40
Paje 2	4	—	4
Paje 1	1	—	1

	Absent	Enlargement	Short	Medium	N
					
Mesoloph					
Carrilanga	9 (50%)	8 (44.44%)	1 (5.56%)	—	18
Nombrevilla 4	2	—	—	—	2
Nombrevilla 3	1	—	—	—	1
Nombrevilla 2	45 (71.43%)	10 (15.87%)	5 (7.94%)	3 (4.76%)	63
Solera	17 (42.50%)	5 (12.50%)	18 (45%)	—	40
Paje 2	2	1	—	1	4


M1






	Open	Cingul	Style	Anteroloph	Low cingul	
						
Protosinus						N
Carrilanga	—	7 (41.18%)	2 (11.76%)	—	8 (47.06%)	17
Nombrevilla 4	—	1	1	—	—	2
Nombrevilla 3	1	—	—	—	—	1
Nombrevilla 2	2 (3.77%)	17 (32.08%)	14 (26.42%)	20 (37.73%)	—	53
Solera	—	6 (15.79%)	19 (50%)	13 (34.21%)	—	38
Paje 2	1	1	1	1	—	4
Paje 1	—	1	—	—	—	1




	Open	Cingul	Style	Two crests	
					
Sinus					N
Carrilanga	13 (72.22%)	1 (5.56%)	4 (22.22%)	—	18
Nombrevilla 4	1	—	—	1	2
Nombrevilla 3	—	—	—	1	1
Nombrevilla 2	42 (67.74%)	8 (12.9%)	—	12 (18.35%)	62
Solera	19 (46.34%)	7 (17.07%)	3 (7.32%)	12 (29.27%)	41
Paje 2	3	—	—	1	4

	Open	Cingul	Low Cingul	
				
Mesosinus				N
Carrilanga	—	9 (47.37%)	10 (52.63%)	19
Nombrevilla 4	—	2	—	2
Nombrevilla 3	—	1	—	1
Nombrevilla 2	—	46 (80.70%)	11 (19.30%)	57
Solera	2 (4.88%)	31 (75.61%)	8 (19.51%)	41
Paje 2	1	1	1	3
Paje 1	—	—	1	1

M2



	Absent	Posterior Paracone	Anterior Anterocone	Double	N
 Anterior ectoloph					
Carrilanga	5 (23.81%)	—	15 (71.43%)	1 (4.76%)	21
Nombrevilla 3	2 (66.67%)	1 (33.33%)	—	—	3
Nombrevilla 2	50 (64.10%)	28 (35.90%)	—	—	78
Solera	8 (19.51%)	—	32 (78.05%)	1 (2.44%)	41
Paje 2	1	—	—	—	1
Paje 1	2	—	—	—	2

	Not Complete	Complete	N
 Anterior ectoloph			
Carrilanga	—	16	16
Nombrevilla 3	1	—	1
Nombrevilla 2	8 (28.57%)	20 (71.43%)	28
Solera	2 (6.06%)	31 (93.94%)	33
Paje 2	—	—	—
Paje 1	—	—	—

M2

	Absent	Enlargement	Short	
Mesoloph				N
Carrilanga	17 (80.95%)	4 (19.05%)	—	21
Nombrevilla 3	3	—	—	3
Nombrevilla 2	76 (90.48%)	4 (4.76%)	4 (4.76%)	84
Solera	27 (62.79%)	12 (27.91%)	4 (9.30%)	43
Paje 2	—	—	1	1
Paje 1	1	4	—	5

	Simple	Double	
Posterior ectoloph			N
Carrilanga	12 (57.14%)	9 (42.86%)	21
Nombrevilla 3	2 (66.67%)	1 (33.33%)	3
Nombrevilla 2	55 (69.62%)	24 (30.38%)	79
Solera	40 (88.89%)	5 (11.11%)	45
Paje 2	1	—	1
Paje 1	2	—	2

	Not Complete	Complete	
Posterior ectoloph			N
Carrilanga	4 (19.05%)	17 (80.95%)	21
Nombrevilla 3	1	2	3
Nombrevilla 2	62 (78.48%)	17 (21.52%)	79
Solera	40 (88.89%)	5 (11.11%)	45
Paje 2	1	—	1
Paje 1	2	—	2

M2

Absent

Present



Entomesoloph

N

Carrilanga	18 (94.74%)	1 (5.26%)	19
Nombrevilla 3	3	—	3
Nombrevilla 2	79 (98.75%)	1 (1.25%)	80
Solera	40 (93.02%)	3 (6.98%)	43
Paje 2	1	—	1
Paje 1	2	—	2



Mesosinus

Open

Low Cingul

Cingul

Two Crests

N

Carrilanga	3 (14.29%)	7 (33.33%)	11 (52.38%)	—	21
Nombrevilla 3	—	—	2	—	2
Nombrevilla 2	2 (2.50%)	9 (11.25%)	69 (86.25%)	—	80
Solera	5 (11.36%)	21 (47.73%)	16 (36.36%)	2 (4.55%)	44
Paje 2	—	—	—	—	—
Paje 1	1	—	—	—	1



Sinus

Open

Cingul


Two Crests




Style




N




Carrilanga	17 (94.44%)	1 (5.56%)	—	—	18
Nombrevilla 3	3	—	—	—	3
Nombrevilla 2	48 (71.64%)	12 (17.91%)	2 (2.99%)	5 (7.46%)	67
Solera	33 (78.57%)	5 (11.90%)	—	4 (9.52%)	42
Paje 2	1	—	—	—	1
Paje 1	2	—	—	—	2

M3











	Absent	Present	N
 Anterior Ectoloph			
Carrilanga	21 (95.45%)	1 (4.55%)	22
Nombrevilla 3	—	—	—
Nombrevilla 2	74 (98.67%)	1 (1.33%)	75
Solera	29	—	29
Paje 2	4	—	4






	Not Complete	Complete	N
 Anterior Ectoloph			
Carrilanga	1	—	1
Nombrevilla 3	—	—	—
Nombrevilla 2	—	1	1
Solera	—	—	—
Paje 2	—	—	—

	Absent	Present	N
 Lingual Aneroloph			
Carrilanga	5 (27.78%)	13 (72.22%)	18
Nombrevilla 3	1	—	1
Nombrevilla 2	66 (92.96%)	5 (7.04%)	71
Solera	6 (22.22%)	21 (77.78%)	27
Paje 2	—	4	4




M3





	Absent	Short	Medium	Large	N
					
Mesoloph					
Carrilanga	17 (89.47%)	2 (10.53%)	—	—	19
Nombrevilla 3	—	—	—	—	—
Nombrevilla 2	73	—	—	—	73
Solera	16 (57.14%)	5 (17.86%)	3 (10.71%)	4 (14.29%)	28
Paje 2	3	1	—	—	4




	Absent	Present	N
			
Neo-entoloph			
Carrilanga	19 (82.61%)	4 (17.39%)	23
Nombrevilla 3	1	—	1
Nombrevilla 2	83	—	83
Solera	26 (96.30%)	1 (3.7%)	27
Paje 2	—	2	2

	Absent	Simple Paracone	Simple Protolophule	Double	N
					
Posterior Ectoloph					
Carrilanga	—	13 (59.09%)	2 (9.09%)	7 (31.82%)	22
Nombrevilla 3	—	—	—	1	1
Nombrevilla 2	—	38 (54.29%)	1 (1.43%)	31 (44.28%)	70
Solera	4 (11.77%)	16 (47.06%)	3 (8.82%)	11 (32.35%)	34
Paje 2	—	—	—	4	4


M3





	Not Complete	Complete	N
 Posterior Ectoloph			
Carrilanga	4 (18.18%)	18 (81.82%)	22
Nombrevilla 3	—	1	1
Nombrevilla 2	21 (30.88%)	47 (69.12%)	68
Solera	16 (61.54%)	10 (38.46%)	26
Paje 2	4	—	4





	Absent	Present	Well Developed	N
 Posteroloph				
Carrilanga	5 (25%)	—	15 (75%)	20
Nombrevilla 3	1	—	—	1
Nombrevilla 2	51 (71.83%)	12 (16.90%)	8 (11.27%)	71
Solera	13 (48.15%)	8 (29.63%)	6 (22.22%)	27
Paje 2	—	—	—	—

	Absent	Present	N
 Enamel-coated valley			
Carrilanga	18	5	23
Nombrevilla 3	1	—	1
Nombrevilla 2	83	—	83
Solera	23	5	28
Paje 2	4	—	4




m1











	Absent	Short	Long	N
 Labial Anterolophid				
Carrilanga	—	5 (21.74%)	18 (78.26%)	23
Nombrevilla 2	1 (1.79%)	33 (58.93%)	22 (39.28%)	56
Solera	—	26 (57.78%)	19 (42.22%)	45
Paje 2	—	3 (60%)	2 (40%)	5
Paje 1	—	1	1	2

	Absent	Enlargement	Short	N
 Mesolophid				
Carrilanga	12 (54.55%)	9 (40.90%)	1 (4.55%)	22
Nombrevilla 2	80 (93.02%)	3 (3.49%)	3 (3.49%)	86
Solera	25 (54.35%)	15 (32.61%)	6 (13.04%)	46
Paje 2	3	1	2	6
Paje 1	—	—	2	2



m1

	Absent	Present	
			
Ectomesolophid			N
Carrilanga	23	—	23
Nombrevilla 2	91	—	91
Solera	38	5	43
Paje 2	4	1	5
Paje 1	1	1	2


	Absent	Metalophulid I	Metalophulid II	Metalophulid I+II	
					
Metalophulid					N
Carrilanga	—	17 (77.27%)	—	5 (22.73%)	22
Nombrevilla 2	3 (3.57%)	47 (55.95%)	—	34 (40.48%)	84
Solera	1 (2.08%)	20 (41.67%)	4 (8.33%)	23 (47.92%)	48
Paje 2	2	2	—	2	6
Paje 1	—	3	—	—	3

	Open	Low Cingulid	Cingulid	Stylid	
					
Mesosinusid					N
Carrilanga	14 (58.33%)	8 (33.33%)	1 (4.17%)	1 (4.17%)	24
Nombrevilla 2	59 (75.64%)	19 (24.36%)	—	—	78
Solera	32 (69.57%)	13 (28.26%)	—	1 (2.17%)	46
Paje 2	5	1	—	—	6
Paje 1	1	1	—	—	2


m1




m2




Posterior root




Undivided




Partially Divided




Completely Divided




	N
Carrilanga	2 (14.29%)
Nombrevilla 3	—
Nombrevilla 2	1 (3.45%)
Solera	3 (18.75%)
Paje 2	—
Paje 1	—



Labial anterolophid




Short




Long


	N
Carrilanga	9 (42.86%)
Nombrevilla 3	—
Nombrevilla 2	—
Solera	3 (6.82%)
Paje 2	1
Paje 1	—



Ectomesolophid








Absent












Present

	N
Carrilanga	23
Nombrevilla 3	3
Nombrevilla 2	73
Solera	49
Paje 2	4
Paje 1	1


m2

		Open 	Cingulid 	Two Crests 	Stylid 	N
Sinusid						
Carrilanga		3 (12%)	20 (80%)	1 (4%)	1 (4%)	25
Nombrevilla 3		1	—	1	—	2
Nombrevilla 2		2 (3.91%)	43 (84.31%)	6 (11.76%)	—	51
Solera		3 (6.25%)	32 (66.67%)	7 (14.58%)	6 (12.50%)	48
Paje 2		1	3	—	—	4
Paje 1		—	1	—	—	1

		Open 	Low Cingulid 	N
Mesosinusid				
Carrilanga		17 (62.96%)	10 (37.04%)	27
Nombrevilla 3		2	1	3
Nombrevilla 2		56 (96.55%)	2 (3.45%)	58
Solera		29 (54.72%)	24 (45.28%)	53
Paje 2		4	1	5
Paje 1		1	—	1

		Absent 	Enlargement 	Short 	Connected 	Double 	N
Mesolophid							
Carrilanga		5 (19.23%)	6 (23.08%)	15 (57.69%)	—	—	26
Nombrevilla 3		1	—	2	—	—	3
Nombrevilla 2		44 (61.11%)	2 (2.78%)	23 (31.94%)	3 (4.17%)	—	72
Solera		9 (16.07%)	—	25 (44.65%)	13 (23.21%)	9 (16.07%)	56
Paje 2		1	—	3	1	—	5
Paje 1		—	—	1	—	—	1

m3



Short

Long



**Labial
Anterolophid**

**N**

Carrilanga	12 (80%)	3 (20%)	15
Nombrevilla 4	—	1	1
Nombrevilla 3	—	3	3
Nombrevilla 2	6 (7.50%)	74 (92.50%)	80
Solera	6 (13.95%)	37 (86.05%)	43
Paje 2	—	2	2

Absent

Short

Medium

Connected

Double







Mesolophid





N

Carrilanga	15 (78.95%)	1 (5.26%)	2 (10.53%)	—	1 (5.26%)	19
Nombrevilla 4	—	1	—	—	—	1
Nombrevilla 3	3	—	—	—	—	3
Nombrevilla 2	67 (66.34%)	28 (27.72%)	6 (5.94%)	—	—	101
Solera	22 (51.16%)	15 (34.88%)	1 (2.33%)	1 (2.33%)	4 (9.30%)	43
Paje 2	3	1	—	—	—	4

m3

	Absent	Present	
			
Ectomesolophid			N
Carrilanga	19	—	19
Nombrevilla 4	1	—	1
Nombrevilla 3	3	—	3
Nombrevilla 2	114	—	114
Solera	38	2	40
Paje 2	4	—	4

	Open	Low Cingulid	
			
Mesosinusid			N
Carrilanga	14 (77.78%)	4 (22.22%)	18
Nombrevilla 4	1	—	1
Nombrevilla 3	3	—	3
Nombrevilla 2	88 (96.70%)	3 (3.30%)	91
Solera	23 (56.10%)	18 (43.90%)	41
Paje 2	3	—	3


	Open	Cingulid	Two Crests	Stylid	
					
Sinusid					N
Carrilanga	7 (36.84%)	10 (52.63%)	—	2 (10.53%)	19
Nombrevilla 4	—	—	—	1	1
Nombrevilla 3	1	1	—	—	2
Nombrevilla 2	21 (26.25%)	54 (67.50%)	4 (5%)	1 (1.25%)	80
Solera	14 (34.15%)	26 (63.41%)	1 (2.44%)	—	41
Paje 2	1	2	—	—	3





Appendix 8.2





Appendix 8.2.




Morphological variability of *Hispanomys nombrevillae* (Cañada 10, Nombrevilla 1 and Nombrevilla 9) and *Hispanomys* cf. *nombrevillae* (Cañada 11, Cañada 9, Cañada 8 and Nombrevilla 10) in the Calatayud-Daroca Basin; number of specimens that have each state of character and the percentage (in brackets) over the total number of specimens (N).

M1










	Open	Low Cingul	Cingul	N
 Anterosinus				
Nombrevilla 1	1 (6.25%)	13 (81.25%)	2 (12.5%)	16
Nombrevilla 9	3	—	1	4






	Posterior Anterocone	Anterior Paracone	Double	N
 Anterior ectoloph				
Nombrevilla 1	2 (13.34%)	6 (40%)	7 (46.66%)	15
Nombrevilla 9	2	—	3	5





	Not Complete	Complete	N
 Anterior ectoloph			
Nombrevilla 1	—	15	15
Nombrevilla 9	1	4	5

M1




	Cingul	Style	Anteroloph	
				
Protosinus				N
Nombrevilla 1	6 (50%)	5 (41.67%)	1 (8.33%)	12
Nombrevilla 9	1	2	2	5

	Open	Cingul	Style	Two crests	
					
Sinus					N
Nombrevilla 1	9 (52.94%)	6 (35.29%)	1 (5.88%)	1 (5.88%)	17
Nombrevilla 9	—	2	2	1	5

	Open	Cingul	Low Cingul	
				
Mesosinus				N
Nombrevilla 1	1 (6.25%)	5 (31.25%)	10 (62.50%)	16
Nombrevilla 9	—	2	2	4

M2



Posterior Paracone

Anterior Anterocone

Double




Anterior
ectoloph



N

Cañada 10	—	—	1	1
Cañadas 8	—	1	—	1
Nombrevilla 1	3 (16.67%)	9 (50%)	6 (33.33%)	18
Nombrevilla 9	—	3 (37.50%)	5 (62.50%)	8

Complete








Anterior
ectoloph



N

Cañada 10	1	1
Cañadas 8	1	1
Nombrevilla 1	18	18
Nombrevilla 9	8	8



M2




	Simple	Double	
			
Posterior ectoloph			N
Cañada 10	—	1	1
Cañadas 8	1	—	1
Nombrevilla 1	2 (10.53%)	17 (89.47%)	19
Nombrevilla 9	—	8	8





	Complete	
		
Posterior ectoloph		N
Cañada 10	1	1
Cañadas 8	1	1
Nombrevilla 1	19	19
Nombrevilla 9	8	8

	Absent	
		
Mesoloph		N
Cañada 10	1	1
Cañadas 8	1	1
Nombrevilla 1	18	18
Nombrevilla 9	7	7


M2

	Absent	Present	
			
Entomesoloph			N
Cañada 10	1	—	1
Cañadas 8	1	—	1
Nombrevilla 1	17	—	17
Nombrevilla 9	4	3	7

	Open	Low Cingul	Cingul	
				
Mesosinus				N
Cañada 10	—	1	—	1
Cañadas 8	1	—	—	1
Nombrevilla 1	12 (66.67%)	6 (33.33%)	—	18
Nombrevilla 9	2 (28.57%)	2 (28.57%)	3 (42.86%)	7

	Open	Cingul	Two Crests	Style	
					
Sinus					N
Cañada 10	1	—	—	—	1
Cañadas 8	1	—	—	—	1
Nombrevilla 1	5 (29.41%)	11 (64.71%)	1 (5.88%)	—	17
Nombrevilla 9	4 (57.14%)	1 (14.29%)	1 (14.29%)	1 (14.29%)	7

M3



Absent


Present



Anterior Ectoloph

N

Cañada 10	—	1	1
Nombrevilla 1	3 (25%)	9 (75%)	12
Nombrevilla 10	2 (66.67%)	1 (33.33%)	3
Nombrevilla 9	—	4	4



Complete

Anterior Ectoloph

N

Cañada 10	1	1
Nombrevilla 1	9	9
Nombrevilla 10	1	1
Nombrevilla 9	4	4

Absent

Present









Lingual Aneroloph




N

Cañada 10	—	1	1
Nombrevilla 1	7	3	10
Nombrevilla 10	3	—	3
Nombrevilla 9	1	2	3

M3





	Absent	Medium	
			
Mesoloph			N
Cañada 10	1	—	1
Nombrevilla 1	10	1	11
Nombrevilla 10	3	—	3
Nombrevilla 9	4	—	4

	Absent	Present	
			
Neo-entoloph			N
Cañada 10	1	—	1
Nombrevilla 1	10	2	12
Nombrevilla 10	3	—	3
Nombrevilla 9	5	—	5

	Simple Paracone	Simple Protolophule	Double	
				
Posterior Ectoloph				N
Cañada 10	—	—	—	0
Nombrevilla 1	7 (77.78%)	2 (22.22%)	—	9
Nombrevilla 10	3	—	—	3
Nombrevilla 9	2	—	2	4

M3





Not Complete Complete

Posterior Ectoloph

			N
Cañada 10	1	—	1
Nombrevilla 1	—	12	12
Nombrevilla 10	—	3	3
Nombrevilla 9	—	5	5




Absent Present Well Developed

Posteroloph

				N
Cañada 10	—	—	—	0
Nombrevilla 1	7 (77.78%)	2 (22.22%)	—	9
Nombrevilla 10	3	—	—	3
Nombrevilla 9	2	—	2	4


Absent Present

Enamel-coated valley

			N
Cañada 10	1	—	1
Nombrevilla 1	9 (81.81%)	2 (18.18%)	11
Nombrevilla 10	3	—	3
Nombrevilla 9	3	1	4




m1



Absent

Short

Long



Labial
Anterolophid



N

Cañada 10	1	—	—	1
Cañada 9	—	1	—	1
Nombrevilla 1	—	6	11	17
Nombrevilla 10	—	1	1	2
Nombrevilla 9	—	4	3	7

Absent

Enlargement

Short



Mesolophid

N

Cañada 10	1	—	—	1
Cañada 9	—	—	1	1
Nombrevilla 1	11 (68.75%)	5 (31.25%)	—	16
Nombrevilla 10	—	—	—	1
Nombrevilla 9	4	—	2	7


m1

	Absent	Present	
Ectomesolophid			N
Cañada 10	1	—	1
Cañada 9	1	—	1
Nombrevilla 1	20	—	20
Nombrevilla 10	1	1	2
Nombrevilla 9	7	1	8


	Metalophulid I	Metalophulid I+II	
Metalophulid			N
Cañada 10	1	—	1
Cañada 9	—	1	1
Nombrevilla 1	16	4	20
Nombrevilla 10	1	1	2
Nombrevilla 9	4	3	7

	Open	Low Cingulid	Cingulid	
Mesosinusid				N
Cañada 10	—	1	—	1
Cañada 9	—	—	1	1
Nombrevilla 1	5 (23.80%)	13 (61.91%)	3 (14.28%)	21
Nombrevilla 10	—	—	—	—
Nombrevilla 9	2 (50%)	1 (25%)	1 (25%)	4

m1



Open
Cingulid

Posterosinusid			N
Cañada 10	1	—	1
Cañada 9	—	1	1
Nombrevilla 1	13 (65%)	7 (35%)	20
Nombrevilla 10	1	—	1
Nombrevilla 9	4	4	8


Cingulid
Low Cingulid
Stylid



Sinusid				N
Cañada 10	1	—	—	1
Cañada 9	1	—	—	1
Nombrevilla 1	17 (89.48%)	1 (5.26%)	1 (5.26%)	19
Nombrevilla 10	1	—	—	1
Nombrevilla 9	7	—	1	8

m2



Partially Divided

Completely Divided

Posterior
root


			N
Cañada 9	—	1	1
Cañadas 8	—	—	0
Nombrevilla 1	1 (14.29%)	6 (85.71%)	7
Nombrevilla 9	1	1	2

Long

Absent



Labial
anterolophid







Cañada 9	1
Cañadas 8	1
Nombrevilla 1	13
Nombrevilla 9	9





Ectomesolophid








Cañada 9	1
Cañadas 8	1
Nombrevilla 1	16
Nombrevilla 9	9


m2

	Cingulid	Two Crests	Stylid	
				N
Sinusid				
Cañada 9	—	1	—	1
Cañadas 8	1	—	—	1
Nombrevilla 1	10 (76.93%)	1 (7.69%)	2 (15.38%)	13
Nombrevilla 9	5 (71.42%)	2 (28.57%)	—	7

	Open	Low Cingulid	Stylid	
				N
Mesosinusid				
Cañada 9	—	—	1	1
Cañadas 8	—	1	—	1
Nombrevilla 1	4 (21.05%)	14 (73.68%)	1 (5.26%)	19
Nombrevilla 9	6 (66.66%)	3 (33.34%)	—	9

	Absent	Short	Connected	Double	
					N
Mesolophid					
Cañada 9	1	—	—	—	1
Cañadas 8	1	—	—	—	1
Nombrevilla 1	7 (35%)	6 (30%)	6 (30%)	1 (5%)	20
Nombrevilla 9	2 (22.22%)	4 (44.44%)	3 (33.33%)	—	9

m3



Absent

Short


Labial
Anterolophid



N

Nombrevilla 1	16 (80%)	4 (20%)	20
Nombrevilla 9	6 (54.54%)	5 (45.45%)	11

Absent



Ectomesolophid

Nombrevilla 1	19
Nombrevilla 9	12

Absent


Short

Medium

Double



Mesolophid



N

Nombrevilla 1	15 (75%)	3 (15%)	—	2 (10%)	20
Nombrevilla 9	6 (46.15%)	6 (46.15%)	1 (7.69%)	—	13

m3

Low Cingulid

Open

**Mesosinusid****N**

Nombrevilla 1	16 (80%)	4 (20%)	20
Nombrevilla 9	4 (40%)	6 (60%)	10

Cingulid

Stylid

Open

Two Crests

**Sinusid****N**


Nombrevilla 1	1 (5%)	17 (85%)	—	2 (10%)	20
Nombrevilla 9	2 (10%)	8 (40%)	1 (5%)	1 (5%)	12

Appendix 8.3

Appendix 8.3.

Morphological variability of *Hispanomys aragonensis* (Pedregueras 1A, Pedregueras 2A and Pedregueras 2C) and *Hispanomys* cf. *aragonensis* (Nombrevilla 22, Nombrevilla 13, Nombrevilla 14 and Nombrevilla 19) in the Calatayud-Daroca Basin; number of specimens that have each state of character and the percentage (in brackets) over the total number of specimens (N).

M1





	Open	Low Cingul	Cingul	N
Anterosinus				
Pedregueras 2C	2(12.5%)	14(87,5%)	0	16
Pedregueras 2A	20(45.45%)	22(50%)	2(4,54%)	44
Pedregueras 1A	1	—	—	1
Nombrevilla 14	—	2	—	2
Nombrevilla 13	1	—	—	1

	Posterior Anterocone	Anterior Paracone	Double	N
Anterior ectoloph				
Pedregueras 2C	10(55.56%)	—	8(44.44%)	18
Pedregueras 2A	3(6.82%)	5(11.36%)	36(81.82%)	44
Pedregueras 1A	—	—	1	1
Nombrevilla 14	—	—	2	2
Nombrevilla 13	—	—	1	1

	Anterior ectoloph	Complete	N
Anterior ectoloph			
Pedregueras 2C		18	18
Pedregueras 2A		44	44
Pedregueras 1A		1	1
Nombrevilla 14		2	2
Nombrevilla 13		1	1

M1



Simple Double

Posterior ectoloph **N**

Pedregueras 2C	3(15.78%)	16(84.21%)	19
Pedregueras 2A	—	40	40
Pedregueras 1A	—	1	1
Nombrevilla 14	—	2	2
Nombrevilla 13	—	1	1





Complete

Posterior ectoloph **N**






Pedregueras 2C	19	19
Pedregueras 2A	40	40
Pedregueras 1A	1	1
Nombrevilla 14	2	2
Nombrevilla 13	1	1






Absent Enlargement Short Medium










Mesoloph **N**


Pedregueras 2C	13(81.25%)	—	—	3(18.75%)	16
Pedregueras 2A	24(63.15%)	4(10.52%)	8(21.05%)	2(5.26%)	38
Pedregueras 1A	1	—	—	—	1
Nombrevilla 14	2	—	—	—	2
Nombrevilla 13	1	—	—	—	1






	Open	Cingul	Style	M1 Anteroloph	N
					
Protosinus					
Pedregueras 2C	—	4(23.53%)	5(29.41%)	8(47.06%)	17
Pedregueras 2A	1(2.22%)	3(6.67%)	18(40%)	23(51.11%)	45
Pedregueras 1A	—	1	—	—	1
Nombrevilla 14	—	—	2	—	2
Nombrevilla 13	—	—	—	—	—




	Open	Cingul	Style	Two Crests	N
					
Sinus					
Pedregueras 2C	15(88.24%)	—	1(5.88%)	1(5.88%)	17
Pedregueras 2A	22(52.38%)	2(4.76%)	14(33.33%)	4(9.52%)	42
Pedregueras 1A	1	—	—	—	1
Nombrevilla 14	2	—	—	—	2
Nombrevilla 13	1	—	—	—	1

	Open	Cingul	Low Cingul	N
				
Mesosinus				
Pedregueras 2C	3(18.75%)	—	13(81.25%)	16
Pedregueras 2A	20(50%)	3(7.5%)	17(42.5%)	40
Pedregueras 1A	1	—	—	1
Nombrevilla 14	—	1	—	1
Nombrevilla 13	1	—	—	1




M2








	Absent	Posterior Paracone	Anterior Anterocone	Double	N
 Anterior ectoloph					
Pedregueras 2C	1(3.45%)	5(17.24%)	6(20.69%)	17(58.62%)	29
Pedregueras 2A	—	5(9.09%)	17(30.91%)	33(60%)	55
Pedregueras 1A	2	—	—	2	4
Nombrevilla 22	2	—	—	—	2

	Complete	N
 Anterior ectoloph		
Pedregueras 2C	28	28
Pedregueras 2A	56	56
Pedregueras 1A	2	2
Nombrevilla 22	—	—



M2





	Simple	Double	
			
Mesoloph			N
Pedregueras 2C	26(96.30%)	1(3.70%)	27
Pedregueras 2A	44(80%)	11(20%)	55
Pedregueras 1A	4	—	4
Nombrevilla 22	2	—	2




	Absent	Short	
			
Posterior ectoloph			N
Pedregueras 2C	9	20	29
Pedregueras 2A	—	54	54
Pedregueras 1A	—	4	4
Nombrevilla 22	—	2	2

	Complete	
		
Posterior ectoloph		N
Pedregueras 2C	29	29
Pedregueras 2A	56	56
Pedregueras 1A	4	4
Nombrevilla 22	2	2


M2

	Absent	Present	
			
Entomesoloph			N
Pedregueras 2C	27(93.10%)	2(6.90%)	29
Pedregueras 2A	50(92.59%)	4(7.40%)	54
Pedregueras 1A	3	1	4
Nombrevilla 22	2	—	2

	Open	Low cingul	Cingul	Style	
					
Mesosinus					N
Pedregueras 2C	13(44.83%)	12(41.37%)	3(10.34%)	1(3.45%)	29
Pedregueras 2A	45(81.82%)	—	10(18.18%)	—	55
Pedregueras 1A	4	—	—	—	4
Nombrevilla 22	—	2	—	—	2

	Open	Cingul	Style	
				
Sinus				N
Pedregueras 2C	25(92.59%)	2(7.41%)	—	27
Pedregueras 2A	48(96%)	2(4%)	4(8%)	50
Pedregueras 1A	4	—	—	4
Nombrevilla 22	2	—	—	2

M3



Absent

Posterior

Anterior





Anterior Ectoloph

N

Pedregueras 2C	28(70%)	6(15%)	6(15%)	40
Pedregueras 2A	26(60.47%)	9(20.93%)	8(18.60%)	43
Pedregueras 1A	2	—	—	2
Nombrevilla 14	—	—	2	2
Nombrevilla 22	2	—	—	2

Not Complete

Complete



Anterior Ectoloph

N

Pedregueras 2C	1(8.3%)	11(91.67%)	12
Pedregueras 2A	1(5.88%)	16(94.12%)	17
Pedregueras 1A	—	—	0
Nombrevilla 14	—	2	2
Nombrevilla 22	—	—	0

Absent

Present



Lingual Aneroloph

N

Pedregueras 2C	30(83.33%)	6(16.67%)	36
Pedregueras 2A	33(84.62%)	6(15.38%)	39
Pedregueras 1A	—	1	1
Nombrevilla 14	2	—	2
Nombrevilla 22	2	—	2

M3



Absent



Short



Medium



Large




Mesoloph


N

Pedregueras 2C	32(84.21%)	4(10.53%)	1(2.63%)	1(2.63%)	38
Pedregueras 2A	33(78.57%)	6(14.29%)	—	3(7.14%)	42
Pedregueras 1A	2	—	—	—	2
Nombrevilla 14	2	—	—	—	2
Nombrevilla 22	—	—	—	2	2

Absent



Present



Neo-entoloph

N

Pedregueras 2C	38(95%)	2(5%)	40
Pedregueras 2A	42	—	42
Pedregueras 1A	2	—	2
Nombrevilla 14	2	—	2
Nombrevilla 22	2	—	2

Absent

Simple
Paracone




Double

Posterior
Ectoloph


N

Pedregueras 2C	1(2.44%)	5(12.20%)	35(85.37%)	41
Pedregueras 2A	—	28(65.11%)	15(34.88%)	43
Pedregueras 1A	—	—	2	2
Nombrevilla 14	—	1	1	2
Nombrevilla 22	—	—	2	2

M3



m1



Short

Long



Labial
Anterolophid





			N
Pedregueras 2C	20(43.48%)	26(56.52%)	46
Pedregueras 2A	27(50.94%)	26(49.06%)	53
Pedregueras 1A	1	1	2
Nombrevilla 14	1	—	1
Nombrevilla 13	1	—	1

Absent

Enlargement

Short

Medium




Mesolophid

					N
Pedregueras 2C	17(36.96%)	11(23.91%)	18(39.13%)	—	46
Pedregueras 2A	10(17.86%)	12(21.43%)	26(46.43%)	8(14.29%)	56
Pedregueras 1A	—	—	3	—	3
Nombrevilla 14	1	—	—	—	1
Nombrevilla 13	—	—	1	—	1

m1

Absent

Present




**Ectomesolophid****N**

Pedregueras 2C	47	—	47
Pedregueras 2A	52	1	53
Pedregueras 1A	3	—	3
Nombrevilla 14	—	—	—
Nombrevilla 13	1	—	1

Absent

Metalophulid I

Metalophulid I+II

**Metalophulid****N**





Pedregueras 2C	1(2.17%)	41(89.13%)	4(8.70%)	46
Pedregueras 2A	—	48(88.89%)	6(11.11%)	54
Pedregueras 1A	—	3	—	3
Nombrevilla 14	—	1	—	1
Nombrevilla 13	—	1	—	1

Open

Low Cingulid

Cingulid

Stylid

**Mesosinusid****N**


Pedregueras 2C	34(72.34%)	9(19.15%)	3(6.38%)	1(2.13%)	47
Pedregueras 2A	35(64.81%)	18(33.33%)	—	1(1.85%)	54
Pedregueras 1A	—	—	2	—	2
Nombrevilla 14	—	—	—	—	—
Nombrevilla 13	—	—	—	—	—

m1



Open

Cingulid

**Posterosinusid****N**

Pedregueras 2C	26(59.09%)	18(40.91%)	44
Pedregueras 2A	45(83.33%)	9(16.67%)	54
Pedregueras 1A	3	—	3
Nombrevilla 14	—	—	—
Nombrevilla 13	1	—	1



Open

Cingulid


Two Crests

Stylid


Sinusid**N**

Pedregueras 2C	4(9.09%)	12(27.27%)	2(4.55%)	26(59.09%)	44
Pedregueras 2A	1(1.92%)	29(55.77%)	5(9.62%)	17(32.69%)	52
Pedregueras 1A	—	3	—	—	3
Nombrevilla 14	—	—	—	—	—
Nombrevilla 13	—	—	—	—	—

m2



Completely Divided

Posterior
root**N**

Pedregueras 2C	4	4
Pedregueras 2A	—	—

Open

Low Cingulid

**Mesosinusid****N**

Pedregueras 2C	11 (39.28%)	17 (60.71%)	28
Pedregueras 2A	12 (26.08%)	34 (73.91%)	46


Short

Long


Labial
anterolophid**N**

Pedregueras 2C	3 (12%)	22 (88%)	25
Pedregueras 2A	5 (13.16%)	33 (86.84%)	38


m2



		N
Pedregueras 2C	29	29
Pedregueras 2A	42	42




					N
Pedregueras 2C	1(3.70%)	13(48.15%)	2(7.41%)	11(40.74%)	27
Pedregueras 2A	3(7.69%)	30(76.92%)	4(10.26%)	2(5.13%)	39



						N
Pedregueras 2C	3(10%)	—	10(33.33%)	12(40%)	5(16.66%)	30
Pedregueras 2A	1(3.23%)	6(19.35%)	21(67.74%)	1(3.23%)	2(6.45%)	31

m3



Short

Large



**Labial
Anterolophid**

**N**

Nombrevilla 19	—	1	1
Pedregueras 2C	3 (8.33%)	33 (91.67%)	36
Pedregueras 2A	—	28	28
Pedregueras 1A	—	2	2
Nombrevilla 14	—	3	3

Absent

Present







Ectomesolophid

N

Nombrevilla 19	1	—	1
Pedregueras 2C	38	—	38
Pedregueras 2A	—	8	8
Pedregueras 1A	2	—	2
Nombrevilla 14	3	—	3

m3




Absent Short Double

Mesolophid **N**

Nombrevilla 19	1	—	—	1
Pedregueras 2C	15(42.9%)	15(42.9%)	5(14.3%)	35
Pedregueras 2A	14(46.67%)	16(53.33%)	—	30
Pedregueras 1A	2	—	—	2
Nombrevilla 14	1	2	—	3






Open Low Cingulid

Mesosinusid **N**

Nombrevilla 19	1	—	1
Pedregueras 2C	27(72.97%)	10(27.02%)	37
Pedregueras 2A	25(83.33%)	5(16.66%)	30
Pedregueras 1A	2	—	2
Nombrevilla 14	2	1	3

Open Cingulid Two Crests Styloid


Sinusid **N**

Nombrevilla 19	—	1	—	—	1
Pedregueras 2C	11(31.43%)	10(23.57%)	1(2.86%)	13(37.14%)	35
Pedregueras 2A	5(17.86%)	14(50%)	1(3.57%)	8(28.57%)	28
Pedregueras 1A	2	—	—	—	2
Nombrevilla 14	—	2	1	—	3

IV. Bioestratigrafía

Chapter 9

Biostratigraphy



CRICETODONTINI FROM THE CALATAYUD-DAROCA BASIN (SPAIN): AN UP-TO-DATE OF THEIR STRATIGRAPHICAL DISTRIBUTIONS

“On a regional scale, obviously, a detailed stratigraphical framework based on local biozonation, supported by superposition, radiometric ages and palaeomagnetostratigraphy, is far superior and in our opinion the best way to proceed

Albert van der Meulen & colaboradores.

9


Resumen

Se presentan las distribuciones estratigráficas de los miembros de la tribu Cricetodontini de la Cuenca de Calatayud-Daroca. Algunos géneros de Cricetodontini fueron utilizados, en la Cuenca de Calatayud-Daroca, para caracterizar ciertas unidades cronoestratigráficas (pisos continentales). Sin embargo, las especies de Cricetodontini no han sido, normalmente, utilizadas en la definición y propuesta de las escalas bioestratigráficas locales. No obstante, estudios recientes en las cuencas de Europa central y Anatolia muestran que las especies de Cricetodontini encontradas allí se pueden utilizar para establecer subdivisiones bioestratigráficas locales. Los resultados presentados en este capítulo muestran que, para la Cuenca de Calatayud-Daroca, la mayoría de las distribuciones estratigráficas de las especies de Cricetodontini están estrechamente vinculadas con la biozonación local. Esto sugiere que podrían ser utilizados como buenos indicadores bioestratigráficos. Por otra parte, la biozona local G3, que es inusualmente larga, podría ser redefinida y subdividida basándonos en las distribuciones de las cuatro especies de Cricetodontini presentes en ella. Sin embargo, se recomienda completar el estudio de toda la fauna de roedores antes de proponer una nueva biozonación.

Referencia

Este artículo señalado abajo recoge parte del contenido de este capítulo, con algunas modificaciones:

López-Guerrero, P., I., García-Paredes, M. A., Álvarez-Sierra, and Peláez-Campomanes, P. 2014. Cricetodontini from the Calatayud-Daroca Basin (Spain): a taxonomical description and up-to-date of their stratigraphical distributions. *Comptes Rendus Palevol*. En prensa.



Abstract

The most updated stratigraphical distributions of the Cricetodontini from the Calatayud–Daroca Basin are presented. Generally, the genera of Cricetodontini were used to define large time intervals and limits between major continental units, whereas the species were neglected in the local biostratigraphical studies. Nevertheless, recent studies in central Europe and Anatolia indicate that the species of Cricetodontini have served as a source to establish biostratigraphical local subdivisions. The results presented herein show that, for the Calatayud–Daroca Basin, most of the stratigraphical distributions of the species of Cricetodontini are strongly linked with the local biozonation. That reveals that they could be used as good biostratigraphical indicators. Moreover, the unusually long local biozone G3 could be redefined and subdivided based on the distributions of the four species of Cricetodontini present in it. However, it is recommended to complete the study of the whole rodent fauna before proposing a new biozonation.

INTRODUCTION

The tribe Cricetodontini is a group of large-sized rodents found in Europe, northern Africa and Asia (Jaeger et al., 1973; De Bruijn and Ünay, 1996; Rummel, 1999). Their biogeographical distribution follows a symmetrical model in which the first occurrence of the tribe is in a small area (region of Anatolia) during the early Miocene (De Bruijn and Ünay, 1996; Rummel, 1999; Álvarez-Sierra et al., 2013). In the middle Miocene there is a gradual expansion into central Europe and France followed by a period of stasis in which the taxon is distributed over most Europe and Asia Minor (De Bruijn and Ünay, 1996; Rummel, 1999; Álvarez-Sierra et al., 2013). Their extinction at the end of the Pliocene was preceded by a period of fragmentation and contraction of their distribution area since the late middle Miocene (De Bruijn and Ünay, 1996; Rummel, 1999; Álvarez-Sierra et al., 2013a; 2013b).

This tribe includes eight genera; three of them—*Cricetodon* Lartet, 1851, *Hispanomys* Mein and Freudenthal, 1971 and *Ruscinomys* Depéret, 1890—are recorded in the Aragonian and Vallesian from the Calatayud-Daroca Basin.

Genera of Cricetodontini have been used, as biostratigraphical indicators, in many studies for the Calatayud-Daroca Basin, (e.g., Daams et al., 1999; Álvarez Sierra et al., 2003; Hernández-Ballarín et al., 2011; Van der Meulen et al., 2011, 2012), although mainly to define large time intervals and limits between major continental units. For instance, the first occurrence (FO) of *Cricetodon* coincides with the beginning of the local biozone E in the Calatayud-Daroca Basin (Daams et al., 1999; Van der Meulen et al., 2012), and the relative abundance of *Cricetodon* and *Hispanomys* characterizes the faunas from the late Aragonian and the early Vallesian in the same basin (Daams et al., 1999).

By contrast, the species of Cricetodontini have been not traditionally used for local biostratigraphical subdivisions in the Calatayud-Daroca Basin. De Bruijn et al. (1993) pointed out that the mosaic morphology displayed by the species of Cricetodontini precludes their use for the proposal of detailed biostratigraphical scales. Nevertheless, the recently exhaustive study of the species of Cricetodontini have served as a source to establish biostratigraphical subdivisions. For instance, the species of *Byzantinia* and *Cricetodon* were used in the local biostratigraphy of the Neogene from Turkey (Ünay and De Bruijn, 1984; Ünay et al., 2003). Furthermore, in the workshop of the Regional Committee of Stratigraphy of the Neogene in May 2012, the biostratigraphy of the uppermost early Badenian (middle Aragonian) from the German Molasse were updated from

those made by Heiig (1997). Three species of *Cricetodon* were used to define three local biozones: *Cricetodon meini*, *Cricetodon aureus* and *Cricetodon sansaniensis* which correspond to OMS units E', F, and ?F (Bhme et al., 2012) (Fig. 9.1a). Moreover, Klin and Kempf (2009) also used a similar approach in the biostratigraphy of the Swiss Molasse. Therefore, it seems that a better knowledge of the species of Cricetodontini could reveal them as good local biostratigraphical and biochronological indicators (Fig. 9.1b). Furthermore, the zonation of Van der Meulen et al. (2012), based mainly on cricetids (e.g., *Megacricetodon*, *Democricetodon*), could be extended towards the late Aragonian and the early Vallesian; indeed, the most updated distribution of the Cricetodontini will be useful in the redefinition of the local biozonation.

Several works have recently focused in the systematics of the tribe Cricetodontini from the Calatayud–Daroca Basin (Lpez-Guerrero et al., 2008, 2009, 2013a, 2014), but a number of localities and species from the earliest part of the late Aragonian still lacked a detailed revision. Thus, the main aim of this study is to evaluate the updated stratigraphical distributions of the Cricetodontini from the Calatayud-Daroca Basin described in chapters 5 to 8 and assess whether or not they are good biostratigraphical indicators.





FIGURE 9.1. Stratigraphical distributions of the species of Cricetodontini in the central Europe Basins. (a). Germany Molasse. Correlation of the local biozones with continental stages and epoch is also present. Data source: Bhme et al. (2012); (b). Swiss Molasse. Localities, correlation with local biozones, the MN system and epoch are also presented. Data source: Klin and Kempf, (2009).



(b)

RECORDED SPECIES OF CRICETODOTINI

After the systematic study of the *Cricetodontini* from the Calatayud-Daroca Basin, eight different species have been found. Their stratigraphic distributions and a summary of their morphological features are presented below.

CRICETODON SORIAE López Martínez, Cárdaba, Salesa,
Hernández Fernández, Cuevas González and Fesharaki, 2006

The most important features of *C. soriae* are: medium to large size; brachydont molars without styles or stylids. The upper molars have short or absent ectolophs, the anterior ectoloph is absent on the M1; presence of split anterocone and anterior + posterior protolophule on the M1, as well as well-developed posteroloph on the M3. The lower molars often show ectomesolophids and low-developed mesolophids; the m1 possess both metalophulid I and II; incomplete metalophulid I and rounded anteroconid; the m3 is large and longer than the m1, and it has long and double mesolophids.

Type locality—Somosaguas N, Madrid Basin, Spain, local biozone E (MN5, middle Aragonian, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone E (MN5, middle Aragonian, middle Miocene). FO: LUM11 (14.06 Ma); LO: LUM20 (13.80 Ma)

CRICETODON SANSANIENSIS Lartet, 1851

The most important features of *C. sansaniensis* are: large size; M1 with an slightly divided anterocone. Ectolophs absent in half of the M1. Short mesoloph present in some M1, slightly more frequent in M2 and M3. The anterior surface of both the paracone and the metacone notched by a vertical groove. Undivided and rounded anteroconid in the m1; the metaconid has a metalophulid II in more than half of the specimens, I+II or only I in the other specimens. Mesolophid is absent or rare in the m1, more frequent in the m2 and the m3, but always short. M1 with four roots.

Type locality—Sansan, France, (MN6, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone F (MN6, upper Aragonian, middle Miocene) FO: LUM22 (13.76 Ma); LO: VT2C (13.68 Ma).

CRICETODON JOTAE Mein and Freudenthal, 1971

The most important features are: poor-developed mesolophs and posterolophs present in the upper molars. The anterior ectoloph of the M1 composed by the anterior ectoloph of the paracone or by the posterior ectoloph of the anterocone. The M1 with posterior ectoloph; composed by the posterior ectoloph of the paracone, by the anterior ectoloph of the metacone, or by both. The ectolophs are not completed. The main valleys frequently blocked by cinguls and styles. The lower molars have short or absent mesolophids, ectomesolophids and lingual anterolophids. Most of the m1 display metalophulid I and metalophulid II together in the same specimen, but some m1 have only metalophulid I or Metalophulid II. The m2 possess two or three roots and the posterior root that can be either partially or completely split.

Type locality—Manchones, Calatayud-Daroca Basin, Spain, local biozone G2, (MN6 upper Aragonian, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone G1-local biozone G3 (MN6-MN7/8, upper Aragonian, middle Miocene). FO: LP5B (13.56 Ma); LO: TOR1.

CRICETODON NIEVEI López-Guerrero, Álvarez-Sierra, García-Paredes
and Peláez-Campomanes, 2014

The most important features of *C. nievei* are: the thick enamel; the presence of styles on the protosinus of the M1; the moderate development of the ectolophs—being incomplete in most specimens—; the moderately simple pattern of the M3; the presence of metalophulid I on the m1 and absence of metalophulid II; the presence of ectomesolophid, and the moderately reduced m3.

Type locality—Toril 3A, Calatayud-Daroca Basin, Spain local biozone G3 (MN7/8, upper Aragonian, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—Local biozone G3 (MN7/8, upper Aragonian, middle Miocene). FO: TOR3A/3B (12.65 Ma); LO: LP5H (12.60

Ma).

HISPANOMYS AGUIRREI (Sesé Benito, 1977 in López Martínez et al., 1977)

Hispanomys aguirrei presents features such as: anterior ectoloph frequently double and complete, but also there are ectolophs simple and not complete; posterior ectoloph frequently complete and composed by both anterior ectoloph of the metacone and posterior ectoloph of the paracone; mesoloph in the upper molars almost always absent; frequently presence of posteroloph in the M3, which is well developed in some specimens; presence of metalophulid I and II together in the same m1 or only the metalophulid I; mesolophid and ectomesolophid highly reduced or absent and three-rooted m2.

Type locality—Escobosa de Calatañazor, Duero Basin, Spain (MN7/8, upper Aragonian, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—Local biozone G3 (MN7/8, upper Aragonian, middle Miocene). FO: PJE1 (12.39 Ma); LO: CARR (11.33 Ma).

HISPANOMYS LAVOCATI (Freudenthal, 1966)

The most important features of *H. lavocati* are: well-developed ectolophs; frequently composed by both anterior and posterior ectolophs and connected; posterior ectoloph on the M3 usually starting from the protolophule instead of from the paracone; axioloph interrupted in several M3; absent metalophulid II in the m1 and m2 always three-rooted.

Type locality—Hostalets de Pierola, Vallés-Penedés Basin, Spain (MN7/8, upper Aragonian, middle Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone G3, (MN7/8, upper Aragonian, middle Miocene). FO: NOM2 (11.87 Ma); LO: NOM4 (11.80 Ma).

HISPANOMYS NOMBREVILLAE (Freudenthal, 1966)

The most important features of *H. nombrevillae* are: four-rooted M1; anterior ectoloph always present, always complete and mainly composed by the posterior ectoloph of the anterocone; anterior ectoloph composed mainly by the anterior ectoloph of the paracone; posterior ectoloph

is always present and composed mainly by the posterior ectoloph of the paracone and the anterior ectoloph of the metacone, and it is always complete; mesoloph absent; valleys open or closed by a low cingulum; M3 highly reduced; mesolophid presents in half of the lower molars; metalophulid I is always present, some also have metalophulid II and most m2 displays three roots completely split.

Type locality—Nombrevilla 1, Calatayud-Daroca Basin, Spain, local biozone H (MN9, lower Vallesian, upper Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone H (MN7/8-MN9, upper Aragonian-lower Vallesian, middle-upper Miocene). FO: NOM9 (11.20 Ma); LO: NOM1 (10.77 Ma).

HISPANOMYS ARAGONENSIS (Freudenthal, 1966)

The most important features of *H. aragonensis* are: relative small size; M1 with four roots; well-developed ectolophs, mainly double and always complete; the posterior ectoloph of the M2 is oblique and connects the metacone to the point where the posterior protolophule reaches the longitudinal crest; mesoloph and enamel-coated valley frequent on the M3; weak lingual and labial posteroloph; valleys mainly blocked by low cingula; weak posteroloph on the M3; well-developed labial anterolophid; mesolophid present in most of the lower molars, it can be double or connected with the metacone; valleys closed by low cingulids; m1 frequently shows a metalophulid I, and in some specimens, the metalophulid II is also present.

Type locality—Pedregueras 2C, Calatayud-Daroca Basin, Spain, local biozone I (MN9, lower Vallesian, upper Miocene).

Stratigraphical distribution in the Calatayud-Daroca Basin—local biozone H-local biozone I, (MN9, lower Vallesian, upper Miocene). FO: NOM13 (10.72 Ma); LO: NOM19 (9.98 Ma).

Discussion

All the material of Cricetodontini from Calatayud-Daroca Basin is now taxonomically assigned, including the most recently recovered specimens (Table 9.1). Eight species of Cricetodontini split in two genera have been recognized; four belong to *Cricetodon* and the other four, to *Hispanomys*. The stratigraphical distribution of Cricetodontini in the Calatayud-Daroca Basin

(Fig. 9.2) ranges from the local biozone E to I for the middle and upper Aragonian, and lower Vallesian (middle-upper Miocene). The oldest species is *C. soriae* and the youngest is *H. aragonensis*. *Hispanomys* replaced *Cricetodon* in the middle of the local biozone G3, around 12.60 and 12.39 Ma.

The stratigraphical distributions of the studied species (Fig. 9.2) generally fit well with the local biozonation; those of *Cricetodon soriae*, *C. sansaniensis* and *H. aragonensis* coincide with the local biozones E, F, and I, respectively and therefore these species could be used as good biostratigraphical markers. On the other hand, the distribution of *Cricetodon jotae* is longer and it ranges about 1 my, covering several local biozones (G1, G2 and partly G3). As it shown in figure 9.2, four species of Cricetodontini are present in local biozone G3 (*C. jotae*, *C. nievei*, *H. aguirrei* and *H. lavocati*). On the light of the distribution of this species and the unusually long duration of the G3 (1 my) the redefinition and subdivision of this biozone is recommended. Nevertheless, we suggest a study of the complete rodent fauna before proposing a new biozonation. The local biozonation built for the Calatayud-Daroca Basin (Daams et al., 1999; Van der Meulen et al., 2012) uses mainly species of cricetids (*Megacricetodon*, *Democricetodon* and *Eumyarion*), comyids (*Ligerimys*), sciurids (*Heteroxerus*) and Insectivores (*Galerix*). Nonetheless, such redefinition is out of the scope of this Ph.D.

The stratigraphical distribution of *Hispanomys nombrevillae* is placed within the local biozone H covering part of the late Aragonian and part of the early Vallesian. This is because of NOM9. This locality yields a small mammal assemblage typical from the biozone H (Álvarez-Sierra et al., 2003; Garcés et al., 2003). The local biozone H is classically correlated to the lower Vallesian (Daams et al., 1999) but the beginning of the Vallesian is marked by the first occurrence of *Hipparion*. NOM9 presents large mammals typical from the Vallesian, but *Hipparion* is absent (Álvarez-Sierra et al., 2003) thus it has to be placed in the late Aragonian. In consequence, the limit of local biozone H does not necessarily coincide with the Aragonian-Vallesian boundary. The age of NOM9 (and the limit of the local biozone H) is 11.21 Ma (Van Dam et al., in press), whereas the earliest occurrence of *Hipparion* in the Calatayud-Daroca Basin is in NOM1, dated in 10.77 Ma (Van Dam et al., in press).

Rögl and Daxner-Höck (1996) dated the oldest *Hipparion* remains at 11.20 Ma; Agustí et al. (2001) defined the *Hipparion*-datum at 11.10 Ma in the Vallès-Penedès Basin; Sen (1997) correlated the first appearance datum (FAD) of *Hipparion* with the chron C5m, at 10.80 Ma, and Krijgsman et al. (1996) placed the boundary of the Vallesian in the Duero Basin (Spain) in 11.10 Ma. Therefore, it is widely accepted that the first occurrence of *Hipparion* can be considered diachronic. According to these data, the *Hipparion*-datum and age of the limit of the Vallesian ranges from 11.20 to 10.77 Ma. Moreover, in the Geologic Time Scale (Hilgen et al., 2012), the astrochronological age for this boundary is 11.20 Ma, which is consistent with the age of NOM9

(11.21 Ma). NOM9 lacks *Hipparion*, but its faunal assemblage is typical of the Vallesian, so the age of NOM9 agrees with the boundary between the Aragonian and the Vallesian.

The evolutionary patterns observed in the Cricetodontini from Calatayud-Daroca Basin agree with the following model: basal morphology in the first registered taxa (*C. soriae*, *C. sansaniensis*), mosaic configuration in most part of the late Aragonian (*C. jotae*, *C. nievei*, *H. aguirrei* and *H. lavocati*) and, finally, derived forms starting in *H. nombrevillae*. Most of the species from the Calatayud-Daroca Basin are endemic and therefore not found elsewhere. However, this morphological patterns are also recognized in the rest of the European Cricetodontini faunas, although not synchronously (Kálin and Kempf, 2009; Álvarez-Sierra et al., 2013a, 2013b; López-Guerrero et al., 2013b, 2014). In line with this, Van der Meulen et al., (2011, 2012) detected several asynchronies studying the rodent bioevents sequence between from the Calatayud-Daroca Basin and central Europe during the early and middle Aragonian. The asynchrony in the evolutionary patterns of the Cricetodontini between Iberian and central Europe Basins provides new criteria that support the asynchronies noted by the works of Van der Meulen et al. (2011, 2012).

The estimated age of the localities containing these species of Cricetodontini spans a time range from 14.06 to 9.98 Ma (Daams et al., 1999; Van Dam et al., 2006, Van Dam et al., in press Table 9.1). Several important paleoclimatic changes have been detected within this interval; firstly, the Miocene Climatic Optimum (MCO), a period with high temperatures and humid tropical and subtropical conditions, is recorded worldwide (Zachos et al., 2001; Shevenell et al., 2004). The MCO predates the first occurrence of *Cricetodon* in the Calatayud-Daroca Basin. The MCO was followed by the Middle Miocene Climatic Transition (MMCT), a period characterized by Antarctic ice growth and gradual global cooling. The MMCT ends in a short-term drop in sea surface temperatures called the Middle Miocene Cooling (MMC). The MMC was triggered by the reestablishment of the eastern Antarctica ice sheet; it has been recognized also in some continental deposits from the Iberian Peninsula (Domingo et al., 2012) and represents the beginning of the cold climatic regime of the Neogene (Lear et al., 2000; Zachos et al., 2001; Shevenell et al., 2004).

Hordijk (2010) noted that these climatic phases (MCO, MMCT and MMC) agreed well with the timing of the main changes in the mammal succession in the Calatayud-Daroca Basin. He pointed out that the gradual cooling of the MMCT is correlated with a prolonged immigration phase which produced an increase of the richness in the small mammal faunas. According to Hordijk (2010) the major turnover in the Calatayud-Daroca Basin took place during local biozones E to G2; consequently the FO of *Cricetodon* in the Calatayud-Daroca Basin (local Biozone E) is part of this event.

Locality	Abb.	Biozone	Age	Species	Locality	Abb.	Biozone	Age	Species
Nombrevilla 19	NOM19	I	9.98	<i>H. cf. aragonensis</i>	Toril 1	TOR1	G3	—	<i>C. jotae</i>
Pedregueras 2C	PED2C	I	10.38	<i>H. agonensis</i>	Alcocer 3	AC3	G3	—	<i>C. jotae</i>
Pedregueras 2A	PED2A	I	10.62	<i>H. agonensis</i>	Alcocer 2	AC2	G3	—	<i>C. jotae</i>
Pedregueras 1A	PDA	I	—	<i>H. agonensis</i>	Alcocer 1	AC1	G3	—	<i>C. jotae</i>
Nombrevilla 14	NOM14	I	10.68	<i>H. cf. aragonensis</i>	Villafeliche 9	VL9	G3	—	<i>C. jotae</i>
Nombrevilla 13	NOM13	I	10.72	<i>H. cf. aragonensis</i>	Arroyo del Val VI	ARV6	G2	—	<i>C. jotae</i>
Nombrevilla 22	NOM22	I	—	<i>H. cf. aragonensis</i>	Las Planas 5K	LP5K	G2	13.08	<i>C. jotae</i>
Cañada 11	CAÑ11	H	—	<i>H. cf. nombrevillae</i>	Las Planas 5L	LP5L	G2	13.16	<i>C. jotae</i>
Cañada 10	CAÑ10	H	—	<i>H. cf. nombrevillae</i>	Borjas	BOR	G2	—	<i>C. jotae</i>
Cañada 9	CAÑ9	H	—	<i>H. cf. nombrevillae</i>	Manchones	MAN	G2	—	<i>C. jotae</i>
Cañada 8	CAÑ8	H	—	<i>H. cf. nombrevillae</i>	Valtalto 1	VT1	G1	13.30	<i>C. jotae</i>
Nombrevilla 1	NOM1	H	10.77	<i>H. nombrevillae</i>	Las Planas 5C	LP5C	G1	13.55	<i>C. jotae</i>
Nombrevilla 10	NOM10	H	11.18	<i>H. cf. nombrevillae</i>	Las Planas 5B	LP5B	G1	13.56	<i>C. jotae</i>
Nombrevilla 9	NOM9	H	11.20	<i>H. nombrevillae</i>	Valalto 2C	VT2C	F	13.68	<i>C. sansanienis</i>
Carrilanga 1	CARR1	G3	11.33	<i>H. aguirrei</i>	Valtalto 2B	VT2B	F	13.70	<i>C. sansanienis</i>
Nombrevilla 4	NOM4	G3	11.80	<i>H. aguirrei/H. lavocati</i>	Armantes 7	ARM7	F	—	<i>C. sansanienis</i>
Nombrevilla 3	NOM3	G3	11.84	<i>H. aguirrei/H. lavocati</i>	Las Umbrias 21	LUM21	F	13.75	<i>C. sansanienis</i>
Nombrevilla 2	NOM2	G3	11.87	<i>H. aguirrei/H. lavocati</i>	Las Umbrias 22	LUM22	F	13.76	<i>C. sansanienis</i>
Solera	SOL	G3	12.01	<i>H. cf. aguirrei</i>	Las Umbrias 20	LUM20	E	13.80	<i>C. soriae</i>
Paje 2	PJE2	G3	12.26	<i>H. cf. aguirrei</i>	Las Planas 4C	LP4C	E	13.88	<i>C. soriae</i>
Paje 1	PJE1	G3	12.39	<i>H. cf. aguirrei</i>	Las Umbrias 19	LUM19	E	13.95	<i>C. soriae</i>
Las Planas 5H	LP5H	G3	12.60	<i>C. nievei</i>	Las Planas 4B	LP4B	E	13.96	<i>C. soriae</i>
Toril 3B	TOR3B	G3	12.65	<i>C. nievei</i>	Las Planas 4BA	LP4BA	E	—	<i>C. soriae</i>
Toril 3A	TOR3A	G3	12.65	<i>C. nievei</i>	Las Planas 4A	LP4A	E	13.98	<i>C. soriae</i>
Toril 2	TOR2	G3	—	<i>C. nievei</i>	Las Umbrias 16	LUM16	E	14.04	<i>C. soriae</i>
					Las Umbrias 11	LUM11	E	14.06	<i>C. soriae</i>

TABLE 9.1. List of the localities from the Calatayud-Daroca Basin with Cricetodontini. The abbreviations used, local biozone and the species present in each locality are also presented. Calibration according with Daams et al. (1999) and Van Dam et al. (2006) Van Dam et al. (in press).




FIGURE 9.2. Stratigraphical distributions of the species of Cricetodontini in the Calatayud-Daroca Basin. Their correlations with local biozones, the MN system, continental stages and epoch are also presented (Data acquired from López-Guerrero et al., 2008, 2009, 2013a, 2013b, 2014).

CONCLUSIONS

With the studies presented in the chapters from section III (Systematic Paleontology), all the material of Cricetodontini from the Calatayud-Daroca Basin has been taxonomically assigned at species level. The fossils revised in this study belongs to eight species. *Cricetodon soriae* from the local biozone E and *Cricetodon sansaniensis* from the local biozone F have basal traits. *Cricetodon jotae*, found in the local biozones G1, G2 and partly G3; *Cricetodon nievei*, *Hispanomys aguirrei* and *Hispanomys lavocati* from the local biozone G3; display a combination of basal and derivate characters. Finally, *Hispanomys nombrevillae* from the local biozone H and *Hispanomys aragonensis* from the local biozone I are derived representatives from the tribe Cricetodontini.

As is evident from the results described above, the stratigraphical distribution of some Cricetodontini from the Calatayud-Daroca Basin (*C. soriae*, *C. sansaniensis*, *H. nombrevillae* and *H. aragonensis*) demonstrates their suitability as biostratigraphical markers for the Calatayud-Daroca Basin. Moreover, we have been able to precise the distribution of the species of Cricetodontini belonging to the long G3. We have recognized the possibility of redefinition and subdivision of this local biozone. However, it is necessary to complete the study of the rest of fauna before proposing a new biozonation.

LITERATURE CITED

- Agustí, J., L. Cabrera, M. Garcés, W. Krijgsman, O. Oms, and J. M. Parés. 2001. A calibrated mammal scale for the Neogene of Western Europe. State of the art. *Earth-Science Reviews* 52:247–260.
- Álvarez-Sierra, M. A., J. P. Calvo, J. Morales, A. Alonso-Zarza, B. Azanza, I. García Paredes, M. Hernández Fernández, A. J. Van der Meulen, P. Peláez-Campomanes, V. Quiralte, M. J. Salesa, I. M. Sánchez, and D. Soria. 2003. El tránsito Aragoniense-Vallesiense en el área de Daroca-Nombrevilla (Zaragoza, España). *Coloquios de Paleontología Volumen Extraordinario* 1:25–33.
- Álvarez-Sierra, M. A., I. García-Paredes, V. Hernández-Ballarín, L.W. Van den Hoek Ostende, K. Hordijk, P. López-Guerrero, A. J. Van der Meulen, A. Oliver, and P. Peláez-Campomanes, P. 2013a. Biochronology and Historical Biogeography: Testing Symmetric versus Asymmetric Time Models; pp. 244 in N. Çağatay, and C. Zabci (eds.), 14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratehys and Black sea. Book of Abstracts. General Directorate of Mineral Research and Exploration (MTA), Istanbul.
- Álvarez-Sierra, M. A., I. García-Paredes, V. Hernández-Ballarín, L.W. Van den Hoek Ostende, K. Hordijk, P. López-Guerrero, A. J. Van der Meulen, A. Oliver, and P. Peláez-Campomanes, P. 2013b. Models of historical biogeography and continental biochronology. *Spanish Journal of Palaeontology* 28:129–138.
- Böhme, M., J. Prieto, B. Reichenbacher, M. Aiglstorfer, and A. Rocholl. 2012. Pre-Riesian hiatus (Styrian tectonic phase) and minor hiatuses in the NAFB; pp. 2-8 in M. Böhme, J. Prieto, B. Reichenbacher, M. Aiglstorfer, and A. Rocholl (eds.), *The Early Miocene Molasse in southern Germany. Field Trip Guide. RCMNS Workshop May 16th-20th, 2012. Tübingen-Bad Krumbad.*
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.

- De Bruijn, H., and E. Ünay. 1996. On the evolutionary history of the *Cricetodontini* from Europe and Asia Minor and its bearing on the reconstruction of migrations and the continental biotope during the Neogene; pp. 227–234 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Saraç, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a discussion of the evolutionary history of the *Cricetodontini*. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 96:151–216.
- Depéret, C., 1890. Les animaux Pliocènes du Roussillon. *Mémoires de la Société Géologique de France* 1:1–63.
- Domingo, L., P. L. Koch, S. T. Grimes, J. Morales, and N. López-Martínez. 2012. Isotopic paleoecology of mammals and the Middle Miocene Cooling event in the Madrid Basin (Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology* 339:98–113.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 69:298–317.
- Garcés, M., W. Krijgsman, P. Peláez-Campomanes, M. A. Álvarez-Sierra, and R. Daams. 2003. *Hipparion* dispersal in Europe: magnetostratigraphic constraints from the Daroca area (Spain). *Coloquios de Paleontología Volumen Extraordinario* 1:171–178.
- Heißig, K., 1997. Mammal faunas intermediate between the reference faunas of MN4 and MN6 from the Upper freshwater Molasse of Bavaria; pp. 537–546 in J. P., Aguilar, S. Legendre, and J. Michaux (eds.) *Congrès BiochroM '97. Mémoires et Travaux EPHE*, Institut Montpellier, Montpellier.
- Hernández-Ballarín, V., A. Oliver, and P. Peláez-Campomanes. 2011. Revisión de las asociaciones de mamíferos del tránsito Aragoniense medio y superior de la Cuenca de Madrid; pp. 183–172 in A. Pérez-García, F. Gascó, J. M. Gasulla, and F. Escaso (eds.), *Viajando a Mundos Pretéritos*. Ayuntamiento de Morella, Morella, Castellón.
- Hernández Fernández, M., J. A. Cárdena, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López-Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.

- Hilgen, F. J., L. J. Lourens, and J. A. Van Dam. 2012. The Neogene Period; pp. 923–978 in F. M. Gradstein, J. G. Ogg, J. G. Schmitz (eds.), *Geologic Time Scale*. Cambridge University Press, Cambridge.
- Hordijk, K., 2010. Perseverance of pikas in the Miocene. *Geologica Ultraiectina Mededelingen van de Faculteit Geowetenschappen departement Aardwetenschappen, Universiteit Utrecht, Utrecht*, 232 p.
- Jaeger, J. J., J. Michaux, and B. David. 1973. Biochronologie du Miocène moyen et supérieur continental du Maghreb. *Comptes Rendus de l'Académie des Sciences Paris, séries D* 277:2477–2480.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254:177–235.
- Krijgsman, W., M. Garcés, C. G. Langereis, R. Daams, J. Van Dam, A. J. Van der Meulen, J. Agustí, and L. Cabrera. 1996. A new chronology for the middle to late Miocene continental record in Spain. *Earth and Planetary Science Letters* 142:367–380.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d'une récapitulation des diverses espèces d'animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d'autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. *J. A. Portes, Auch*, 41 pp.
- Lear, C. H., H. Elderfield, and P. A. Wilson. 2000. Cenozoic deep-sea temperatures and global ice volumes from Mg/Ca in benthic foraminiferal calcite. *Science* 287:269–272.
- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013a. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89:573–592.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra, and P. Peláez-Campomanes, P. 2013b. Cricetodontini from the Calatayud-Daroca Basin (Miocene, Spain); pp. 240 in N. Çağatay, and C. Zabcı (eds.), 14th RCMNS Congress. Neogene to Quaternary Geological Evolution of Mediterranean, Paratethys and Black sea. Book of Abstracts. General Directorate of Mineral Research and Exploration (MTA), Istanbul.
- López-Guerrero, P., M. A. Álvarez-Sierra, R. López-Antoñanzas, A. Oliver Pérez, and P. Peláez-

- Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). *Palaeontologica Nova* 8:259–271.
- López-Guerrero, P., I. García-Paredes, M. A. Álvarez-Sierra, R. López-Antoñanzas, and A. Oliver. 2009. Cricetodontini (Rodentia, Mammalia) from the upper Aragonian and lower Vallesian of the Toril-Nombrevilla section (Middle and Upper Miocene, Calatayud-Daroca basin, Zaragoza, Spain). *Journal of Vertebrate Paleontology* 29(3, supplement):161A.
- López-Martínez, N., C., Sesé Benito, and J. L. Sáenz-García. 1977. La microfauna (Rod., Insect., Lagomorph y Reptilia) de las fisuras del Mioceno medio de Escobosa de Calatañazor (Soria, España). *Acta Geologica Hispanica* 12:47–73.
- Maridet, O., and S. Sen. 2012. Les Cricetidae du gisement de Sansan; pp. 29–64 in S. Peigné and S. Sen (eds.), *Les mammifères du gisement de Sansan. Mémoires du Muséum national d'Histoire naturelle*, Paris.
- Mein, P., and M. Freudenthal. 1971. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Rögl, F., and G. Daxner-Höck. 1996. Late Miocene Paratethys Correlations; pp. 47–55 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The evolution of western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- Rummel, M. 1999. Tribe Cricetodontini; pp. 359–364 in G. Rößner and K. Heißig (eds.), *The Miocene Land Mammals of Europe*. Verlag Dr. Friedrich Pfeil, München.
- Sen, S. 1996. Present State of Magnetostratigraphic Studies in the Continental Neogene of Europe and Anatolia; pp. 56–63 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The evolution of western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- Shevenell, A., J. P. Kennett, and D. W. Lea. 2004. Middle Miocene Southern Ocean cooling and Antarctic cryosphere expansion. *Science* 305:1766–1770.
- Ünay, E., H. De Bruijn, and G. Saraç. 2003. A preliminary zonation of the continental Neogene of Anatolia based on rodents; pp. 539–547 in J.W.F. Reumer, W. Wessels (eds.), *Distribution and migration of Tertiary Mammals in Eurasia. A volume in honour of Hans de Bruijn*. Deinsea, Rotterdam.
- Ünay, E., and H. De Bruijn. 1984. On some Neogene Rodent assemblages from both sides of the Dardanelles, Turkey. *Newsletter of Stratigraphy* 13:119–132.

- Van Dam, J., W. Krijgsman, H. A. Abels, M. A. Álvarez-Sierra, I. García-Paredes, P. López-Guerrero, P. Peláez-Campomanes y D. Ventura. In press. Updated chronology for middle to late Miocene mammal sites of the Daroca area (Calatayud-Montalbán Basin), Spain. DOI: <http://dx.doi.org/10.1016/j.geobios.2014.07.002>.
- Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. Van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, and P. Peláez-Campomanes. 2012. Updated Aragonian biostratigraphy: Small Mammal distribution and its implications for the Miocene European Chronology. *Geologica Acta* 10:1–24.
- Van der Meulen, A. J., I. García-Paredes, M. A. Álvarez-Sierra, L. W. Van den Hoek Ostende, K. Hordijk, A. Oliver, P. López-Guerrero, V. Hernández-Ballarín, and P. Peláez-Campomanes. 2011. Biostratigraphy or Biochronology? Lessons from the Early and Middle Miocene small Mammal events in Europe. *Geobios* 44:309–321.
- Zachos, J., M. Pagani, L. Sloan, E. Thomas, and K. Billups. 2001. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693.

V. Filogenia

Chapter 10

Phylogeny



FIRST PHYLOGENETIC APPROACH TO THE CRICETODONTINI (RODENTIA, MAMMALIA) FROM THE MIOCENE AND PLIOCENE OF EUROPE AND ASIA MINOR

“Knowledge is not a series of self consistent theories that converges towards an ideal view; it is rather an ever increasing ocean of mutually incompatible (and perhaps even incommensurable) alternatives”.

Paul Karl Feyerabend

Resumen

10

La taxonomía de los géneros de Cricetodontini es controvertida y en muchas ocasiones, se ha discutido ampliamente la adscripción genérica de algunas de sus especies. Además, el establecimiento de las relaciones filogenéticas de Cricetodontini es complicado debido al patrón morfológico en mosaico que muestran algunos de sus representantes. Se ha llevado a cabo un análisis cladístico de 51 especies pertenecientes a cinco géneros de Cricetodontini. En el cladograma consenso de los resultados, *Cricetodon kasapligili* se sitúa como la especie más basal del *in-group*. *Deperetomys* y *Byzantinia* son monofiléticos y *Cricetodon* es parafilético. Además, se necesitan datos suplementarios para poder establecer las relaciones filogenéticas de *Ruscinomys* e *Hispanomys*. Los géneros estudiados presentan la siguiente distribución geográfica: *Deperetomys* estaba presente en Anatolia y Europa Central, *Cricetodon* se distribuyó ampliamente, mientras *Byzantinia* estaba fuertemente relacionado con Anatolia, así como *Ruscinomys* e *Hispanomys* con la Península Ibérica. Se pueden sacar una serie de conclusiones paleobiogeográficas a partir de la observación del cladograma datado. Prestando atención a la distribución geográfica y cronológica de la tribu Cricetodontini se pueden observar tres eventos principales de migración dentro del grupo: el primero (18 Ma) fue aquel en el que *Cricetodon* salió de Anatolia. En el segundo (15 Ma) otra rama de *Cricetodon* migró fuera de Anatolia y se dispersó a través de Europa. En el último evento (14 Ma), la tribu Cricetodontini emigró al suroeste de Europa. Este último evento es coincidente con la primera aparición de los géneros *Hispanomys* y *Ruscinomys* en Europa, de *Byzantinia* en Asia Menor, así como con la última aparición de *Deperetomys* en Europa.



Abstract

The taxonomy of the tribe Cricetodontini is controversial and the generic adscription of some species has been largely discussed. Besides, the establishment of the phylogenetic relationships of Cricetodontini is complicated due to the mosaic evolutionary pattern displayed by some of their representatives. A cladistics analysis involving 51 species of five genera of Cricetodontini has been carried out. *Cricetodon kasapligili* emerges as the most basal species of the in-group. *Deperetomys* and *Byzantinia* are monophyletic, *Cricetodon* is paraphyletic, and the analysis needs of supplemental data to establish the phylogenetic relationships of *Ruscinomys* and *Hispanomys*. The studied genera were geographically distributed as follows: *Deperetomys* was present in Anatolia and Central Europe, *Cricetodon* was widely distributed whereas *Byzantinia* was strongly related with Anatolia, and *Ruscinomys* and *Hispanomys* with the Iberian Peninsula. Based in a dated cladogram a number of paleobiogeographical conclusions can be interpreted. The geographical and chronological distribution of Cricetodontini indicates three main migration events within the group: the first one (18 Ma) took *Cricetodon* out of Anatolia. The second one (15 Ma) took another branch of *Cricetodon* outside Anatolia and dispersed it through Europe. In the last event (14 Ma), the Cricetodontini migrated to south-western Europe; which is coincident with the first occurrence of the genera *Hispanomys* and *Ruscinomys* in Europe and *Byzantinia* in Asia Minor as well as the last occurrence of *Deperetomys* in Europe.

INTRODUCTION

The tribe Cricetodontini includes cricetid rodents found in Europe, Northern Africa, and Asia (Jaeger and Michaux, 1973; De Bruijn and Ünay, 1996; Rummel, 1999). They are well distinguished by the large size of their molars and by the presence of one or two highly variable ectolophs. The Cricetodontini were originated in the early Miocene and the latest known record is of Pliocene age (De Bruijn et al., 1993; García-Alix, 2006).

According to Rummel (1999), the tribe includes eight genera: *Cricetodon* Lartet, 1851; *Hispanomys* Mein and Freudenthal, 1971a; *Ruscinomys* Déperet, 1890; *Byzantinia* De Bruijn, 1976; *Deperetomys* Mein and Freudenthal, 1971a; *Zramys* Jaeger and Michaux, 1973; *Meteamys* De Bruijn and Saraç, 1992; and *Eumyarion* Thaler, 1966. Among them, *Cricetodon*, *Hispanomys*, *Ruscinomys* and *Deperetomys* are the most abundant and studied representatives of the tribe in Europe. *Cricetodon* is replaced by *Byzantinia* in Turkey (De Bruijn et al., 1993), becoming the most common Cricetodontini in Anatolia. Given that these five genera are widely distributed in Europe and Asia Minor and they lumps a high number of species, our analysis will be focused on them.

The taxonomical classification of the Cricetodontini has been largely discussed. For example, *Deperetomys* was assigned to several subfamilies by different authors (Mein and Freudenthal, 1971a; Ünay, 1989; De Bruijn et al., 1993; Maridet and Ni, 2013). Apart from this, as explained in chapter 5, *Cricetodon* firstly included a large number of European which are now included in other cricetid groups. However, recently, Sen and Erbajeva (2011) suggest that the taxonomical revision of *Cricetodon* is still necessary. In addition, as remarked in the section III (Systematic Paleontology) the generic assignment is, sometimes, very complicated because the differences among some Cricetodontini are not clear (De Bruijn, 1976; Sen and Ünay, 1979; Ünay, 1980; Ünay and De Bruijn, 1984; De Bruijn et al., 1993; De Bruijn and Ünay 1996; Casanovas-Vilar, 2007; Prieto et al., 2010, 2014; López-Guerrero et al., 2014).

There are several phylogenetic approaches that deal with the Cricetodontini from southwestern Europe (*Cricetodon*, *Hispanomys* and *Ruscinomys*). Figure 10.1 represents the different phylogenies proposed by various authors based in several criteria, morphology and size basically. Similarly, there are cladistic analyses including some species of Cricetodontini (Sen and Erbajeva, 2011) or *Cricetodon* (Bi, 2005). Unfortunately, the phylogeny made by Bi (2005) is unpublished and it will not be discussed in the present chapter.




FIGURE 10.1. Former phylogenies proposed for the tribe Cricetodontini: (a). Freudenthal, 1966; (b). Mein and Freudenthal, 1971a; (c). Van de Weerd, 1976; (d). Agustí, 1980.

The main aim of this work is to test the phylogenetic relationships of the species from the Calatayud-Daroca Basin presented in chapters 5, 6, 7 and 8. Additionally, the phylogenetic hypotheses on Eurasian Cricetodontini proposed by previous authors will be tested. Another objectives are to reassess the generic ascription of some species and, finally, to describe a general overview of the paleobiogeographical and evolutionary history of the Cricetodontini.

MATERIAL & METHODS

Parimony analyses have been performed to assess the phylogenetic affinities including the whole array of Miocene Cricetodontini of Europe, Anatolia and China. The in-group contains 51 operational taxonomic units (OTU) representing the following genera of Miocene and Pliocene Cricetodontinae: *Cricetodon*, *Hispanomys*, *Ruscinomys*, *Byzantinia* and *Deperetomys*. The rodent *Democricetodon franconicus* (Falhbusch, 1966), considered as a primitive cricetid (Sen and Erbajeva, 2011), has been selected as out-group. The data matrix includes 103 characters considered as homologous (Appendix 10.1). The characters have been mainly based on dental morphology (Fig. 10.2) and size. Data has been collected from fossils and photographs of the type series as well as from literature, resulting in the largest morphometrical dataset used in a Cricetodontinae cladistic analysis to date. Table 10.1 provides all the information about the studied material. Each character has been chosen to fit the criteria established by Gomes-Rodrigues et al. (2009): it is included in the different diagnoses proposed for the studied genera, it is widely used in taxonomy and its variation shows independence between upper and lower teeth. Non-preserved characters (features not observed due to taphonomical problems or not described in the original publications, the information might be included in the future) has been coded as “?” and non-applicable characters (features not found in the OTU, the information will never be included) has been coded as “-”. The data matrix has a very low proportion of missing data: only 9.62% of the total cells contain question marks.

Size has been coded in three states of character (“large”, “medium” and “small” size of the M1). The limits among them are the percentiles (P) calculated for the means of the studied sample. Hence, “large” are those sizes which have means larger than the P75 value. Between P75 and P50 are “medium” ones, and “small” are those smaller than P50. Similar procedure has been used in the codification of the metric indexes. For the length m2/length m3 index, only two codes are used; “less than the P50” and “more than the P50”. These limits are specific for our studied sample. That means that if any further studies include new species or new measurements of the studied ones, they should be recalculated. Table 10.2 presents the percentiles calculated for size and the metric indexes. Within the 103 characters, 56 are binomial and 47 are multistate. Characters scored as having multiple states are interpreted as polymorphic. Multistate characters are unordered and no character is weighted.

Species	Locality	Original reference	Age	References (For the Age)	code
<i>R. europaeus</i>	Perpignan	Depéret, 1890	MN14/15	Rummel, 1999	*
<i>R. schaubi</i>	Los Mansuetos	Crusafont and Villalta, 1956	MN14	Rummel, 1999	-
<i>H. adroveri</i>	Casa del Acero	Agustí, 1986	MN12	Rummel, 1999	-
<i>H. baixasi</i>	Lo Fournas 16-M	Aguilar et al., 2007	MN11	Aguilar et al., 2007	-
<i>H. peralensis</i>	Peralejos C	Van de Weerd, 1976	MN10	Van de Weerd, 1976	*
<i>H. moralesi</i>	Batallones 10	López-Antoñanzas et al., 2010	MN10	López-Antoñanzas et al., 2010	+
<i>H. mediterraneus</i>	Montredon	Aguilar, 1982	MN10	Aguilar, 1982	-
<i>H. daamsi</i>	Can Missert	Agustí et al., 2005	MN9	Gómez Cano et al., 2011	+
<i>H. aragonensis</i>	Pedregueras 2C	Freudenthal, 1966	MN9	Daams et al., 1999	*
<i>H. nombrevillae</i>	Nombrevilla 1	Freudenthal, 1966	MN9	Daams et al., 1999	*
<i>H. thaleri</i>	Can Llobateres	Hartenberger, 1965	MN9	Agustí et al., 1984	-
<i>H. bijugatus</i>	La Grive-Saint Alban, L3	Mein and Freudenthal, 1971a	MN7/8-MN9	López-Antoñanzas and Mein, 2009	+
<i>H. dispectus</i>	Hostalets de Pierola	Agustí, 1980	MN7/8-MN9	Agustí, 1980	+
<i>H. decedens</i>	La Grive-Saint Alban, L5'	Schaub, 1925	MN7/8	López-Antoñanzas and Mein, 2009	+
<i>H. lavocati</i>	Hostalets de Pierola	Freudenthal, 1966	MN7/8	Agustí, 1980	*
<i>H. aguirrei</i>	Escobosa de Calatañazor	López-Martínez et al., 1977	MN7/8	López-Martínez et al., 1977	*
<i>H. castelnovi</i>	Castelnou 6	Aguilar et al., 1994	MN6?	Aguilar et al., 1994	-
<i>D. hangi</i>	Giggenhausen	Fahlbusch, 1964	MN7/8	De Bruijn et al., 1993	+
<i>D. intermedius</i>	Harami 1	De Bruijn et al., 1987	MN2	De Bruijn et al., 1993	+
<i>D. anatolicus</i>	Kılçak 0"	De Bruijn et al., 1993	MN1	De Bruijn et al., 1993	+
<i>'C' fandli</i>	Gratkorn	Prieto et al., 2010	MN7/8	Prieto et al., 2010	*
<i>'C'. klariankae</i>	Felsőtárkány-Felnémet	Hír, 2007	MN7/8	Hír, 2007	*
<i>C. jumaensis</i>	Petersbuch 18	Rummel, 2001	MN7/8	Rummel, 2001	-
<i>C. bolligeri</i>	Petersbuch 10	Rummel, 1995	MN7/8	Rummel, 1995	*
<i>C. cariensis</i>	Sarıçay	Sen and Ünay, 1979	MN7/8	Sen and Ünay, 1979	-
<i>C. albanensis</i>	La Grive-Saint Alban	Mein and Freudenthal, 1971a	MN7/8	Mein and Freudenthal, 1971a	*
<i>C. nievei</i>	Toril 3A	López-Guerrero et al., 2014	MN7/8	Álvarez-Sierra et al., 2003	*
<i>C. engesseri</i>	Chräzerentobl 505m	Rummel and Kálin, 2003	MN6-MN7/8	Rummel and Kálin, 2003	-
<i>C. sansaniensis</i>	Sansan	Schaub, 1925	MN6	De Bruijn et al., 1992	*
<i>C. pasalarensis</i>	Pasalar	Tobien, 1978	late MN6	Peláez-Campomanes and Daams, 2002	-
<i>C. jotae</i>	Manchones	Mein and Freudenthal, 1971a	MN6, biozone	De Bruijn et al., 1992	*
<i>C. hungaricus</i>	Hasznos	Kordos, 1986	MN6	De Bruijn et al., 1993	-
<i>C. caucasicus</i>	Belometchetskaya	Argyropulo, 1938	late MN6	De Bruijn et al., 1992	-
<i>C. soriae</i>	Somosaguas	Hernández-Fernández et al., 2006	MN5, biozone E	Hernández-Fernández et al., 2006	*
<i>C. meini</i>	Vieux-Collonges	Freudenthal, 1963	MN5	Mein and Freudenthal, 1971b	*
<i>C. candirensis</i>	Candir	Tobien, 1978	MN5	De Bruijn et al., 2003	-
<i>C. aureus</i>	Vieux-Collonges	Mein and Freudenthal, 1971b	MN5	Mein and Freudenthal, 1971b	*
<i>C. volkeri</i>	Songlinzhuan	Wu et al., 2009	early Miocene	Wu et al., 2009	-
<i>C. aliveriensis</i>	Aliveri	Klein Hofmeijer and de Bruijn, 1989	MN4	Klein Hofmeijer and de Bruijn, 1988	+
<i>C. wanhei</i>	Dingshanyanchi	Qiu, 2010	middle Miocene	Qiu, 2010	-
<i>C. tobieni</i>	Horlak 1a	De Bruijn et al., 1993	MN3-MN4	Tobien, 1978	-
<i>C. kasapligilli</i>	Keseköy	De Bruijn et al., 1993	MN3	De Bruijn et al., 1993	+
<i>C. versteegi</i>	Kılçak 3a	De Bruijn et al., 1993	MN1	De Bruijn et al., 1993	+
<i>B. pikermiensis</i>	Pikermi (Chomateri)	De Bruijn, 1976	MN12	Rummel, 1999	-
<i>B. hellenicus</i>	Samos Island	Freudenthal, 1970	MN11	Kostopoulos, 2009	-
<i>B. nikosi</i>	Biodrak	De Bruijn, 1976	Vallesian	Rummel, 1999	-
<i>B. dardanellensis</i>	Bayraktepe II	Ünay, 1980	middle Vallesian	Rummel, 1999	-
<i>B. ozansoyi</i>	Bayraktepe I	Ünay, 1980	late aragonian	Rummel, 1999	-
<i>B. sofçaensis</i>	Sofça	Tobien, 1978	MN8	Rummel, 1999	-
<i>B. bayraktepensis</i>	Bayraktepe I	Ünay, 1980	late aragonian	Rummel, 1999	-
<i>B. eskihissarensis</i>	Yeni Eskihsar	Tobien, 1978	middle Miocene	Rummel, 1998	-

TABLE 10.1. List of the species included in the in-group of the cladistic analysis. The observation code is present in the sixth column: *: personal observation; +: data acquired from recent publications with accurate and detailed descriptions and good quality photographs, and -: data taken from vague descriptions or only one photograph.




Figure 10.2. Terminology of the parts of the cheek teeth of Cricetodontini. Only the morphology of the upper molars and the m1 has been drawn; however, this nomenclature can be applied for the rest of the lower molars. (Modified after López-Guerrero et al., 2013.)

Character states have been defined as follow:

1. Size - Length of the M1: 0 = less than 3.2mm, 1 = between 3.2mm and 3.38mm, 2 = more than 3.38mm.
2. Groove of the upper incisor: 0 = absent, 1 = present.
3. Hypsodonty: 0 = low, 1 = high.
4. Shape of the cusps: 0 = 'inflated', 1 = slender, 2 = fused with the ectolophs.
5. Sinuous ectolophs: 0 = absent, 1 = present.
6. Contour: 0 = straight, 1 = sinuous.
7. Wear: 0 = plain, 1 = concave.

8. Upper molars: Enamel-coated valley: 0 = present, 1 = absent.
9. M1 - Anterocone: 0 = simple/undivided, 1 = slightly divided, 2 = strongly divided.
10. M1 - Anterior valley between the lobes of the anterocone: 0 = absent, 1 = present.
11. M1 - Lingual spur of the anterolophule: 0 = absent, 1 = present.
12. M1 - Protocone spur: 0 = mostly absent, 1 = present.
13. M1 - Labial spur of the anterolophule: 0 = present at least in some specimens, 1 = always absent.
14. M1 - Lingual anteroloph: 0 = mostly absent, 1 = absent.
15. M1 - Protostyle: 0 = absent, 1 = present in some specimens, 2 = always present.
16. M1 - Protosinus: 0 = at least partially closed, 1 = open, 2 = absent.
17. M1 - Anterior ectoloph: Posterior ectoloph of the anterocone: 0 = absent in some specimens, 1 = present, 2 = present and curved towards the anterosinus.
18. M1 - Anterior ectoloph: Anterior ectoloph of the paracone: 0 = absent at least in some specimens, 1 = present.
19. M1 - Anterior ectoloph: Connection: 0 = absent, 1 = present in some specimens.
20. M1 - Posterior ectoloph: Anterior ectoloph of the metacone: 0 = absent in some specimens, 1 = always present.
21. M1 - Posterior ectoloph: Posterior ectoloph of the paracone: 0 = absent in some specimens, 1 = always present, 2 = present and curved towards the mesosinus.
22. M1 - Posterior ectoloph: Connection: 0 = always incomplete, 1 = complete in some specimens, 2 = connected through the long mesoloph.
23. M1 - Anterolingual cingulum: 0 = absent, 1 = present in some specimens, 2 = always present.
24. M1 - Labial posteroloph: 0 = well developed, 1 = poorly developed, 2 = absent.
25. M1 - Anterosinus closed by a cingulum: 0 = present in more than 50% of the specimens, 1 = present in less than 50% of the specimens, 2 = always absent.
26. M1 - Mesosinus closed by a cingulum: 0 = present in more 50% of the specimens, 1 = present in less than 50% of the specimens, 2 = always absent.
27. M1 - Sinus closed by a cingulum: 0 = mostly complete, 1 = mostly incomplete, 2 = always absent.
28. M1 - Protolophule: 0 = anterior present in some specimens, 1 = double, anterior and posterior always present, 2 = only the posterior is present.
29. M1 - Sinus direction: 0 = anterior, 1 = transverse, 2 = posterior.
30. M1 - Style on the sinus: 0 = present at least in some specimens, 1 = always absent.
31. M1 - Entomesoloph: 0 = present at least in some specimens, 1 = always absent.
32. M1 - Mesoloph: 0 = long, 1 = short, 2 = absent.
33. M1 - Lingual posteroloph: 0 = well developed, 1 = poorly developed, 2 = absent.
34. M1 - Roots: 0 = three, 1 = four, 2 = five.

35. M2 - Lingual anteroloph: 0 = well developed, 1 = short, 2 = absent.
36. M2 - Anterior ectoloph: 0 = absent, 1 = present in some specimens, 2 = always present.

37. M2 - Anterior ectoloph: Connection: 0 = absent, 1 = present in some specimens, 2 = always connected.
38. M2 - Posterior ectoloph: Anterior ectoloph of the metacone: 0 = always absent, 1 = present in some specimens, 2 = always present.
39. M2 - Posterior ectoloph: Posterior ectoloph of the paracone: 0 = absent in some specimens, 1 = always present.
40. M2 - Posterior ectoloph: Connection: 0 = always incomplete, 1 = complete in some specimens or always complete, 2 = connected through the mesoloph.
41. M2 - Anterosinus closed by a cingulum: 0 = present, 1 = connected through the mesoloph, 2 = absent.
42. M2 - Mesosinus closed by a cingulum: 0 = present, 1 = narrow platform + low cingul, 2 = absent.
43. M2 - Sinus closed by a cingulum: 0 = absent, 1 = present.
44. M2 - Protocone spur: 0 = mostly absent, 1 = present.
45. M2 - Style on the sinus: 0 = absent, 1 = present in some specimens, 2 = always present.
46. M2 - Entomesoloph: 0 = present in some specimens, 1 = always absent.
47. M2 - Mesoloph: 0 = absent, 1 = present, 2 = long.
48. M2 - Lingual posteroloph: 0 = well developed, 1 = poorly developed, 2 = absent.

49. M3 - Lingual anteroloph: 0 = present, 1 = weak, 2 = absent.
50. M3 - Labial anteroloph: 0 = absent, 1 = present in some specimens, 2 = always present.
51. M3 - Anterior ectoloph: Anterior ectoloph of the paracone: 0 = absent, 1 = present in some specimens, 2 = always present, 3 = it is certainly the posterior ectoloph of the anteroloph.
52. M3 - Anterior ectoloph: Connection: 0 = always absent, 1 = present.
53. M3 - Posterior ectoloph: Anterior ectoloph of the metacone: 0 = absent, 1 = present in some specimens.
54. M3 - Posterior ectoloph: Posterior ectoloph of the paracone: 0 = absent in some specimens, 1 = always present.
55. M3 - Posterior ectoloph of the M3 curved: 0 = absent, 1 = present.
56. M3 - Posterior ectoloph: Connection: 0 = never connected, 1 = connected in at least one specimen.
57. M3 - Labial cingulum: 0 = present, 1 = absent.
58. M3 - Lingual cingulum: 0 = complete, 1 = incomplete, 2 = absent.
59. M3 - Neo-entoloph: 0 = present in some specimens, 1 = always absent.
60. M3 - Mesoloph: 0 = long, 1 = short, 2 = absent.
61. M3 - Metaloph: 0 = anterior, 1 = transversal, 2 = posterior.
62. M3 - Axioloph: 0 = continuous, 1 = interrupted.
63. M3 - Posteroloph: 0 = present, 1 = absent.
64. M3 - Posterosinus: 0 = present, 1 = absent.
65. M3 - Length M1/ Length M3: 0 = less than 1.62, 1 = between 1.62 and 1.78, 2 = more than 1.78.

66. m1 - Situation of the anteroconid: 0 = in the middle of the longitudinal axis, 1 = labially displaced.
67. m1 - Shape of the anteroconid: 0 = elongated, 1 = rounded.
68. m1 - Labial anterolophid: 0 = absent, 1 = present in some specimens and short, 2 = present in some specimens and well developed.
69. m1 - Lingual anterolophid: 0 = present, 1 = absent.
70. m1 - Spur of the anterolophulid: 0 = present, 1 = absent.
71. m1 - Metalophulid I, solely: 0 = absent, 1 = interrupted, 2 = present.
72. m1 - Metalophulid II, solely: 0 = absent, 1 = present, 2 = transversally displace.
73. m1 - Metalophulid I and II together: 0 = absent, 1 = present.
74. m1 - Mesosinusid closed by a cingulid: 0 = absent, 1 = frequently present (more than 50%), 2 = always present.
75. m1 - Mesolophid: 0 = long, 1 = short, 2 = absent.
76. m1 - Ectomesolophid: 0 = absent, 1 = interrupted, 2 = present.

77. m1 - Sinusid: 0 = open, 1 = closed.
78. m1 - Spur on the entoconid and metaconid: 0 = both always absent, 1 = both present in some specimens.
79. m1 - Hypolophulid connected with the mesolophid: 0 = absent, 1 = present.
80. m1 - Protosinusid: 0 = absent, 1 = present.
81. m1 - Styloid on the sinus: 0 = present in some specimens, 1 = absent.
82. m1 - Protostylid: 0 = present in some specimens, 1 = absent.
83. m2 - Labial anterolophid: 0 = present, 1 = absent.
84. m2 - Lingual anterolophid: 0 = present in some specimens, 1 = absent.
85. m2 - Mesolophid: 0 = long, 1 = short, 2 = absent, 3 = connected.
86. m2 - Mesosinusid closed by a cingulid: 0 = absent, 1 = present.
87. m2 - Sinusid: 0 = closed, 1 = open.
88. m2 - Hypolophulid connected with the mesolophid: 0 = absent, 1 = present.
89. m2 - Protosinusid: 0 = well developed, 1 = poorly developed or absent.
90. m2 - Styloid on the sinus: 0 = present in some specimens, 1 = always absent.
91. m2 - Ectomesolophid: 0 = present, 1 = absent.
92. m2 - Posterolophid: 0 = long, 1 = short.
93. m2 - Number of roots: 0 = two, 1 = three.
94. m3 - Labial anterolophid: 0 = present, 1 = absent.
95. m3 - Lingual anterolophid: 0 = absent, 1 = present in some specimens, 2 = always present.
96. m3 - Metalophid: 0 = posterior, 1 = double, 2 = anterior.
97. m3 - Mesolophid: 0 = long, 1 = short, 2 = absent, 3 = connected.
98. m3 - Hypolophulid connected with the mesoloph: 0 = absent, 1 = present.
99. m3 - Mesosinusid closed by a cingulid: 0 = absent, 1 = present in more or less the 50% of the specimens, 2 = always present.
100. m3 - Protosinusid: 0 = absent, 1 = partially closed, 2 = closed.
101. m3 - Styloid on the sinus: 0 = present, 1 = absent.
102. Length m1/length m3: 0 = less than 1.10, 1 = between 1.10 and 1.14, 2 = more than 1.14.
103. Length m2/length m3: 0 = less than 1.06, 1 = more than 1.06.

A parsimony analysis was conducted with TNT software (Golobof et al., 2008). A traditional search using Wagner trees (1000 replicates, TBR algorithm) was performed. Bremer support values were calculated as a measure of node robustness. This index examines how many additional steps are required for losing a branch in the consensus cladogram (Morrone, 2000; Bremer, 1988; 1994).

	LM1	LM1/LM3	Lm1/Lm3	Lm2/Lm3
Min	2.01	1.32	0.94	0.94
P50	3.20	1.62	1.10	1.06
P75	3.38	1.78	1.14	—
Max	5.75	2.53	1.28	1.23

TABLE 10.2. Values of the percentiles calculated for the size and indexes of the studied sample. Based on the mean length of each locality. Abbreviations: Min, minimum; P50, percentile 50; P75, percentile 75; Max, maximum; L, length.

RESULTS & DISCUSSION

RESULTS

A traditional search have recovered four most-parsimonious trees (MPT), with a total length of 773 steps (Consistency index (CI): 0.194; Retention index (RI): 0.578) and five clades (Fig. 10.3). Our phylogenetic analysis explores the monophyly and composition of Cricetodontini, testing the hypotheses of relationships proposed by previous authors (mentioned in chapters 5, 6, 7 and 8, and some of the illustrated in figure 10.1). Present work does not intend to be a complete phylogenetic study of the whole tribe Cricetodontini, because we have just focused on several taxa that have been historically controversial.

The strict consensus presents two politomies involving the Node J and *Byzantinia* remained unresolved (Fig. 10.3). The distributions of the character states for the internal and terminal nodes are detailed in figures 10.4, 10.5 and 10.6.

In our phylogenetic analysis, three main groups clearly emerge within Cricetodontini. One of the genera analysed, *Cricetodon*, results clearly paraphyletic. The three groups are distributed in two main clades (Node C) and a group of basal species.

The species *Cricetodon kasapligili* and *C. versteegi* are basal, forming successive sister species to the two main clades (Node C).

The Node A is composed by nine OTUs belonging to *Cricetodon* and *Deperetomys*. *Cricetodon aliverensis*, *C. aureus*, *C. jumaensis* and *C. engesseri*, are successive sister species to the Node D. *Cricetodon nievei* is sister species to *C. bolligeri* (Node B) which is diagnosed by the following synapomorphies: presence of anterolingual cingulum; entomesoloph present in some M1 and M2, and absence of posteroloph. Node B is sister group to clade *Deperetomys* (Node E)

The Node F contains 32 species. The basal positions within Node F are occupied by three successive sister species to Node J. Node J displays a basal politomy composed by *Cricetodon pasalarensis*, Node I and Node K.

Node I is defined by: *Cricetodon soriae*, *C. sansaniensis*, *C. meini* and *C. jotae*.

Node K is composed by four species (*Cricetodon albanensis*, *C. hungaricus*, *C. candirensis* and *C. cariensis*) forming successive sister species to Node O. Clade O is composed by two main clades,

C. caucasicus + *Byzantinia* (Node M) and '*Cricetodon*' *fandli* + *Hispanomys* + *Ruscinomys* (Node P).

DISCUSSION

The systematics of Cricetodontini are controversial (Rummel, 1999; Prieto et al., 2010; Sen and Erbajeva, 2011; López-Guerrero et al., 2013). Within the five studied genera, *Cricetodon* Lartet, 1851 is the first described. As discussed in chapter 5, *Cricetodon* originally included some species that are now ascribed to other cricetid genera. However, in the cladistic analysis performed by Sen and Erbajeva (2011), *Cricetodon* was paraphyletic and these authors questioned whether the content of *Cricetodon* is homogeneous. They pointed out that *Cricetodon* lumps 20 species and its chronostratigraphic range covered the entire early and middle Miocene, with dental patterns differing greatly from one species to another. Thus, they suggested that a revision of the species of *Cricetodon* was still necessary. Our results, that also point out that *Cricetodon* is paraphyletic, are in agreement with those of Sen and Erbajeva (2011). This problem will be further addressed in this chapter.

In Node A *Cricetodon aliverensis*, *C. aureus*, *C. jumaensis* and *C. engesseri* are successive sister species to the Node D, which contains Node B (*C. nievei* + *C. bolligeri*) and Node E (*Deperetomys anatolicus*, *D. intermedius* and *D. hagni*). The support of Node A is weak, the synapomorphies that diagnose it—presence of a protocone spur in the M1; well-developed lingual anteroloph in the M2; anterior ectoloph present in some M2; style present in the sinus of some M2 and the length m1/length m3 index between 1.10 and 1.14—are important characters because they show low intraspecific variability; however, they suffer reversions and are also present in other Cricetodontini. Therefore, we cannot fully confirm the phylogenetic relationship between *Deperetomys* and the species of *Cricetodon* from Node A. Other authors (Ünay, 1989; De Bruijn et al., 1993; Maridet and Ni, 2013) related *Deperetomys* with *Eumyarion*, *Meteamys*, and *Mirrabella*, from Asia. Consequently, we consider that the addition of these genera into the analysis will help to understand the relationships within Node A.

The species of Node F share the presence of entomesoloph in some M2 and a weak lingual anterolophid in the m3. In our MPT, *Cricetodon tobieni* is the most basal species in Node F, together with *C. volkeri* and *C. wanhei*. These OTUs are successive sister groups to the rest of the clade F.

Node J is a polytomy involving *C. pasalarensis*, Node I and Node K. The original trees offer two possible options of *C. pasalarensis*, either at the base of the Node I (Fig. 10.3a) or as sister group to Node K (Fig. 10.3b). Unfortunately, we do not have enough criteria to support none of them and the position of *Cricetodon pasalarensis* remains uncertain.

The monophyly of Node I (*C. sansaniensis*, *C. soriae*, *C. meini* and *C. jotae*) is supported by the following synapomorphies: labial spur of the anterolophule always absent in the M1; protocone spur always present in the M2; entomesoloph on the M2 always absent; the lingual posteroloph in the M2 poorly developed; presence of two crests on the lingual cingulum of the M3 and absence of cingulid closing the mesosinusid in the m3. In chapter 5, we reported that *C. soriae* may be related to the Anatolian species of *Cricetodon* based on the presence of common features (De Bruijn et al., 1993; López-Guerrero et al., 2013). Our MPT rejects this hypothesis as *C. soriae* is placed in a basal position in the Node I. The results of our analysis indicate that *Cricetodon soriae* is more closely related to the species of *Cricetodon* from Europe than to the Anatolian ones.

Node K include species of *Cricetodon*, *Byzantinia*, *Hispanomys* and *Ruscinomys*. Several authors (Ünay, 1980; Aguilar, 1982) proposed that *Hispanomys* was related with the Turkish genus *Byzantinia*, most likely sharing a common ancestor from Eastern Europe during the middle Miocene. Our phylogenetic analysis shows that *Byzantinia* and *Hispanomys* are sister groups. Besides, the Turkish species *Cricetodon cariensis* is basal to *Byzantinia* and *Hispanomys*, suggesting a common Asian origin for these genera. The monophyly of *C. caucasicus* + *Byzantinia* (Node M) is supported by: high hypsodonty; absence of protostyle in M1; anterior ectoloph of the metacone always present in the M1; absence of cingulum in the mesosinus of the M1; weak lingual anteroloph in the M3 and sinusid closed in the m1. Although these are important features, they are also very variable among Cricetodontini.

The clade *Byzantinia* is defined as the one comprising *B. sofçaensis* + *B. hellenicus*, their most recent common ancestor and all its descendants. *Byzantinia* (Node N) is diagnosed by: concave wear; the presence of an anterior valley between the two lobes of the anterocone in the M1, the absence of lingual posterolophid in the m2, the labially situated anteroconid in the m1 and the closed protosinusid in the m3.

The monophyly of '*Cricetodon*' *fandli* + *Hispanomys* + *Ruscinomys* (Node P) is supported by four synapomorphies: absence of enamel-coated valley; presence of style on the sinus of some M1 and mesoloph absent in the M1 but present in the M2. This clade is the sister group to *C. caucasicus* + *Byzantinia*.

The monophyly of *H. lavocati*, *H. aguirrei*, *H. decedens* + *H. bijugatus* (Node Q) is supported by: entomesoloph present in some M1 and anterior ectoloph present in some M2. The clade including the rest of *Hispanomys* and *Ruscinomys* (Node S) is defined by: protostyle and anterior protolophule present in some M1; absent anterosinus and well-developed lingual posteroloph in the M2, and lingual anteroloph absent and anterior ectoloph of the paracone present in some M3. Node S is supported by six synapomorphies, whereas there are two in Node Q. This points out the more conservative character of the species *H. lavocati*, *H. aguirrei*, *H. decedens* and *H. bijugatus* in

contrast to the rest of the species of *Hispanomys* and *Ruscinomys*.

'*Cricetodon*' *klariankae* is placed as sister group of *H. daamsi* (Node T). This relationship is supported by the length of the M1 larger than 3.38mm; protostyle always present, open protosinus and sinus without style in the M1 and short posterolophid in the m2. The monophyly of Node R is supported by: absence of the lingual spur of the anterolophule; presence of complete anterior ectoloph and protocone spur in the M2; absence of mesoloph in the M2; presence of a stylid in the mesosinus of some m1 and length m1/length m3 index higher than 1.14. '*Cricetodon*' *klariankae* was related with '*C. fandli*' and also with *H. bijugatus* (Prieto et al., 2010, 2014). The results of our analysis reject these hypotheses.

Hispanomys and *Ruscinomys* display a similar morphology and some species were ascribed to both genera (e.g., *H. thaleri*). The species of *Ruscinomys* are placed with several derived species of *Hispanomys* (Fig. 10.3) showing that the monophyly of *Hispanomys* is highly dubious. The analysis needs the addition of the remaining species of *Ruscinomys* and supplemental morphological data, especially features present in *Ruscinomys*, to assess the possible synonymy with *Hispanomys*. Therefore, we decided to maintain the status of both genera.

In general, the synapomorphies present in the main nodes from our MPT reveal changes in the morphological pattern (Fig. 10.7). The basal forms (Fig. 10.7, labels 1-3) show well-developed transversal structures (mesoloph, entomesoloph, posteroloph, protocone spur, mesolophid and ectomesolophid). The cingular formations are present and therefore, the contour is straight. These basal forms also present a non-reduced third molar. In the more derived forms, (Fig. 10.7, labels 4-6) the pattern generally becomes simpler (the cingular formations and stylids are less frequent) and the transversal lophs (ectolophs) more developed. The third molar starts to reduce its length. In the derived forms, the cingular formations and transversal lophs disappear, so the contour is sinuous. Additionally, the longitudinal lophs are completely formed and the general pattern is selenodont-like with high hypsodonty.

CRICETODONTINI FROM THE CALATAYUD-DAROCA BASIN

The relationships of the Cricetodontini from the Calatayud-Daroca Basin and the hypotheses previously stated (Fig. 10.1) will be discussed. In our MPT, *Cricetodon soriae* is the most basal taxa of Node I and it is closely related with the other southwestern Europe species *C. sansaniensis*, *C. meini* and *C. jotae*. As discussed above, our analysis rejects the hypothesis discussed in chapter 5 which, according to De Bruijn et al. (1993), states that *C. soriae* is closely related to the species from Asia Minor.

Cricetodon sansaniensis was included by some authors (Mein and Freudenthal, 1971a; Maridet and Sen, 2012) in the anagenetic lineage *C. aureus*–*C. sansaniensis*–*C. albanensis*. In our MPT, *C. sansaniensis* is a member of Node I, whereas *Cricetodon aureus* is placed in Node A. Therefore, *C. aureus* and *C. sansaniensis* are not directly related.

According to the opinion of Mein and Freudenthal (1971a), *Cricetodon jotae* is related to *C. meini*. Our results support this relationship because *C. jotae* and *C. meini* are sister species. *Cricetodon jotae* is the most derived species of Node I and also the most recent member of the clade (late Aragonian) (Daams et al., 1999; López-Guerrero et al., in press). Its main morphological innovations lie almost exclusively in the M1 and M2 (Fig. 10.5). Thus, in this clade, the most derived morphology is that of the upper molars of *C. jotae*.

As discussed in chapter 7, *C. nievei* shares more features with the species from the central Europe basins than with those from the Calatayud-Daroca Basin (López-Guerrero et al., 2014). In our MTP, *C. nievei* is sister group to *C. bollikeri* (endemic from central Europe) and both are grouped with other central European species (*C. engesseri*, *C. jumaensis* and *D. hagni*), which agrees with the phylogenetic relationship discussed before.

Hispanomys aguirrei is situated in Node P among the other species of *Hispanomys* (Fig. 10.3) indicating that its assignation to *Hispanomys* is correct. As discussed in chapters 6, 7 and 8, *C. jotae* was pointed as the ancestor of *H. aguirrei* (Fig. 10.1d). However, the mosaic morphological patterns described for both species hindered the evaluation of their phylogenetic relationships. Our MPT rejects this association, since both species are placed separately. *Cricetodon jotae* is sister species to *C. meini* (Node I) and *Hispanomys aguirrei* is sister species to *H. lavocati* within Node Q (Fig. 10.3). As detailed in chapter 8, *Hispanomys lavocati* shows a general pattern more derived than *H. aguirrei* (Fig. 10.6) and they could be related, our MPT support this relationship.

The generic assignation and phylogeny of *H. lavocati* were discussed by many authors (See Chapter 8). *Hispanomys lavocati* is situated in the strict consensus (Fig. 10.3) as a member of *Hispanomys* closely related with the other southwestern Europe *Hispanomys* from the late Aragonian. This supports its ascription to *Hispanomys*.

There are two phylogenetic hypotheses proposed for *H. lavocati*. Some authors (Mein and Freudenthal, 1971a; Agustí, 1982) related *H. lavocati* with the southwestern European species *Cricetodon albanensis* or *Cricetodon* cf. *sansaniensis* from Armantes 7 (Fig. 10.1a; 10.1b; 10.1d). De Bruijn et al. (1993) and De Bruijn and Ünay (1996) noted that *H. lavocati* shares several dental characteristics with the Asiatic representatives of Cricetodontini and they interpreted this species as a late Miocene immigrant from Asia Minor. Our MPT rejects both phylogenetic hypotheses since *H. lavocati* appears closely related to other species of *Hispanomys* from southwestern Europe.

The phylogeny of *Hispanomys nombrevillae* was studied by many authors (Mein and Freudenthal, 1971a; Agustí, 1982) and is discussed in chapter 8. It was related with *H. bijugatus* (Fig. 10.1b) and also considered as the descendant of *H. aguirrei* (Fig. 10.1d). In our MPT *H. nombrevillae* is sister species to Node U, which includes the rest of derived *Hispanomys*. According to this, *Hispanomys nombrevillae* shares more features with *H. aragonensis* than with *H. aguirrei*/*H. bijugatus*.

Hispanomys aragonensis was reported as the descendant of the French species *H. decedens* (Mein and Freudenthal, 1971a; Agustí, 1982; López-Antoñanzas and Mein, 2011; Fig. 10.1b, 1d) and also, as the ancestor of *H. peralensis* (Fig. 10.1c). *Hispanomys decedens* possesses some features (ectolophs less developed, and both metalophulids I and II together in the same m1) that makes it a plausible ancestor of *H. aragonensis*. However, in our MPT, these two species are situated in separate clades (Node Q and Node U), rejecting the relationship between *H. decedens* and *H. aragonensis*. According to our MPT, *H. aragonensis* is the most basal OTU to Node V, which includes *H. peralensis*. Van de Weerd (1976) stated that *H. peralensis* could be the descendant of *H. aragonensis* because it possesses more derived features such as: five roots on the M1; highly hypsodont teeth; the labial anteroloph less developed and absence of metalophulid II. Our phylogenetic analysis supports this relationship.

COMMENTS ON SYSTEMATICS

As concluded in the chapters of section III (Systematic Paleontology), there are some open questions regarding the taxonomical status of several Cricetodontini found in the Calatayud-Daroca Basin. Similar problems have been noted in some species in central Europe (Prieto et al., 2010, 2014 and references therein). Based on the results of our phylogenetic analysis (Fig. 10.6) and the detailed observation of several Cricetodontini, we discuss below the taxonomy of some taxa.

Sen and Erbajeva (2011) suggested that *Cricetodon versteegi* and *C. kasapligili* have a simple pattern shared with the other “primitive” genera *Eucricetodon*, *Democricetodon* and *Spanocricetodon*. The position of *Cricetodon versteegi* and *C. kasapligili* in our MPT indicates that they should not be included in the genus *Cricetodon*, rather they appear to represent a paraphyletic array of stem taxa. Therefore, due to their primitive morphology and their basal position, we classified them as Cricetodontinae *incertae sedis*.

The inclusion of *Deperetomys* in the tribe Cricetodontini was discussed before (Ünay, 1989; De Bruijn et al., 1993; Maridet and Ni, 2013). In our MTP, the clade *Deperetomys* possesses some synapomorphies very exclusive and well supported (Fig. 10.3). These are: slender cusps; posterior ectoloph of the anterocone in the M1 present and curved towards the mesosinus; anterior ectoloph

of the metacone always present in the M1; connection of the posterior ectoloph in the M1 through the mesoloph; long mesoloph in the M1; anterior ectoloph of the metacone present in some M2; lingual posteroloph absent in the M2; hypolophulid connected with the mesolophid in the lower molars; elongated anteroconid, well-developed lingual anterolophid, long mesolophid and closed sinusid in the m1 and closed protosinusid in the m3.

This supports the hypotheses that consider *Deperetomys* very different from the rest of Cricetodontini. However, *Deperetomys* is placed, in our MTP, among other representatives of Cricetodontini, thus we maintain its inclusion in this tribe. Nevertheless, it is recommended to include the genera *Meteamys*, *Mirrabella* (subfamily Cricetopinae) and *Eumyarion* (subfamily Cricetodontinae) in future analysis to properly assess the suprageneric status of *Deperetomys*.

Node I contains the type species *Cricetodon sansaniensis* and we consider the species included in it as the unequivocal members of *Cricetodon*. According to the strict consensus (Fig. 10.3) it would seem that any other species of *Cricetodon* placed elsewhere in our MPT should not be assigned to *Cricetodon*. However, we consider that a proper assessment of this affirmation would require the inclusion of the older genera of Cricetodontini: *Zramys*, *Meteamys* and *Eumyarion*. In consequence, we resolved to maintain the generic ascription to *Cricetodon* as they currently are.

Cricetodon hungaricus, *C. candirensis* and *C. cariensis* are successive sister groups to Node O. De Bruijn et al. (1993) suggested a synonymy between *C. hungaricus* and *C. candirensis*. Besides that, *Cricetodon candirensis* and *C. cariensis* are species from Asia Minor originally described as *Turkomys* Tobien, 1978. This genus was synonymised with *Cricetodon* by De Bruijn et al. (1993). The paraphyletic position of these three taxa suggests that they do not belong to *Cricetodon*. The validity of the genus *Turkomys* should be therefore re-considered. Unfortunately, the original description of *Turkomys* made by Tobien (1978) is vague and we have not had the opportunity to study the material of these species, so we lack the information to support any taxonomical opinion.

The monophyly of *C. caucasicus* + *Byzantinia* (Node M) is supported by important features which are also very variable characters among Cricetodontini (see discussion). We consider that the synapomorphies present in Node N are the ones that characterize *Byzantinia*: concave wear; presence of an anterior valley between the two lobes of the anterocone in the M1; absence of lingual posterolophid in the m2; labially situated anteroconid in the m1 and closed protosinusid in the m3. Therefore we do not consider *C. caucasicus* as member of the genus *Byzantinia* maintaining its inclusion in *Cricetodon*.

In the clade '*Cricetodon*' *fandli* + *Hispanomys* + *Ruscinomys*, '*C. fandli*' stands out for being the most basal off-shoot of Node P. Prieto et al. (2010) found affinities with both *Cricetodon* and *Hispanomys* and decided not to include it in any genus (Prieto et al., 2010; 2014). In our MPT, the position of '*C. fandli*' is clearly basal to clade *Hispanomys* + *Ruscinomys*. This could lead us to assign

'*C.*' *fandli* to *Hispanomys*. However, the monophyly of '*Cricetodon*' *fandli* + *Hispanomys* + *Ruscinomys* (Node P) is weakly supported and we prefer not to change the taxonomical assignment of the species '*C.*' *fandli* and recommend a taxonomical study.

'*Cricetodon*' *klariankae* stands out for being placed among the species of *Hispanomys*. Hír (2007) pointed out that '*C.*' *klariankae* differs from *Hispanomys* by its well-developed anterolophulid and metalophulid II (which is always present with metalophulid I in '*C.*' *klariankae*) and by its non-reduced M3. However, these traits are also present in the basal species of *Hispanomys*. Recently, Prieto et al. (2014) questioned its original inclusion in the genus *Cricetodon*. The position of '*C.*' *klariankae* in our MPT agrees with the doubts of Prieto et al. (2014) and, going further, we recommend its inclusion in *Hispanomys*.

PALEOBIOGEOGRAPHY AND EVOLUTIONARY HISTORY OF THE CRICETODONTINI

Figure 10.7 shows a calibrated cladogram with the geographical and chronostratigraphical information of the studied species. This figure provides a better picture about the relationships of the species through time and space. Appendix 10.2 present all the occurrences included in the analysis and the age of the localities.

The cladogram obtained has been dated using the *Paleo-Tree package* for the programming language and software environment *R*. The procedure needs for a numerical time interval, therefore we calculate the ages of the stratigraphical distributions of the studied species. Table 10.3 shows the information about the distributions as well as their "Calculated Oldest Date" (COD) and "Calculated Youngest Date" (CYD). The COD/CYD of the species has been obtained by means of different sources (See references in table 10.3). The criteria to establish the limits used in each case were as follows:

- 1. The COD and CYD of the species present in only one locality with numerical dating are the limits of the chron in which they are placed.
- 2. The COD and CYD of the species present in one or more sites without numerical dating:
 - 2.1. In the case of the European localities, the COD and CYD are the numerical limits of the MN in which they were placed. To avoid mistakes due to the diachronic character of the MN system (Gómez Cano et al., 2011; Van der Meulen et al., 2011, 2012), we used the numerical limits of the MN calculated for the basin in which the sites are

located. When the specific limits of the MN system for the basin were not available, we used the general limits proposed by Hilgen et al. (2012).

- 2.2. In the case of the Chinese localities, we used the age limits of the geologic formations containing the localities.
- 3. The COD and CYD of the species present in more than one locality, with all of them dated are the dates of the first and last occurrence of the species.

Figure 10.7 shows that the centre of origin of the studied Cricetodontini is Anatolia, where the tribe was present since the early Miocene. *Deperetomys* extends through Asia Minor and central Europe, *Cricetodon* is wide spread across Europe and Asia Minor, whereas *Byzantinia* is strongly linked with Asia Minor. *Hispanomys* + *Ruscinomys* are linked with southwestern Europe.

Three main intervals of migration of the group can be observed, which correspond to several biotic events. The first one starts around 18 Ma, it took *Cricetodon* and *Deperetomys* out of Anatolia giving rise to *C. aliverensis* and to the central European-like species such as *C. nievei*. This first interval is coincident with the ‘Creodont event’ described by Van der Made (1999). The second one starts around 15 Ma; it took another branch of *Cricetodon* out of Anatolia and dispersed it through Europe. The hypothetical common ancestor of the Iberian species, *C. soriae* and *C. jotae*, took part in this migration. This second interval is coincident with the ‘Hominid event’ described by Van der Made (1999). The last event is placed around 14 Ma and involved the migration of hypothetical basal species from Anatolia to southwestern Europe. *Hispanomys aguirrei*, *H. lavocati*, *H. nombrevillae* and *H. aragonensis* arose from species involved in this migration event. As noted in chapter 5, Van der Made (1999) described the ‘*Conohyus/Pliopithecus* event’ at the same age. This last migration event is coincident with a rise in the diversity of the tribe in Asia and Europe, having its maximum at the MN7/8. This diversification coincides also with the first occurrence of the genus *Hispanomys* (*H. decedens*) and *Deperetomys* (*D. hagni*) in Europe, and *Byzantinia* in Asia Minor.




FIGURE 10.3. Phylogenetic relationships of Cricetodontini within Cricetidae (Rodentia). A single strict consensus tree (773 steps CI: 0.194; RI: 0.578) has been obtained from four most parsimonious trees using TNT v. 1.1 (Goloboff et al., 2008). The pair of numbers above and below a branch represents absolute and relative Bremer values respectively (according to Goloboff et al., 2008). A detailed character/taxa matrix is provided in the Appendix 10.1. The two possibilities in the original trees for the polytomy in Node H are present in (a) and (b).

Node A	Node G	Node M	Node T
12 0→1	28 0→2	3 0→1	1 0→2
35 0→1	39 0→1	15 2→0	15 1→2
36 0→1		20 0→1	16 0→1
45 0→2		26 0→2	30 0→1
102 0→1	Node H	49 0→1	92 0→1
	34 0→1	77 0→1	
	91 0→1		
Node B		Node N	Node U
23 1→2		7 0→1	12 1→0
31 1→0		42 0→1	27 0→2
46 1→0		48 1→2	35 1→2
63 0→1	Node I	66 0→1	38 1→2
	13 0→1	99 1→2	73 1→0
	44 0→1		90 0→1
	46 0→1		
Node C	48 2→1	Node O	
23 2→0	18 0→1	1 1→0	
55 0→1	99 2→0	37 0→2	
91 1→0			Node V
			25 0→1
Node D	Node J	Node P	26 0→1
30 1→0	15 0→2	8 0→1	42 0→2
64 0→1	24 0→1	30 1→0	44 1→0
71 0→2	59 0→1	32 0→2	59 0→1
73 1→0	75 1→2	47 1→0	63 0→1
	84 0→1		74 1→0
	93 0→1		86 1→0
Node E		Node Q	
4 0→1		31 1→0	
17 0→2		36 2→1	
20 0→1	Node K		
22 0→2	1 0→2	Node R	
32 1→0	53 0→1	11 1→0	
38 0→1	56 0→1	37 2→1	
48 1→2	76 2→0	44 0→1	
67 1→0		60 0→2	
68 1→2		81 1→0	
75 1→0		102 0→2	
77 0→1			
79 0→1	Node L		
88 0→1	1 2→1	Node S	
98 0→1	4 0→1	15 2→1	
100 1→2	17 0→1	28 2→0	
	19 0→1	41 0→2	
	33 2→1	48 1→0	
Node F	48 2→1	49 0→2	
46 1→0	78 1→0	51 0→1	
95 1→2	97 0→1		

FIGURE 10.4. Character distribution for the discussed internal nodes.

<i>B. bayraktensis</i>	<i>B. pikermiensis</i>	<i>C. candirensis</i>	<i>C. jotae</i>	<i>C. pasalarensis</i>	<i>C. wolkeri</i>
1 0→1	11 0→1	3 0→1	12 0→1	21 0→1	29 1→0
26 2→0	45 0→2	15 2→1	23 0→1	32 1→2	32 1→0
29 1→0	47 0→2	18 0→1	28 1→0	71 0 or 2→1	83 0→1
99 0→2	49 1→2	20 0→1	33 2→1	95 2→1	
102 0→1	58 0→2	24 1→0	35 0→1	97 0→2	
	61 2→1	51 0→1	36 0→1		
	63 1→0	65 0→1	37 0→1	<i>C. sansaniensis</i>	
<i>B. dardanellensis</i>		84 1→0	38 0→1	1 0→2	
20 1→0	<i>B. sofçaensis</i>		49 0→1	28 2→0	
38 2→1	13 1→0	<i>C. cariensis</i>	76 2→0	63 0→1	' <i>C. fandli</i>
65 0→2	18 0→1	16 0→1	77 0→1	85 1→0	18 0→1
69 1→0	28 2→0	36 2→0	86 2→0	97 0→1	35 1→0
80 1→2	33 1→2	56 1→0	90 1→0		43 1→0
	35 1→0	69 0→1	97 0→2	<i>C. soriae</i>	64 0→1
<i>B. eskhisarensis</i>	49 1→2	81 1→0		15 2→0	74 0→2
1 0→1	54 1→0	89 0→1		23 0→2	92 1→0
7 1→0	68 0→1		<i>C. jumaensis</i>	28 2→1	
10 1→0	84 1→0		9 2→1	30 1→0	
13 1→0	89 0→1	<i>C. caucasicus</i>	22 0→1	59 1→0	
18 0→1		4 1→0	35 0→1	61 0→1	
21 1→2	<i>C. albanensis</i>	9 2→1	56 0→1	74 1→0	
22 1→2	26 0→2	17 1→0	75 1→0	76 2→0	
32 2→1	27 0→1	29 1→0	91 0→1	77 0→1	' <i>C. klariankae</i>
40 1→2	30 1→0	85 1→2		80 0→1	14 0→1
51 0→2	45 0→1				33 1→0
61 2→1	54 1→0			<i>C. tobieni</i>	35 1→0
69 0→1	60 0→1 or 2	<i>C. engesseri</i>	<i>C. kasapligili</i>	30 1→0	71 2→0
73 0→1	65 0→1	1 1→2	35 0→1	31 1→0	77 0→1
76 0→2	102 0→1	24 1→0	72 0→1	62 0→1	80 1→2
78 0→1		28 0→2	74 2→0	103 0→1	81 0→1
80 1→2	<i>C. aliverensis</i>	29 1→0			89 0→1
102 0→1	13 0→1	33 2→1	<i>C. meini</i>	<i>C. versteegi</i>	90 0→1
<i>B. hellenicus</i>	32 1→0	65 0→1	59 1→0	29 1→2	
No autoapomorphies	86 2→0	72 0→1	65 0→1	32 1→0	
<i>B. nikosi</i>	<i>C. aureus</i>	81 0→1	68 0→1	65 0→2	
4 1→2	8 1→0		78 1→0	69 0→1	
11 0→1	26 0→1	<i>C. hungaricus</i>	82 1→0	84 0→1	
16 0→2	29 1→0	14 1→0	91 1→0		
44 0→1	68 1→0	15 2→0	93 1→0	<i>C. nievei</i>	<i>C. wanhei</i>
64 1→0	72 0→1	23 0→2		1 1→0	25 0→2
	74 1 or 2→0	37 0→1		13 0→1	26 0→2
<i>B. ozansoyi</i>	<i>C. bolligeri</i>	68 0→1		23 2→0	27 0→2
9 2→1	9 2→1	70 1→0		28 0→2	35 0→1
10 1→0	29 1→0			33 2→1	41 0→2
27 0→2	32 1→2			48 1→0	45 0→2
32 2→1	42 0→2			51 0→1	49 0→1 or 2
51 0→3	65 0→1			53 0→1	55 1→0
69 0→1	68 1→0			56 0→1	78 1→0
77 1→0	72 0→1			84 0→1	99 2→1
80 1→2	95 1→0			97 0→1	102 0→1

FIGURE 10.5. Character distribution for each studied species of *Byzantinia* and *Cricetodon*.

<i>D. anatolicus</i>	<i>H. baixasi</i>	<i>H. decedens</i>	<i>H. lavocati</i>
1 1→0	36 2→0	8 1→0	1 0→2
	37 2→0	19 1→0	3 0→1
	62 1→0	21 1→0	48 1→2
<i>D. hangi</i>	65 2→1	44 1→0	51 0→1
1 1→2	73 0→1	47 0→1	61 0→1
3 0→1	102 2→1	53 1→0	81 0→1
23 2→0		56 1→0	86 0→2
24 1→0	<i>H. bijugatus</i>	61 0→1	99 0→1
38 1→2	1 0→1	74 0→2	102 2→1
46 0→1	2 1→0	76 0→2	103 1→0
49 1→0	11 0→1	78 0→1	
61 0→2	13 1→0	<i>H. dispectus</i>	<i>H. peralensis</i>
78 1→0	27 0→1	8 1→0	9 2→1
95 1→2	35 1→0	27 0→1	15 1→0
	42 0→2	29 1→0	20 0→1
<i>D. intermedius</i>	51 0→2	31 1→0	57 1→0
18 0→1	59 1→0	35 1→2	81 0→1
19 0→1	71 2→0	47 0→1	
26 0→2	82 1→0	60 2→0	<i>H. thaleri</i>
63 1→0	101 1→0	81 0→1	25 1→0
64 1→0	102 2→0	85 1→0	29 1→2
	103 1→0		31 1→0
<i>H. adroveri</i>	<i>H. castelnovi</i>	<i>H. mediterraneus</i>	37 2→1
29 1→0	9 2→1	27 2→0	47 0→1
65 2→1	18 0→1	33 0→1	99 2→1
	20 0→1	68 0→1	102 2→1
	30 0→1	70 1→0	103 1→0
	39 1→0	73 0→1	
<i>H. aguirrei</i>	85 1→2	90 1→0	<i>R. europaeus</i>
9 2→1	92 1→0		16 1→2
64 0→1	99 0→2	<i>H. moralesi</i>	25 1→2
72 0→1	102 2→1	29 1→2	26 1→2
74 0→1		30 0→1	30 0→1
91 1→0	<i>H. daamsi</i>	53 1→0	33 0→1
92 1→0	20 0→1	80 1→2	48 0→2
95 0→1	26 0→1	85 1→2	97 0→1
101 1→0	28 0→2	87 0→1	
	38 1→2		<i>R. schaubi</i>
<i>H. aragonensis</i>	51 1→0		29 1→2
8 1→0	53 1→0	<i>H. nombrevillae</i>	76 0→2
94 0→1	73 1→0	63 0→1	103 1→0
95 0→2	75 2→1	64 0→1	
96 2→1	78 0→1		
102 2→1	95 0→2		
	103 1→0		

FIGURE 10.6. Character distribution for each studied species of *Deperetomys* and *Hispanomys*.

FINAL REMARKS & CONCLUSIONS

Three main clades can be observed in our cladistic analysis which correspond to *Deperetomys*, *Byzantinia*, and *Hispanomys* + *Ruscinomys*. The cladogram reveals that *Deperetomys* and *Byzantinia* are monophyletic, *Cricetodon* is paraphyletic and the species of *Hispanomys* together with *Ruscinomys* form a clade. There are two polytomies, one involves several species of *Cricetodon* and the other one various species of *Byzantinia*.

The morphological patterns change towards a simpler occlusal surface: from complicated forms displaying brachyodont and bunodont teeth with high density of transversal crests, to forms with prevalence of longitudinal ridges in hypsodont and selenodont-like molars.


Regarding the species from the Calatayud-Daroca Basin, they are grouped as follows:

Cricetodon soriae, *C. sansaniensis* and *C. jotae* are grouped together in a clade with *C. meini*. *Cricetodon nievei* is clustered with the species from central Europe, clearly separated from the other Cricetodontini from the Calatayud-Daroca Basin. The species of *Hispanomys* are split in two clades: *H. aguirrei* and *H. lavocati* are basal, and *H. nombrevillae* and *H. aragonensis* are derived.

The main taxonomical conclusions are:

Deperetomys is included in the tribe Cricetodontini, although its relationship with the species of *Cricetodon* from the Node A is questionable. Besides, we recommend a new taxonomical study of the species originally ascribed to *Turkomys* to evaluate the synonymy *Turkomys*–*Cricetodon*. This might resolve the paraphyletic status of *Cricetodon*. Moreover, the revision of '*C.*' *fandli* is needed as it is related with *Hispanomys* in our MPT. We suggest the ascription of '*C.*' *klariankae* to *Hispanomys*.

After this cladistic study, many questions about the phylogeny and taxonomy of the Cricetodontini are resolved. However, some taxa need supplementary morphological data (especially those of *Ruscinomys*) to establish clearer differences between these genera. Moreover, the inclusion of the remaining species of *Deperetomys*, *Byzantinia* and *Ruscinomys*, as well as new genera like *Meteomys*, *Mirabella* and *Eumyarion* is strongly recommended to confirm some of the interpretations mentioned above and to better understand the evolution of Cricetodontini.



Species	COD	CYD	References used to calculate the COD/CYD	Criteria
<i>R. europaeus</i>	5.30	3.20	García-Alix, 2006	2.1
<i>R. schaubi</i>	7.75	6.00	Van Dam et al., 2006/Morales et al., 2013	3
<i>H. adroveri</i>	8.05	6.80	Morales et al., 2013/García-Alix, 2006	3
<i>H. baixasi</i>	8.12	6.66	Gómez Cano et al., 2011	2.1
<i>B. nikosi</i>	9.00	8.70	Koufos, 2013	1
<i>H. mediterraneus</i>	9.61	8.12	Gómez Cano et al., 2011	2.1
<i>H. thaleri</i>	9.70	9.96	Agustí et al., 1997	1
<i>H. peralensis</i>	9.85	8.85	Van Dam et al., 2006	3
<i>B. hellenicus</i>	9.90	6.80	De Bruijn et al., 2013/Garcés et al., 1998	2.1
<i>H. moralesi</i>	9.94	8.80	Van Dam et al., 2006	2.1
<i>H. aragonensis</i>	11.72	9.98	Van Dam et al., in press	3.1
<i>H. daamsi</i>	11.10	9.88	Agustí et al., 2005/Gómez Cano, 2011	2.1
<i>B. pikermiensis</i>	11.20	7.00	De Bruijn et al., 2013/Hilgen et al., 2012	2.1
<i>B. dardanellensis</i>	11.20	9.90	Hilgen et al., 2012	2.1
<i>H. nombrevillae</i>	11.20	10.77	Van Dam et al., in press	3
<i>H. lavocati</i>	11.87	11.11	Van Dam et al., in press/Carmona et al., 2011	3
<i>C. wolkeri</i>	12.00	11.00	Wang et al., 2013	2.2
<i>'C'. fandli</i>	12.16	12.07	Prieto et al., 2010	1
<i>'C'. klariankae</i>	12.50	12.00	Hír and Kókay, 2010	3
<i>B. bayraktepensis</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>B. eskhisarensis</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>B. ozansoyi</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>B. sofcanensis</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>C. cariensis</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>D. hagni</i>	12.50	11.00	Hilgen et al., 2012	2.1
<i>H. aguirrei</i>	12.39	11.33	Van Dam et al., in press	3
<i>H. bijugatus</i>	12.67	9.70	Agustí et al., 2001/Vagengein and Tesakov, 2013	3/2.1
<i>H. decedens</i>	12.70	11.10	Hilgen et al., 2012	3
<i>H. dispectus</i>	12.75	11.00	Agustí et al., 2001/Casanovas Vilar et al., 2006	2.1
<i>C. albanensis</i>	12.75	12.50	Agustí et al., 2001	2.1/3
<i>C. nievei</i>	12.65	12.60	Van Dam et al., 2006/Van Dam et al., in press	3
<i>C. jatae</i>	13.56	12.65	Van Dam et al., 2006/Van Dam et al., in press	3
<i>H. castelnovi</i>	13.80	12.75	Gómez Cano et al., 2011	2.1
<i>C. bolligeri</i>	13.80	13.00	Hilgen et al., 2012/Kálin and Kempf, 2009	2.1
<i>C. engesseri</i>	13.90	13.80	Kálin and Kempf, 2009	2.1
<i>C. jumaensis</i>	14.00	13.40	Kálin and Kempf, 2009	3
<i>C. soriae</i>	14.06	13.80	Van Dam et al., 2006	3
<i>C. sansaniensis</i>	14.10	13.68	Kálin and Kempf, 2009/Van Dam et al., 2006	3
<i>C. caucasicus</i>	14.20	12.85	Hilgen et al., 2012	2.1
<i>C. hungaricus</i>	14.20	12.85	Hilgen et al., 2012	2.1
<i>C. paslarensis</i>	14.20	12.85	Hilgen et al., 2012	2.1
<i>C. aureus</i>	15.05	14.00	Kálin and Kempf, 2009	3
<i>D. franconicus</i>	16.12	14.84	Van Dam et al., 2006	3
<i>C. candirensis</i>	16.40	14.95	Reichenbacher et al., 2013/Kálin and Kempf, 2009	2.1
<i>C. meini</i>	16.45	14.90	Reichenbacher et al., 2013	2.1
<i>C. aliveriensis</i>	19.72	18.75	Koufos, 2013	1
<i>C. wanhei</i>	19.00	16.00	Wang et al., 2013	2.2
<i>C. tobieni</i>	19.50	16.40	Hilgen et al., 2012	2.1
<i>C. kasapligili</i>	19.50	17.00	Hilgen et al., 2012	2.1
<i>D. intermedius</i>	21.70	19.50	Hilgen et al., 2012	2.1
<i>C. versteegi</i>	23.00	21.70	Hilgen et al., 2012	2.1
<i>D. anatolicus</i>	23.00	21.70	Hilgen et al., 2012	2.1

TABLE 10.3. Species and ages for their Calculated Oldest Date (COD) and Calculated Youngest Date (CYD) applied to date the cladogram. The fourth column provides the references used to calculate the COD/CYD of each taxon. The criteria used in each taxon is present in the fifth column (methodology explained in “Paleobiogeography and Evolutionary History of Cricetodontini” section).

LITERATURE CITED

- Aguilar, J. P. 1982. Contributions à l'étude des Micromammifères du gisement Miocène supérieur de Montredon (Hérault) 2. Les rongeurs. *Palaeovertebrata* 12:75–140.
- Aguilar, J. P., M. Calvet, and J. Michaux. 1994. Les rongeurs de Castelnou 6 (Pyrénées-orientales, France) et les corrélations entre faunes ibériques et françaises au Miocène moyen. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 192:109–131.
- Aguilar, J. P., V. Lazzari, J. Michaux, M. Sabatier, and M. Calvet. 2007. Lo Fournas 16-M (Miocène supérieur) et Lo Fournas 16-P (Pliocène moyen), nouvelles localités karstiques à Baixas, Sud de la France. Partie I- Description et implications géodynamiques. *Géologie de la France* 1:55–62.
- Agustí, J. 1980. La asociación de *Hispanomys* y *Cricetodon* (Rodentia, Mammalia) en el Mioceno superior del Vallès-Penedès (Cataluña, España). *Acta Geologica Hispanica* 15:51–60.
- Agustí, J. 1982. Tendencias evolutivas de la línea *Cricetodon-Ruscinomys* (Rodentia, Mammalia) en la Península Ibérica. *Acta Geologica Hispanica* 17:103–111.
- Agustí, J. 1986. Nouvelles espèces de Cricétidés vicariantes dans le Turolien moyen de Fortuna (Province de Murcia, Espagne). *Geobios* 19:5–11.
- Agustí, J., I. Casanovas-Vilar, and M. Furió. 2005. Rodents, insectivores and chiropterans (Mammalia) from the late Aragonian of Can Missert (middle Miocene, Vallès-Penedés Basin, Spain). *Geobios* 38:575–583.
- Agustí, J., S. Moyà-Solà, and J. Pons Moyà. 1984. Mammal distribution dynamics in the eastern margin of the Iberian Peninsula during the Miocene. *Paléobiologie continentale* 14:33–46.
- Agustí, J., Ll. Cabrera, M. Garcés, and J. M. Parés. 1997. The Vallesian mammal succession in the Vallès-Penedès basin (northeast Spain): Paleomagnetic calibration and correlation with global events. *Palaeogeography, Palaeoclimatology, Palaeoecology* 133:149–180.
- Agustí, J., L. Cabrera, M. Garcés, W. Krijgsman, O. Oms, and J. M. Parés. 2001. A calibrated mammal scale for the Neogene of Western Europe. State of the art. *Earth-Science Reviews* 52: 247–260.

- Álvarez-Sierra, M. A., J. P. Calvo, J. Morales, A. Alonso-Zarza, B. Azanza, I. García Paredes, M. Hernández Fernández, A. J. Van der Meulen, P. Peláez-Campomanes, V. Quirarte, M. J. Salesa, I. M. Sánchez, and D. Soria. 2003. El tránsito Aragoniense-Vallesiense en el área de Daroca-Nombrevilla (Zaragoza, España). *Coloquios de Paleontología Volumen Extraordinario* 1:25–33.
- Argyropulo, A. I. 1938. On the fauna of Tertiary Cricetidae of the USSR. *Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS* 20:223–226.
- Bi, S. 2005. Evolution, systematics and functional anatomy of Cricetodontini (Cricetidae, Rodentia, Mammalia) from the Northern Juggar Basin, northwestern China. Ph.D. dissertation, Howard University, Washington, 156 pp.
- Bremer, K. 1988. The limits of amino acid sequence data in angiosperm phylogenetic reconstruction. *Evolution* 42:795–803.
- Bremer, K. 1994. Branch support and tree stability. *Cladistics* 10:295–304.
- Carmona, R., D. M. Alba, I. Casanovas-Vilar, M. Furió, M. Garcés, J. V. Bertó Mengual, J. Galindo, and À. H. Luján. 2011. Intervención paleontológica en la serie del Mioceno Superior del Ecoparc de Can Mata (cuenca del Vallès-Penedès, NE de la Península Ibérica) pp. 65–74 in A. Pérez-García, F. Gascó, J.M. Gasulla and F. Escaso (eds.), *Viajando a mundos pretéritos*. Morella: Ayuntamiento de Morella.
- Casanovas-Vilar, I., and J. Agustí. 2007. Ecogeographical stability and climate forcing in the Late Miocene (Vallesian) rodent record of Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 248:169–189.
- Casanovas-Vilar, I., M. Furió, and J. Agustí. 2006. Rodents, Insectivores and Paleoenvironment associated to the first-appearing Hipparionine horses in the Vallès-Penedès Basin (North-eastern Spain). *Beiträge zur Paläontologie* 30:89–107.
- Crusafont, M., and J. Villalta. 1956. Un nouveau *Ruscinomys* du Pontien espagnol et sa position systématique. *Comptes Rendus Sommaires de la Société Géologique de France* 7.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.
- De Bruijn, H. 1976. Vallesian and Turolian rodents from Biotia, Attica and Rhodes (Greece). *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen* 79:361–384.

- De Bruijn, H., and G. Saraç. 1992. Early Miocene rodent faunas from the Eastern Mediterranean area. Part II. *Mirabella* (Paracricetodontinea, Muroidea). Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 95:25–40.
- De Bruijn, H., and E. Ünay. 1996. On the evolutionary history of the Cricetodontini from Europe and Asia Minor and its bearing on the reconstruction of migrations and the continental biotope during the Neogene; pp. 227–234 in R. L. Bernor, V. Fahlbusch, H.W. Mittmann (eds.), *The Evolution of Western Eurasian Neogene Mammal Faunas*. Columbia University Press, New York.
- De Bruijn, H., E. Ünay, and K. Hordijk. 2013. A review of the Neogene Succession of the Muridae and Dipodidae from Anatolia, with special reference to taxa Known from Asia and/or Europe; pp. 566–582 in X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Saraç, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a Discussion of the Evolutionary History of the Cricetodontini. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 96:151–216.
- De Bruijn, H., E. Ünay, G. Saraç, and G. K. Klein Hofmeijer. 1987. An unusual new Eucricetodontine from the lower Miocene of the eastern Mediterranean. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 90:129–132.
- De Bruijn, H., L. W. Van den Hoek Ostende, M. R. Kristkoiz-Boon, K. Theocharopoulos, and E. Ünay. 2003. Rodents, lagomorphs and insectivores, from the middle Miocene hominoid locality of Çandır (Turkey). *Courier Forschungsinstitut Senckenberg* 240:51–87.
- De Bruijn, H., R. Daams, V. Fahlbusch, G. Daxner-Höck, L. Ginsburg, P. Mein, and J. Morales. 1992. Report of the RCMNS working group on fossil mammals, Reisenburg 1990. *Newsletters of Stratigraphy* 26:65–118.
- Depéret, C. 1890. Les animaux Pliocènes du Roussillon. *Mémoires de la Société Géologique de France* 1:1–63.
- Fahlbusch, V. 1964. Die Cricetiden (Mammalia) der Oberen Süßwasser-Molasse Bayerns. *Bayerische Akademie Wissenschaften. Mathematisch-naturwissenschaftliche Klasse. München* 118:1–135.
- Fahlbusch, V. 1966. Cricetidae (Rod. Mam.) aus der Mittelmiozänen Spalten Füllung Erkertshofen bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie. München* 6:109–131.

- Freudenthal, M. 1963. Entwicklungsstufen der miozänen Cricetodontinae (Mam. Rod.), Mittelspaniens und ihre stratigraphische Bedeutung. *Beaufortia*, 10:51–157.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 69:298–317.
- Freudenthal, M. 1970. A new *Ruscinomys* (Mammalia, Rodentia) from the late Tertiary (Pliocene). *American Museum Novitates* 2402:1–10.
- García-Alix, A. 2006. Bioestratigrafía de los depósitos continentales de la transición Mio-Plioceno de la cuenca de Granada. Ph.D. dissertation, Universidad de Granada, 429 pp.
- Goloboff, P. A., J. S. Farris, and K. C. Nixon. 2008. TNT, a free program for phylogenetic analysis. *Cladistics* 24:774–786.
- Gomes Rodrigues, H., L. Marivaux, and M. Vianey-Liaud. 2009. Phylogeny and systematic revision of Eocene Cricetidae (Rodentia, Mammalia) from Central and East Asia: on the origin of cricetid rodents. *Journal of Zoological Systematics and Evolutionary Research* 48:259–268.
- Gómez Cano, A. R., M. Hernández Fernández, and M. A. Álvarez-Sierra. 2011. Biogeographic provincialism in rodent faunas from the Iberocceanian Region (southwestern Europe) generates severe diachrony within the Mammalian Neogene (MN) biochronologic scale during the Late Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 307:193–204.
- Hartenberger, J. L. 1965. Les Cricetidae (Rod.) de Can Llobateres (Néogène d'Espagne). *Bulletin De La Societe Geologique De France* 7:487–498.
- Hernández Fernández, M., J. A. Cárdenas, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Hilgen, F. J., L. J. Lourens, and J. A. Van Dam. 2012. The Neogene Period; pp. 923–978 in F. M. Gradstein, J. G. Ogg, J. G. Schmitz (eds.), *Geologic Time Scale*. Cambridge University Press, Cambridge.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőtárkány-Felnémet (Northern Hungary). *Fragmenta Paleontologica Hungarica* 24–25:16–24.

- Hír, J., and J. Kókay. 2010. A systematic study of the middle-late Miocene rodents and lagomorphs (Mammalia) of Felsőtárkány 3/8 and 3/10 (Northern Hungary) with stratigraphical relations. *Geodiversitas* 32:307–329.
- Jaeger, J. J., J. Michaux, and B. David. 1973. Biochronologie du Miocène moyen et supérieur continental du Maghreb. *Les Comptes Rendus de l'Académie des sciences, séries D* 277:2477–2480.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254:177–235.
- Kordos, L. 1986. Upper miocene hamsters (Cricetidae, Mammalia) of Hasznos and Szentendre. *Magyar Allami Földtani Intézet Jelentése Az* 1984:523–553.
- Kostopoulos, D. S. 2009. The Pikermian event: Temporal and spatial resolution of the Turolian large mammal fauna in SE Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology* 274:82–95.
- Koufos, G. D. 2013. Neogene Mammal Biostratigraphy and Chronology of Greece; pp 595–628 in X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- Klein Hofmeijer, G., and H. De Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece). Part 8: The Cricetidae. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 91:185–204.
- Lartet, E. 1851. Notice sur la colline de Sansan, suivie d'une récapitulation des diverses espèces d'animaux vertébrés fossiles, trouvés soit à Sansan, soit dans d'autres gisements du terrain tertiaire miocène dans le Bassin Sous-Pyrénéen. J. A. Portes, Auch, 41 pp.
- López-Antoñanzas, R., and P. Mein. 2009. First detailed description of *Hispanomys bijugatus* Mein & Freudenthal, 1971 (Rodentia, Cricetodontinae) from the Upper Aragonian of La Grive-Saint Alban (France): Biostratigraphical implications. *Geobios* 42:783–796.
- López-Antoñanzas, R., and P. Mein. 2011. First detailed description of *Hispanomys decedens* (Rodentia) from the Middle Miocene of La Grive-Saint Alban (France). *Swiss Journal of Geosciences* 104:345–353.
- López-Antoñanzas, R., P. Peláez-Campomanes, M. A. Álvarez-Sierra, and I. García-Paredes. 2010. New species of *Hispanomys* (Rodentia, Cricetodontinae) from the Upper Miocene of Batallones (Madrid, Spain). *Zoological Journal of the Linnean Society* 160:725–747.

- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89: 573–592.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. In press. Cricetodontini from the Calatayud-Daroca Basin (Spain): a taxonomical description and up-to-date of their stratigraphical distributions. *Comptes Rendus Palevol*. DOI: <http://dx.doi.org/10.1016/j.crpv.2014.07.002>.
- López-Martínez, N., C. Sesé Benito, and J. L. Sanz García. 1977. La microfauna (Rodentia, Insectivora, Lagomorpha y Reptilia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno/Cuaternario* 8:47–73.
- Maridet, O., and X.J. Ni. 2013. A new Cricetid rodent from the early Oligocene of Yunnan, China, and its evolutionary implications for early Eurasian Cricetids. *Journal of Vertebrate Paleontology* 33:185–194.
- Maridet, O., and S. Sen. 2012. Les Cricetidae du gisement de Sansan; pp. 29–64 in S. Peigné and S. Sen (eds.), *Les mammifères du gisement de Sansan. Mémoires du Muséum national d'Histoire naturelle*, Paris.
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Morales, J., P. Peláez-Campomanes, J. Abella, P. Montoya, L. Gibert, G. Scott, J. L. Cantalapiedra, and O. Sanisidro. 2013. The Ventian mammal age (Latest Miocene): present state. *Spanish Journal of Palaeontology* 28:149–160.
- Morrone, J. J. 2000. *El lenguaje de la cladística*. Universidad Nacional Autónoma de México, México D. F., 109 pp.
- Peláez-Campomanes, P., and R. Daams. 2002. Middle Miocene rodents from Pasalar, Anatolia, Turkey. *Acta Paleontologica Polonica* 47:125–132.
- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria):

- a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.
- Prieto, J., C. Angelone, I. Casanovas-Vilar, M. Gross, J. Hír, L. W. Van den Hoek Ostende, L. C. Maul, and D. Vasilyan. 2014. The small mammals from Gratkorn: an overview. *Palaeobiology and Palaeoenvironments* 94:135–162.
- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata PalAsiatica* 48:27–47.
- Reichenbacher, B., W. Krijgsman, Y. Lataster, M. Pippèrr, C. G. C. Van Baak, L. Chang D. Kálin, J. Jost, G. Doppler, D. Jung, J. Prieto, H. Abdul Aziz, M. Böhme J. Garnish, U. Kirscher, and V. Bachtadse. 2013. A new magnetostratigraphic framework for the Lower Miocene (Burdigalian/Ottnangian, Karpatian) in the North Alpine Foreland Basin. *Swiss Journal of Geosciences* 106:309–334.
- Rummel, M. 1995. *Cricetodon bolligeri* n. sp. ein neuer Cricetide aus dem Obermiozän von Petersbuch bei Eichstätt. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 35:109–123.
- Rummel, M. 1999. Tribe Cricetodontini; pp. 359–364 in G. Rößner, and K. Heißig (eds.), *The Miocene land mammals of Europe*. Verlag Dr. Friedrich Pfeil, München.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde B* 311:1–6.
- Rummel, M., and D. Kálin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Mittelmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Schaub, S. 1925. Die hamsterartigen Nagetiere des Tertiärs und ihre lebenden Verwandten. *Abhandlungen der Schweizerischen paläontologischen Gesellschaft* 45:1–114.
- Sen, S., and M. A. Erbajeva. 2011. A new species of *Gobicricetodon* Qiu, 1996 (Mammalia, Rodentia, Cricetidae) from the Middle Miocene of Aya Cave, Lake Baikal. *Vertebrata PalAsiatica* 49:257–274.
- Sen, S., and E. Ünay. 1979. Sur quelques Cricetodontini (Rod.) du Miocène moyen d'Anatolie. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* 82:293–301.
- Thaler, L. 1966. Les rongeurs fossiles du Bas-Languedoc dans leurs rapports avec l'histoire des faunes et la stratigraphie du Tertiaire d'Europe. Ph.D. dissertation, Faculté des Sciences de Paris, Paris, 295 pp.
- Tobien, H. 1978. New species of Cricetodontini (Rod. Mam.) from the Miocene of Turkey.

- Mainzer Geowissenschaftliche Mitteilungen Geologisches Landesamt Rheinland-Pfalz 6:209–219.
- Ünay, E. 1980. The Cricetodontini (Rod. Mam.) from the Bayraktepe section (Çanakkale, Turkey). *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen, Series B* 83:399–418.
- Ünay, E. 1989. Rodents from the Middle Oligocene of Turkish thrace. *Utrecht micropaleontological bulletins. Special publication. Utrecht* 5:1–119.
- Ünay, E., and H. de Bruijn. 1984. On some Neogene Rodent assemblages from both sides of the Dardanelles, Turkey. *Newsletters Stratigraphy* 13:119–132.
- Van Dam, J., W. Krijgsman, H. A. Abels, M. A. Álvarez-Sierra, I. García-Paredes, P. López-Guerrero, P. Peláez-Campomanes y D. Ventura. In press. Updated chronology for middle to late Miocene mammal sites of the Daroca area (Calatayud-Montalbán Basin), Spain. DOI: <http://dx.doi.org/10.1016/j.geobios.2014.07.002>.
- Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L.W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. Van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodent faunas of the Mio-Pliocene continental sediments of the Teruel-Alfambra region, Spain. *Utrecht micropaleontological bulletins. Special publication. Utrecht* 2:1–218.
- Van der Made, J. 1999. Intercontinental relationship Europe-Africa and the Indian Subcontinent; pp. 359–364 in G. Rößner and K. Heißig (eds.), *The Miocene Land Mammals of Europe*. Verlag Dr. Friedrich Pfeil, München.
- Vangengeim, E. A., and S. Tesakov. 2008. Maeotina Mammalian Localities of Eastern Paratethys: Magnetostratigraphy and Position in European Continental Scales. *Stratigraphy and Geological Correlation* 16:437–450.
- Wang, X., Q. Li, Z.-d. Qiu, G.-p. Xie, B.-y. Wang, Z.-x. Qiu, Z.-j. Tseng, G. T. Takeuchi, and T. Deng. 2013. Neogene Mammalian Biostratigraphy and Geochronology of the Tibetan Plateau; pp. 243–292 in X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.

Appendix 10.1

Dental character matrix (non-preserved "?" and non applicable "-" characters).

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52
D. francanicus	0	?	0	0	0	?	1	1	0	0	?	?	1	0	0	1	0	0	0	0	0	0	2	0	0	0	0	0	0	1	1	1	2	0	0	0	0	0	0	0	0	0	1	0	0	1	2	0	2	0	0	
H. castelnovi	0	?	0	0	?	0	1	1	0	1	1	?	?	1	0	1	1	1	1	1	1	?	?	1	0	0	0	0	0	1	1	1	2	1	1	1	1	2	1	1	0	1	?	0	1	0	2	2	1	1		
H. H. decedens	0	1	0	1	0	081	0	2	0	0	0	?	1	2	0	1	0	0	0	0	0	1	1	1	1	0	0	0	2	0	0	0	1	1	1	1	1	2	1	1	0	1	?	0	1	0	2	2	1	1		
H. H. bijugatus	1	0	0	1	0	1	0	1	2	0	1	0	0	1	2	0	1	0	1	0	1	0	2	1	0	0	0	1	2	1	0	2	1	1	0	1	1	2	1	1	0	2	1	0	0	1	1	2	2	1		
H. H. lavacati	2	1	1	0	1	0	1	2	0	1	0	1	0	2	0	1	0	1	0	1	0	1	1	1	1	0	0	2	0	0	081	1	1	1	1	1	1	1	1	1	0	0	1	1	0	0	2	0	2	1	0	
H. H. respecti	0	?	0	1	0	1	0	2	0	?	0	?	1	0	0	1	0	1	?	0	0	1	?	0	0	0	0	0	0	0	2	1	2	2	1	1	2	2	1	1	0	1	1	0	1	0	2	2	1	1		
H. H. aguirrei	0	1	0	081	0	081	0	1	1	0	0	1	1	0	2	0	1	0	1	0	1	1	1	1	0	0	0	2	0	0	1	1	1	1	1	1	1	1	1	1	1	0	0	1	1	0	0	1	0	2	0	-
H. H. daamsi	2	?	1	1	0	1	0	1	2	0	0	?	1	0	0	1	1	?	0	1	1	?	0	0	0	0	0	0	0	1	1	2	1	?	1	2	2	1	1	2	0	1	1	0	0	1	0	2	0	-		
H. H. namorevilleae	0	1	1	1	0	1	0	1	2	0	0	?	1	0	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2	1	1	1	2	2	1	1	2	0	1	0	1	0	0	2	2	1		
H. H. thaleri	1	1	1	2	0	1	0	1	1	0	?	?	1	1	1	0	1	1	1	1	1	1	0	0	1	1	0	0	1	2	2	1	0	2	0	1	2	2	1	?	1	?	1	?	1	1	?	?	2	1	1	
H. H. bogosensis	0	?	1	2	0	1	0	0	2	0	0	1	1	0	1	0	1	0	1	0	1	0	0	0	0	0	0	0	2	0	1	1	2	0	1	1	2	2	1	1	2	0	0	1	0	0	2	2	1	1		
H. H. axiosai	1	?	1	?	0	1	0	1	2	0	?	0	1	1	0	1	0	1	0	1	0	1	1	0	0	1	0	0	1	2	0	1	1	2	0	1	1	2	0	2	1	1	2	0	0	1	0	2	2	0	-	
H. H. mediterraneus	1	?	?	?	0	1	0	1	1	0	?	?	1	1	?	0	1	1	1	1	1	?	?	1	1	?	1	1	0	2	1	1	2	1	1	2	2	?	?	1	2	2	1	?	?	0	1	?	2	1	1	
H. H. moralesi	0	?	1	2	0	1	0	1	2	0	0	?	1	0	1	0	1	0	1	0	1	1	2	2	1	0	1	1	2	2	1	2	0	1	2	2	2	2	1	1	2	0	0	1	0	0	2	2	0	-		
H. H. peralensis	0	1	1	2	0	1	0	1	1	0	0	0	?	?	0	0	1	0	1	1	1	1	0	0	1	0	0	1	1	2	2	0	1	2	0	182	2	2	2	2	1	1	2	0	0	1	0	2	2	1		
H. H. adroveri	2	?	1	2	0	1	0	1	2	0	0	?	?	1	0	1	1	1	1	1	1	1	0	0	1	1	0	1	1	2	2	0	1	2	0	2	1	2	2	1	2	2	0	?	1	0	0	2	0	2	1	
R. R. schaubi	2	1	1	2	0	1	0	1	2	0	?	?	1	1	0	1	1	1	1	1	1	1	0	0	1	1	0	1	2	2	1	1	2	0	2	2	2	2	2	1	1	2	1	0	?	1	0	2	2	1		
R. R. aporeus	0	0	0	0	0	0	1	182	0	1	0	1	0	1	2	0	0	-	0	0	0	0	1	0	0	0	0	1	2	1	1	1	2	1	0	0	0	0	0	0	0	1	?	0	1	1	0	2	0	-		
C. C. meini	0	0	0	0	0	0	1	2	0	1	1	1	0	2	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	1	0	0	0	1	1	1	1	0	2	0	-	
C. C. jatae	0	0	0	0	0	1	2	0	1	1	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	1	1	1	1	1	1	1	1	0	0	0	1	1	1	1	1	2	0	-	
C. C. samsanicus	2	0	0	0	0	0	1	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	2	0	0	0	0	0	0	0	1	0	1	1	0	2	0	0	-		
C. C. versteegi	0	?	0	0	0	0	1	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	2	?	?	0	0	0	0	0	0	?	?	2	0	0	0	0	0	0	0	0	1	0	1	1	0	2	0	-		
C. C. kasaplilgili	0	?	0	0	0	0	1	081	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	1	2	0	1	0	0	0	0	0	0	0	1	0	0	2	0	2	1		
C. C. alverlensis	0	?	?	0	0	0	1	2	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	2	0	0	0	0	0	0	0	0	1	0	1	2	0	0	-	
C. C. tabieni	0	?	0	0	0	0	1	1	0	1	0	1	0	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	081	081	1	1	0	2	0	0	0	0	0	0	0	0	0	1	0	2	0	-	
C. C. wanhel	0	?	?	0	0	0	0	1	081	0	?	?	0	0	1	0	0	0	0	0	0	0	0	0	2	2	2	2	2	1	1	1	1	1	1	1	1	1	1	1	1	1	0	0	1	1	1	2	0	0	-	
C. C. pasalensis	0	?	0	0	0	0	1	0	?	?	?	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	?	?	2	0	0	0	0	0	0	0	0	0	0	1	0	1	2	0	0	-		
C. C. wolkeri	0	?	0	0	?	0	0	?	1	0	?	?	0	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	
"C". C. fandli	0	?	?	1	0	1	0	1	2	0	1	0	?	1	2	?	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	1	2	1	0	2	2	1	0	0	0	0	0	0	0	0	0	2	0	-		
C. C. nievei	0	?	0	0	0	0	1	2	0	0	1	1	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	-	
C. C. bolligeri	1	1	0	0	0	0	1	1	0	0	1	0	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	-	
C. C. candiensis	1	?	1	1	0	1	0	2	0	1	0	0	?	?	0	1	1	1	1	1	1	1	?	?	?	?	0	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
C. C. jumaensis	1	?	0	0	0	0	1	1	0	?	?	0	0	0	0	?	0	0	0	0	0	1	-	1	0	0	0	0	0	1	1	1	081	2	1	1	0	0	0	1	0	?	?	?	?	?	?	?	?	?		
C. C. cariensis	1	?	0	1	0	081	0	2	0	?	?	0	1	2	1	0	?	0	1	0	1	0	0	082	2	1	1	0	0	1	1	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	-
C. C. aureus	1	?	0	0	0	0	2	0	1	0	0	?	?	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	-
"C". C. klariankae	2	?	1	1	0	1	0	1	2	0	?	?	1	1	0	1	1	0	1	0	1	1	0	0	0	0	0	0	0	0	1	1	2	0	1	0	2	1	1	1	2	1	1	0	1	0	0	2	1	1		
C. C. albanensis	2	?	0	0	0	0	1	2	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	182	182	1	0	0	0	0	0	0	0	0	0	0	0	0	2	0	-			
C. C. engesseri	2	?	0	0	0	0	1	2	0	?	?	?	0	0	2	0	0	-	0	1	0	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?
C. C. sariae	0	?	0	0	0	0	1	2	0	?	?	?	1	0	0	0	-	0	?	0																																

	Species	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102																																																																																																																																															
0	<i>D. franciscus</i>	?	?	1	?	1	0	0	0	?	?	0	0	1	0	1	2	0	0	2	0	0	?	?	0	0	?	?	0	1	1	0	0	1	0	0	0	0	0	1	1	0	0	0	2	2	2	0	2	1	2	1																																																																																																																																														
0	<i>H. D. castelnoi</i>	?	1	?	1	0	0	1	2	0	?	?	0	0	0	1	0	1	2	0	0	0	?	?	0	0	?	?	0	1	0	1	0	0	1	2	0	0	0	0	1	1	0	0	2	2	0	2	2	0	-	-	1	2	1																																																																																																																																											
0	<i>H. decedens</i>	0	1	?	0	1	1	1	2	1	0	0	0	2	0	1	0	1	0	1	2	0	1	2	182	2	0	1	0	0	0	1	1	0	0	1	2	0	0	0	1	1	0	0	2	2	1	0	0	1	1	2	1																																																																																																																																													
1	<i>H. H. bijugatus</i>	1	1	1	1	1	0	2	0	0	0	1	0	1	0	1	0	1	0	1	0	2	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	1	1	0	0	2	1	0	0	1	0	0	0																																																																																																																																														
1	<i>H. H. lavacati</i>	1	1	1	1	?	081	1	082	1	1	0	0	1	0	1	0	1	2	0	0	0	2	0	0	0	0	0	0	1	1	0	0	1	1	1	0	0	0	1	1	1	0	0	2	1	0	1	1	1	0	0																																																																																																																																														
? 1	<i>H. H. respectu</i>	? 1	1	1	0	0	1	0	-	0	0	0	0	1	0	1	0	1	2	0	1	2	2	?	?	0	0	0	1	1	0	1	1	0	1	0	0	0	1	1	?	?	0	0	2	2	0	0	1	1	2	1																																																																																																																																														
1	<i>H. H. aguirrei</i>	1	1	0	1	?	0	1	2	0	1	0	1	0	1	0	1	0	1	2	1	1	2	0	0	0	0	0	0	1	0	1	0	1	182	0	0	0	0	1	0	0	0	1	2	182	0	0	1	0	2	1																																																																																																																																														
0	<i>H. adamsi</i>	0	1	1	1	0	1	2	1	?	0	1	0	1	0	1	0	1	2	1	0	0	2	1	0	0	1	0	1	1	?	0	0	0	1	1	?	0	0	0	?	?	0	2	2	081	0	0	1	0	2	1																																																																																																																																														
1	<i>H. H. nombrevillae</i>	1	1	0	1	1	2	0	2	2	1	1	1	1	1	0	1	1	0	1	2	182	0	0	0	0	0	0	1	0	0	1	1	182	1	0	0	0	0	1	1	0	1	2	182	0	0	1	0	2	1																																																																																																																																															
? 1	<i>H. thaleri</i>	? 1	?	1	?	?	1	?	?	?	?	1	1	2	0	1	0	0	1	2	0	?	?	182	0	0	?	?	0	1	1	0	0	?	1	1	0	0	1	1	?	?	0	0	2	0	0	1	1	1	0	1																																																																																																																																														
1	<i>H. H. aragonensis</i>	1	1	0	1	1	2	0	2	1	1	0	0	2	0	1	0	1	1	182	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	1	1	2	1	0	1	1	0	1	1	0	1	1																																																																																																																																														
1	<i>H. H. baixosi</i>	1	1	0	1	2	1	2	2	0	1	0	0	1	0	1	0	1	2	0	1	1	182	0	0	0	0	0	0	1	0	1	1	0	0	1	1	0	0	1	1	1	0	2	182	0	0	2	1	0	1	1																																																																																																																																														
? 1	<i>H. H. mediterraneus</i>	? 1	0	1	1	2	1	2	2	1	081	0	2	0	1	1	0	2	0	1	?	?	2	0	0	0	?	?	0	?	0	?	?	?	182	?	0	0	?	0	1	1	1	0	?	2	0	0	2	1	1	2	1																																																																																																																																													
0	<i>H. H. H. moralesi</i>	0	1	0	1	2	1	2	2	?	1	1	2	0	1	0	1	0	1	2	0	0	2	0	0	0	0	2	0	1	0	1	0	1	2	0	1	0	0	1	1	0	0	2	1	0	0	1	0	2	1																																																																																																																																															
1	<i>H. H. peralensis</i>	1	1	0	1	0	2	0	2	1	1	0	0	0	0	1	0	1	0	1	182	0	0	0	0	0	0	0	0	0	1	1	0	0	1	1	0	0	1	1	1	0	0	2	182	0	0	2	1	0	1	1																																																																																																																																														
1	<i>H. H. adroveri</i>	1	1	0	1	2	?	2	?	1	1	1	1	0	1	0	1	1	2	0	0	0	?	?	0	0	?	?	0	0	2	1	1	1	0	1	0	0	1	1	1	1	1	0	1	1	0	0	1	0	1	1																																																																																																																																														
1	<i>R. R. schaubi</i>	1	1	?	1	2	1	2	1	?	1	1	2	0	1	0	1	1	2	0	0	0	1	0	081	2	0	?	?	2	1	1	0	0	0	1	1	0	0	1	?	?	1	1	0	2	0	?	?	2	0	0	2	0																																																																																																																																												
0	<i>R. R. eupareus</i>	0	1	?	1	2	1	2	?	?	1	1	2	0	1	0	1	1	0	1	2	1	1	2	182	2	0	0	0	0	0	0	0	0	1	1	0	0	1	1	0	0	0	1	2	0	0	1	0	1	0	0																																																																																																																																														
0	<i>C. meini</i>	0	0	-	1	1	0	0	0	0	1	0	0	0	0	1	0	1	0	1	2	1	2	182	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	2	0	0	1	0	0	1																																																																																																																																														
0	<i>C. jotae</i>	0	0	-	1	1	1	0	0	0	0	0	0	0	0	1	0	1	0	1	2	1	2	2	0	1	1	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	1	2	0	0	1	1	0	0	0																																																																																																																																													
0	<i>C. salsaniensis</i>	0	1	1	0	1	1	0	0	0	1	0	0	0	0	0	1	0	1	1	1	1	2	2	0	1	0	0	1	0	0	1	0	0	1	0	0	0	0	0	0	0	1	0	1	2	0	0	1	1	0	0																																																																																																																																														
0	<i>C. versteegi</i>	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	?	0	0	1	2	2	?	0	1	0	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	?																																																																																																																																												
? ? ? ? ? ? ? ? ? ?	<i>C. kasapligilli</i>	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ? ? ? ? ? ? ? ? ?	? ?

Appendix 10.2

Appendix 10.2.

Information about the occurrences of the species of Cricetodontini used to discuss the paleobiogeography and evolutionary history of the tribe. The references on the sixth column gather the papers where the occurrence is reported and those which provided the date of the locality. Abbreviations: A. Minor, Asia Minor; SW. Europe, southwestern Europe; C. Europe, central Europe; E. Europe, eastern Europe.

Species	Locality	Age	Country	Region	References
<i>B. bayraktepenis</i>	Bayraktepe I	MN7/8	Turkey	A. Minor	Ünay, 1980; Grandjean, 2004
<i>B. dardanellensis</i>	Bayraktepe II	MN9	Turkey	A. Minor	Ünay, 1980; Wessels 2009
<i>B. eskihissarensis</i>	Yeni Eskihisar	Middle Miocene	Turkey	A. Minor	Ünay, 1980; Rummel, 1998
<i>B. hellenicus</i>	Hayranlı 1	MN10/MN11	Turkey	A. Minor	De Bruijn et al., 2013
<i>B. hellenicus</i>	Düzyayla	MN12	Turkey	A. Minor	De Bruijn et al., 2013
<i>B. hellenicus</i>	Samos Island	MN12	Greece	A. Minor	Freudenthal, 1970; Kostopoulos, 2009
<i>B. nikosi</i>	Biodrak	MN10	Greece	A. Minor	De Bruijn, 1976; Koufos, 2013
<i>B. ozansoy</i>	Bayraktepe I	MN7/8	Turkey	A. Minor	Ünay, 1980; Grandjean, 2004
<i>B. ozansoy</i>	Bagıçı	MN7/8	Turkey	A. Minor	Ünay, 1980; Grandjean, 2004
<i>B. pikermiensis</i>	pikermi (Chomateri)	MN12	Greece	A. Minor	De Bruijn, 1976; Koufos, 2013
<i>B. pikermiensis</i>	Altıntaş	MN9	Anatolia	A. Minor	De Bruijn et al., 2013
<i>C. albanensis</i>	Buenaventura 3	MN7/8 or MN9	Spain	SW. Europe	García-Moreno, 1988
<i>C. albanensis</i>	La Parrilla	MN7/8 or MN9	Spain	SW. Europe	García-Moreno, 1988
<i>C. albanensis</i>	La Grive-Saint Alban	MN7/8	France	SW. Europe	Mein and Freudenthal, 1971a
<i>C. aliveriensis</i>	Aliveri	MN4	Greece	A. Minor	Klein Hofmeijer and De Bruijn, 1988
<i>C. aureus</i>	Vieux-Collonges	MN5	France	SW. Europe	Mein and Freudenthal, 1971b
<i>C. aureus</i>	Castelnov-Barbieu	MN5	France	SW. Europe	Bulut et al., 1992
<i>C. aureus</i>	Laimering 3	MN6	Germany	C. Europe	Heissig, 2006
<i>C. aureus</i>	Laimering 5	MN7/8	Germany	C. Europe	Heissig, 2006
<i>C. aureus</i>	Pont levoy-Theney	MN5	France	SW. Europe	Daxner-Höck, 2003
<i>C. aureus</i>	Noyant-Sous-Le-Lude	MN5	France	SW. Europe	Ginsburg, 2001
<i>C. aff. aureus</i>	Petersbuch 38	MN6	Germany	C. Europe	Rummel, 2000
<i>C. aff. aureus</i>	Petersbuch 33	MN6	Germany	C. Europe	Rummel, 2000
<i>C. aff. aureus</i>	Petersbuch 32	MN6	Germany	C. Europe	Rummel, 2000
<i>C. aff. aureus</i>	Petersbuch 39	MN6	Germany	C. Europe	Rummel, 2000
<i>C. aff. aureus</i>	Steinberg	MN6	Germany	C. Europe	Bolliger, 1994
<i>C. aff. aureus</i>	Goldberg	MN6	Germany	C. Europe	Daxner-Höck, 2003
<i>C. aff. aureus</i>	Rümikon	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Untere-wagenburg	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Wiezholz	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Uzwill-Nutzenbuech	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Camentobel	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Rutzentobel 550m	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Schwamendingen	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. aff. aureus</i>	Sagentobel	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. cf. aureus</i>	Laimering 4a	MN6	Germany	C. Europe	Heissig, 2006
<i>C. cf. aureus</i>	Laimering 4b	MN6	Germany	C. Europe	Heissig, 2006
<i>C. cf. aureus</i>	Blanquatière 3	MN5	France	SW. Europe	Aguilar, 2010
<i>C. bolligeri</i>	Petersbuch 10	MN8	Germany	C. Europe	Rummel, 2000
<i>C. candirensis</i>	Berdik-1	middle/late Miocene	Turkey	A. Minor	De Bruijn et al., 2003
<i>C. candirensis</i>	Candir	MN5	Turkey	A. Minor	De Bruijn et al., 2003
<i>C. cariensis</i>	Sarıçay	MN7	Turkey	A. Minor	Sen and Ünay, 1979
<i>C. caucasicus</i>	Belometchetskaya	late MN6	Georgia	A. Minor	De Bruijn et al., 1992
<i>C. engesseri</i>	Mettlen-Weid	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. engesseri</i>	Schauenberg-Langriet	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. engesseri</i>	Chräzerentobel 523m	MN7	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. engesseri</i>	Chräzerentobel 505m	MN7	Swiss	C. Europe	Rummel and Kälin, 2003
<i>C. cf. engesseri</i>	Kirrborg-Tongrube	late MN6	Germany	C. Europe	Seehuber, 2008
<i>C. cf. engesseri</i>	Kirrborg-Sandgrube	late MN6	Germany	C. Europe	Seehuber, 2008
<i>C. hungaricus</i>	Hasznos	MN6	Hungary	E. Europe	De Bruijn et al., 1993
<i>C. cf. hungaricus</i>	Zambal	MN6	Turkey	A. Minor	Sen and Ünay, 1979
<i>C. jotae</i>	Povóa de Santarem	MN6	Portugal	SW. Europe	Antunes and Mein, 1977
<i>C. jotae</i>	Chloës	MN6	Portugal	SW. Europe	Antunes and Mein, 1977
<i>C. jotae</i>	Arroyo del Val 6	MN6	Spain	SW. Europe	Mein and Freudenthal, 1971a
<i>C. jotae</i>	Manchones	MN6, biozone G2	Spain	SW. Europe	De Bruijn et al., 1992
<i>C. jotae</i>	Povóa de Santarem	MN6	Portugal	SW. Europe	Antunes and Mein, 1977

Species	Locality	Age	Country	Region	References
<i>C. jotae</i>	Las Planas 5K	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Las Planas 5L	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Borjas	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Valalto 1	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Las Planas 5C	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Las Planas 5B	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Toril 1	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Alcocer 3	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Alcocer 2	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Alcocer 1	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jotae</i>	Villafeliche 9	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. jumaensis</i>	Schmidrüti	MN6	Swiss	C. Europe	Kálin and Kempf, 2009
<i>C. jumaensis</i>	Petersbuch 35	MN8	Germany	C. Europe	Rummel, 2001
<i>C. jumaensis</i>	Petersbuch 18	MN8	Germany	C. Europe	Prieto, 2007
<i>C. jumaensis</i>	Petersbuch 6	MN8	Germany	C. Europe	Prieto, 2007
<i>C. jumaensis</i>	Petersbuch 48	MN8	Germany	C. Europe	Prieto, 2007
<i>C. ? jumaensis</i>	Anwill	MN8	Swiss	C. Europe	Engesser, 1972
<i>C. kasapligili</i>	Keseköy	MN3	Turkey	A. Minor	De Bruijn et al., 1993
<i>'C'. klariankae</i>	Felsőtárkány-Felnémet	MN7/8	Hungary	E. Europe	Hír, 2007; Hír and Kókay, 2010
<i>'C'. ? klariankae</i>	Gaweinstal	MN9	Austria	C. Europe	Harzhauser et al., 2011; Prieto et al., 2014
<i>'C'. fandli</i>	Gratkorn	MN9	Austria	C. Europe	Prieto et al., 2010
<i>C. meini</i>	Mühlbach2	late MN5	Austria	C. Europe	Daxner-Höck, 2003
<i>C. meini</i>	Grund	late MN5	Austria	C. Europe	Daxner-Höck, 2003
<i>C. meini</i>	Komotini	MN4/MN5	Greece	A. Minor	Daxner-Höck, 2003; Koufos, 2013
<i>C. meini</i>	Petersbuch 41	MN5	Germany	C. Europe	Rummel, 2000
<i>C. meini</i>	Ebershausen	MN5	Germany	C. Europe	Bolliger, 1994
<i>C. meini</i>	Ziemetshausen	MN5	Germany	C. Europe	Bolliger, 1994
<i>C. meini</i>	Burg-Balzhausen	MN5	Germany	C. Europe	Seehuber, 2008
<i>C. meini</i>	Vieux-Collonges	MN5	France	SW. Europe	Mein and Freudenthal, 1971b
<i>C. meini</i>	Castelnov-Barbieu	MN5	France	SW. Europe	Bulot et al., 1992
<i>C. meini</i>	Unterneufl 1b	MN6	Germany	C. Europe	Heissig, 2006
<i>C. meini</i>	Unterneufl 1a	MN6	Germany	C. Europe	Heissig, 2006
<i>C. meini</i>	Mülbach1	late MN5	Austria	C. Europe	Daxner-Höck, 2003
<i>C. meini</i>	Antonios	MN4/MN5	Greece	A. Minor	Vasileiadou and Koufos, 2005; Koufos,
<i>C. meini</i>	Litke 1	MN5	Hungary	E. Europe	Hír, 2014
<i>C. meini</i>	Litke2	MN5	Hungary	E. Europe	Hír, 2014
<i>C. meini</i>	Popovac	MN5	Serbia	E. Europe	Alaburic and Markovic, 2010
<i>C. meini</i>	Mala Miliva	MN5	Serbia	E. Europe	Markovic, 2003
<i>C. meini</i>	Lararevac	MN6	Serbia	E. Europe	Markovic, 2008, 2010
<i>C. aff. meini</i>	Belometchetskaya	late MN6	Georgia	A. Minor	Pickford et al., 2000
<i>C. aff. meini</i>	Edelbeuren-Maurerkopf	MN5	Germany	C. Europe	Daxner-Höck, 2003
<i>C. aff. meini</i>	Strakonice	MN5	Czech Republic	E. Europe	Daxner-Höck, 2003
<i>C. aff. meini</i>	Wannentobel	MN5-6	Germany	C. Europe	Daxner-Höck, 2003
<i>C. aff. meini</i>	Gallenbach 2a	MN6	Germany	C. Europe	Bolliger, 1994
<i>C. cf. meini</i>	Hohenraunau	early MN6	Germany	C. Europe	Seehuber, 2008
<i>C. cf. meini</i>	Tuzluca-Kars	MN4-MN5	Armenia	A. Minor	Sen et al., 2011
<i>C. cf. meini</i>	Blanquatière 3	MN5	France	SW. Europe	Aguilar, 2010
<i>C. nievei</i>	Toril 3B	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2014
<i>C. nievei</i>	Toril 3A	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2014
<i>C. nievei</i>	Las Planas 5H	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2014
<i>C. nievei</i>	Toril 2	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2014
<i>C. pasalarensis</i>	Pasalar	late MN6	Turkey	A. Minor	Peláez-Campomanes and Daams, 2002
<i>C. pasalarensis</i>	Hosköy	MN6	Turkey	A. Minor	Ünay, 1980
<i>C. pasalarensis</i>	Mürefte	MN6	Turkey	A. Minor	Ünay, 1980
<i>C. cf. pasalarensis</i>	Zambal	MN6	Turkey	A. Minor	De Bruijn et al., 2013

Species	Locality	Age	Country	Region	References
<i>C. sansaniensis</i>	Niderwis	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. sansaniensis</i>	Zeglingen	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. sansaniensis</i>	Sansan	MN6	France	SW. Europe	De Bruijn et al., 1992
<i>C. sansaniensis</i>	Valalto 2C	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. sansaniensis</i>	Valato 2B	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. sansaniensis</i>	Armantes 7	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. sansaniensis</i>	Las Umbrías 21	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. sansaniensis</i>	Las Umbrías 22	MN6	Spain	SW. Europe	López-Guerrero et al., in press
<i>C. sansaniensis</i>	Duredos 1	MN7/8	Spain	SW. Europe	García-Moreno, 1987
<i>C. sansaniensis</i>	Simancas 2	MN7/8	Spain	SW. Europe	García-Moreno, 1987
<i>C. cf. sansaniensis</i>	Kirrberg-Tongrube	late MN6	Germany	C. Europe	Seehuber, 2008
<i>C. cf. sansaniensis</i>	Tschöplhof	MN6	Germany	C. Europe	Rummel and Kälin, 2003
<i>C. cf. sansaniensis</i>	Göttschlag	MN6	Germany	C. Europe	Bolliger, 1994
<i>C. cf. sansaniensis</i>	Oeschgraben	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. cf. sansaniensis</i>	Mettlen 4	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. cf. sansaniensis</i>	Mühlrüti Pt 806	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. cf. sansaniensis</i>	Tschöplhof 720m	MN6	Swiss	C. Europe	Kälin and Kempf, 2009
<i>C. cf. sansaniensis</i>	Armantes 7	MN6	Spain	SW. Europe	Freudenthal, 1966
<i>C. cf. sansaniensis</i>	Petersbuch 31	MN7	Germany	C. Europe	Rummel, 2000
<i>C. cf. sansaniensis</i>	Laimering 1b	MN7/8	Germany	C. Europe	Bolliger 1994
<i>C. cf. sansaniensis</i>	Laimering 4b	MN7/8	Germany	C. Europe	Bolliger 1994
<i>C. cf. sansaniensis</i>	Imenberg	MN8	Swiss	C. Europe	Bolliger 1994
<i>C. soriae</i>	Las Umbrías 20	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Planas 4C	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Umbrías 19	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Planas 4B	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Planas 4A	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Umbrías 16	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Las Umbrías 11	MN5	Spain	SW. Europe	López-Guerrero et al., 2013
<i>C. soriae</i>	Somosaguas	MN5, biozone E	Spain	SW. Europe	Hernández-Fernández et al., 2006
<i>C. tobieni</i>	Horlak 1a	MN3-MN4	Turkey	A. Minor	Tobien, 1978
<i>C. tobieni</i>	Horlak 1b	MN3-MN4	Turkey	A. Minor	De Bruijn et al., 1993
<i>C. tobieni</i>	Kaplangi 1-2	MN4	Turkey	A. Minor	De Bruijn et al., 2013
<i>C. verteegi</i>	Kilçak 3a	MN1	Turkey	A. Minor	De Bruijn et al., 1993
<i>C. verteegi</i>	Kilçak 0	MN1	Turkey	A. Minor	De Bruijn et al., 1993
<i>C. verteegi</i>	Kargi 2	Oligocene	Turkey	A. Minor	De Bruijn et al., 2013
<i>C. aff. versteegi</i>	Kilçak 3b	MN1	Turkey	A. Minor	De Bruijn et al., 1993
<i>C. volkeri</i>	Dingshanyanchi	MN7/8	China	Asia	Wu et al., 2009; Qiu and Qiu, 2013
<i>C. wanhei</i>	onglinzhuan/Xiacaowang fr	early Miocene	China	Asia	Qiu and Qiu, 2013
<i>C. wanhei</i>	Sihong	MN3-MN4	China	Asia	Qiu, 2010; Qiu and Qiu, 2013
<i>D. anatolicus</i>	Kilçak 0-3	MN1	Turkey	A. Minor	De Bruijn et al., 2013
<i>D. anatolicus</i>	Kargi 2	Oligoceno	Turkey	A. Minor	De Bruijn et al., 2013
<i>D. intermedius</i>	Harami1	MN2	Turkey	A. Minor	De Bruijn et al., 2013
<i>H. adroveri</i>	Jun-2C	MN12	Spain	SW. Europe	García-Alix, et al., 2008
<i>H. adroveri</i>	Casa del Acero	MN12	Spain	SW. Europe	Agustí, 1986; García-Alix., 2008
<i>H. adroveri</i>	CR-17	MN12	Spain	SW. Europe	García-Alix et al., 2008
<i>H. aguirrei</i>	Carrilanga 1	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. aguirrei</i>	Nombrevilla 2	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. aguirrei</i>	Solera	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. aguirrei</i>	Escobosa de Calatañazor	MN7/8	Spain	SW. Europe	López-Martínez et al., 1977
<i>H. cf. aguirrei</i>	Nombrevilla 4	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. cf. aguirrei</i>	Nombrevilla 3	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. cf. aguirrei</i>	Paje 2	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. cf. aguirrei</i>	Paje 1	MN7/8	Spain	SW. Europe	López-Guerrero et al., in press

Species	Locality	Age	Country	Region	References
<i>H. aragonensis</i>	ECM	MN9	Spain	SW. Europe	Alba et al., 2006; Carmona et al., 2011
<i>H. aragonensis</i>	Pedregueras 2A	MN9	Spain	SW. Europe	Mein and Freudenthal, 1971a
<i>H. aragonensis</i>	Pedregueras 1A	MN9	Spain	SW. Europe	Mein and Freudenthal, 1971a
<i>H. aragonensis</i>	Pedregueras 2C	MN9	Spain	SW. Europe	Daams et al., 1999; Freudenthal, 1966
<i>H. aragonensis</i>	Torremormojón 4	MN9	Spain	SW. Europe	Álvarez-Sierra, 1983
<i>H. cf. aragonensis</i>	Nombrevilla 19	MN9	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. cf. aragonensis</i>	Nombrevilla 14	MN9	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. cf. aragonensis</i>	Nombrevilla 13	MN9	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. cf. aragonensis</i>	Nombrevilla 22	MN9	Spain	SW. Europe	López-Guerrero et al., in press
<i>H. baixasi</i>	Lo Fournas 16-M	MN11	France	SW. Europe	Aguilar et al., 2007
<i>H. bijugatus</i>	La Grive-Saint Alban, L3	MN8	France	SW. Europe	López-Antoñanzas and Mein, 2009
<i>H. bijugatus</i>	La Grive-Saint Alban, L5'	MN8	France	SW. Europe	López-Antoñanzas and Mein, 2009
<i>H. cf. bijugatus</i>	Bullendorf	MN9	Austria	C. Europe	Daxner-Höck, 1996
<i>H. castelnovi</i>	Castelnou 6	MN6?	France	SW. Europe	Aguilar et al., 1994
<i>H. daamsi</i>	Abocador de can Mata	MN7/8 or MN9	Spain	SW. Europe	Alba et al., 2006
<i>H. daamsi</i>	Can Missert	MN9	Spain	SW. Europe	Agustí et al., 2005; Gómez Cano et al., 2011
<i>H. decedens</i>	La Grive-Saint Alban, L5'	MN7/8	France	SW. Europe	López-Antoñanzas and Mein, 2011
<i>H. decedens</i>	La Grive-Saint Alban, M	MN7/8	France	SW. Europe	López-Antoñanzas and Mein, 2011
<i>H. decedens</i>	Abocador de can Mata	MN7/8 or MN9	Spain	SW. Europe	Alba et al., 2006
<i>H. dispectus</i>	Hostalets de Pierola	MN8-MN9	Spain	SW. Europe	Agustí, 1980
<i>H. dispectus</i>	Creu Conill 20	MN9	Spain	SW. Europe	Casanovas et al., 2006
<i>H. dispectus</i>	Creu Conill 22	MN9	Spain	SW. Europe	Casanovas et al., 2006
<i>H. lavocati</i>	San Quirze	MN7	Spain	SW. Europe	López-Antoñanzas and Mein, 2009
<i>H. lavocati</i>	Nombrevilla 2	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. lavocati</i>	Hostalets de Pierola	MN7/8	Spain	SW. Europe	Agustí, 1980
<i>H. cf. lavocati</i>	Nombrevilla 4	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. cf. lavocati</i>	Nombrevilla 3	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2008
<i>H. mediterraneus</i>	Montredon	MN10	France	SW. Europe	Aguilar, 1982
<i>H. moralesi</i>	Batallones 10	MN10	Spain	SW. Europe	López-Antoñanzas et al., 2010
<i>H. nombrevillae</i>	Nombrevilla 9	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2009
<i>H. nombrevillae</i>	Cañada 10	MN9	Spain	SW. Europe	López-Guerrero et al., 2011
<i>H. nombrevillae</i>	Nombrevilla 1	MN9	Spain	SW. Europe	Freudenthal 1966; Daams et al., 1999
<i>H. cf. nombrevillae</i>	Nombrevilla 10	MN7/8	Spain	SW. Europe	López-Guerrero et al., 2009
<i>H. cf. nombrevillae</i>	Cañada 11	MN9	Spain	SW. Europe	López-Guerrero et al., 2011
<i>H. cf. nombrevillae</i>	Cañada 9	MN9	Spain	SW. Europe	López-Guerrero et al., 2011
<i>H. cf. nombrevillae</i>	Cañada 8	MN9	Spain	SW. Europe	López-Guerrero et al., 2011
<i>H. peralensis</i>	Peralejos C	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. peralensis</i>	Peralejos D	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. peralensis</i>	Peralejos B	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. peralensis</i>	Peralejos 4	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. peralensis</i>	Masia del BarboA	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. peralensis</i>	Masia del BarboB	MN10	Spain	SW. Europe	Van de Weerd, 1976; Van Dam et al., 2006
<i>H. thaleri</i>	Can Llobateres	MN9	Spain	SW. Europe	Hartenberger, 1965; Casanovas-Vilar and Agustí, 2007
<i>H. thaleri</i>	Riu Ripoll	MN9	Spain	SW. Europe	Agustí and Gilbert, 1979
<i>R. europaeus</i>	Villalba atal rio	MN14	Spain	SW. Europe	Adrover, 1988
<i>R. europaeus</i>	Layna	MN14	Spain	SW. Europe	García-Alix et al., 2008
<i>R. europaeus</i>	Rambla seca-1,-2,-3	MN14	Spain	SW. Europe	García-Alix et al., 2008
<i>R. cf. europaeus</i>	Peralejos E	MN14	Spain	SW. Europe	Adrover, 1988

References

Adrover, R., P. Mein, and E. Moissenet. 1988. Contribución al conocimiento de la fauna de roedores del Plioceno de la región de Teruel. *Teruel* 79:91–151.

Aguilar, J. P. 1982. Contributions à l'étude des Micromammifères du gisement Miocène supérieur

- de Montredon (Hérault) 2. les rongeurs. *Palaeovertebrata* 12:75–140.
- Aguilar, J. P., M. Calvet, and J. Michaux. 1994. Les rongeurs de Castelnou 6 (Pyrénées-orientales, France) et les corrélations entre faunes ibériques et françaises au Miocène moyen. *Neues Jahrbuch Fur Geologie Und Palaontologie-Abhandlungen* 192:109–131.
- Aguilar, J. P., V. Lazzari, J. Michaux, M. Sabatier, and M. Calvet. 2007. Lo Fournas 16-M (Miocène supérieur) et Lo Fournas 16-P (Pliocène moyen), nouvelles localités karstiques à Baixas, Sud de la France. Partie I- Description et implications géodynamiques. *Géologie de la France* 1:55–62.
- Aguilar J. P., J. Michaux, and V. Lazzari. 2010. Nouvelles faunes de rongeurs (Mammalia, Rodentia) d'âge Miocène moyen en Languedoc-Roussillon (Sud de la France); biostratigraphie et corrélations. *Geodiversitas* 32:501–513.
- Agustí, J. 1980. La asociación de *Hispanomys* y *Cricetodon* (Rodentia, Mammalia) en el Mioceno superior del Vallès-Penedès (Cataluña, España). *Acta Geologica Hispanica* 15:51–60.
- Agustí, J. 1986. Nouvelles espèces de Cricétidés vicariantes dans le Turolien moyen de Fortuna (Province de Murcia, Espagne). *Geobios* 19:5–11.
- Agustí, J., and J. Gilbert. 1979. Nuevos datos sobre los Roedores e Insectívoros (Mammalia) del mioceno superior del Vallés (Cataluña, España). *Paleontologia i Evolució* 14:17–20.
- Agustí, J., I. Casanovas-Vilar, and M. Furió. 2005. Rodents, insectivores and chiropterans (Mammalia) from the late Aragonian of Can Missert (middle Miocene, Vallés-Penedés Basin, Spain). *Geobios* 38:575–583.
- Alaburic, S., and Z. Markovic. 2010. New records of mammal remains from the Miocene sediments of the cement mine in Popovac (Serbia). *Bulletin of the Natural History Museum* 3:95–103.
- Alba, D. M., Moyá-Solá, S., Casanovas-Vilar, I., Galindo, J., Robles, J.M., Rotgers, C., Furió, M., Angelone, C., Köhler, M., Garcés, M., Cabrera, L., Almécija, S., Obrado, P. 2006. Los vertebrados fosiles del Abocador de Can Mata (els Hostalets de Pierola, l'Anoia, Cataluna), una sucesión de localidades del Aragoniense superior (MN6 y MN7+8) de la cuenca del Vallès-Penedès. *Campanas 2002-2003, 2004 y 2005. Estudios Geológicos* 62: 295–312.
- Álvarez-Sierra, M. A. 1983. Paleontología y bioestratigrafía del Mioceno superior del sector central de la Cuenca del Duero. Estudio de los micromamíferos de la serie de Torremormojón (Palencia). Memoria de grado de licenciatura. Universidad Complutense de Madrid, Madrid, 188 pp.

- Antunes, M., and Mein, P., 1977. Contributions à la paléontologie du miocène moyen continental du bassin du Tage. III. Mammifères-póvoa de Santarem, Pero Filho et Choes (Secorio) conclusiones generales. *Ciências da Terra (UNL)* 2:143–165.
- Bolliger, T. V., 1994. Die Obere SüBwassermolasse in Bayern und der Ostschweiz: bio-und lithostratigraphische Korrelationen. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und histor. Geologie* 34:09–144.
- Bulot, C., L. Ginsburg, and P. Tassy. 1992. Le gisement à mammifères miocènes de Castelnau-d'Arbieu (Gers). Données nouvelles et implications biostratigraphiques. *Comptes rendus de l'Académie des sciences, Séries IIA* 314:533–537.
- Carmona, R., D.M. Alba, I. Casanovas-Vilar, M. Furió, M. Garcés, J.V. Bertó Mengual, J. Galindo, and À.H. Luján. 2011. Intervención paleontológica en la serie del Mioceno Superior del Ecoparc de Can Mata (cuenca del Vallès-Penedès, NE de la península Ibérica) pp. 65–74 in A. Pérez-García, F. Gascó, J.M. Gasulla and F. Escaso (eds), *Viajando a mundos pretéritos*. Morella: Ayuntamiento de Morella.
- Casanovas-Vilar, I., and J. Agustí. 2007. Ecogeographical stability and climate forcing in the Late Miocene (Vallesian) rodent record of Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology* 248:169–189.
- Casanovas-Vilar, I., M. Furió, and J. Agustí. 2006. Rodents, Insectivores and Paleoenvironment associated to the first-appearing Hipparionine horses in the Vallès-Penedès Basin (North-eastern Spain). *Beiträge zur Paläontologie* 30:89–107.
- Daams, R., A. J. Van der Meulen, M. A. Álvarez-Sierra, P. Peláez-Campomanes, J. P. Calvo, M. A. Alonso Zarza, and W. Krijgsman. 1999. Stratigraphy and sedimentology of the Aragonian (Early to Middle Miocene) in its type area (North-Central Spain). *Newsletters on Stratigraphy* 37:103–139.
- Daxner-Höck, G. 2003. *Cricetodon meini* and other rodents from Muehlbach and Grund, Lower Austria (Middle Miocene, late MN5). *Annalen des Naturhistorischen Museums in Wien* 104A:267–291.
- Daxner-Höck, G., V. Fahlbusch, L. Kordos, and W. Wu. 1996. The late Neogene cricetid rodent genera *Neocricetodon* and *Kowalskia*; pp. 220–226 in Bernor, R. L, V. Fahlbusch, H. W. Mittmann (eds.), *The evolution of western Eurasian Neogene Mammal Faunas*. Columbia University Press. New York.
- De Bruijn, H. 1976. Vallesian and Turolian rodents from Biotia, Attica and Rhodes (Greece). *Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen* 79:361–384.

- De Bruijn, H., and G. Saraç. 1992. Early Miocene rodent faunas from the Eastern Mediterranean area. Part II. *Mirabella* (Paracricetodontinea, Muroidea). Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 95:25–40.
- De Bruijn, H., E. Ünay, and K. Hordijk. 2013. A review of the Neogene Succession of the Muridae and Dipodidae from Anatolia, with special reference to taxa Known from Asia and/or Europe; pp. 566–582 in X. Wang, L. Flynn and M. Fortelius (eds.), Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology. Columbia University Press, New York.
- De Bruijn, H., V. Fahlbusch, G. Saraç, and E. Ünay. 1993. Early Miocene Rodent Faunas from the Eastern Mediterranean Area 3. The Genera *Deperetomys* and *Cricetodon* with a Discussion of the Evolutionary History of the Cricetodontini. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, Series B 96:151–216.
- De Bruijn, H., L. W. Van den Hoek Ostende, M. R. Kristkoiz-Boon, K. Theocharopoulos, and E. Ünay. 2003. Rodents, lagomorphs and insectivores, from the middle Miocene hominoid locality of Çandir (Turkey). Courier Forschungsinstitut Senckenberg 240:51–87.
- Engesser, B. 1972. Die obermiozäne Säugetier Fauna von Anwil (Baselland). Ph.D. dissertation, Tätigk. Naturwiissenschaftlichen Fakultät der Universität Basel, Basel, 363 pp.
- Freudenthal, M. 1966. On Mammalian Fauna of *Hipparion*-Beds in Calatayud-Teruel Basin (Prov. Zaragoza Spain) I. Genera *Cricetodon* and *Ruscinomys* (Rodentia). Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B 69:298–317.
- Freudenthal, M. 1970. A new *Ruscinomys* (Mammalia, Rodentia) from the late Tertiary (Pikermian). American Museum Novitates 2402:1–10.
- García-Alix, A., 2006. Bioestratigrafía de los depósitos continentales de la transición Mio-Plioceno de la cuenca de Granada. Ph.D. dissertation, Universidad de Granada, 429 pp.
- García-Alix, A., Minwer-Barakat, R., Martín-Suarez, E., Freudenthal, M., 2007. New data on Mio-Pliocene Sciuridae (Rodentia, Mammalia) from southern Spain. Comptes Rendus Palevol 6:269–279.
- García-Alix, A., Minwer-Barakat, R., Martín-Suárez, E., Freudenthal, M., 2008. Muridae from the Mio-Pliocene boundary in the Granada Basin (southern Spain). Biostratigraphic and phylogenetic implications. Neues Jahrbuch für Geologie und Paläontologie Abhandlungen 248:183–215.
- García-Moreno, E. 1987. Roedores y Lagomorphos del Mioceno de la Zona central de la Cuenca del Duero. Sistemática, Bioestratigrafía y Paleoecología. Ph.D. dissertation, Universidad Complutense, Madrid, Madrid, 258 pp.

- Gómez Cano, A. R., M. Hernández Fernández, and M. A. Álvarez-Sierra. 2011. Biogeographic provincialism in rodent faunas from the Iberocccitanian Region (southwestern Europe) generates severe diachrony within the Mammalian Neogene (MN) biochronologic scale during the Late Miocene. *Palaeogeography, Palaeoclimatology, Palaeoecology* 307:193–204.
- Grandjean, E. 2004. Le genre *Byzantinia* (Rodentia, Cricetidae) du Miocène d’Anatolie centrale. Systématique et phylogénie. Mémoire de Maîtrise de Biologie des Populations et des Ecosystèmes Module Systématique, Evolution et Biodiversité. Université Pierre et Marie Curie, Paris, 37 pp.
- Ginsburg, L. 2001. Les faunes de mammifères terrestres du Miocène moyen des Faluns du bassin de Savigné-sur-Lathan (France). *Geodiversitas* 23:381–394.
- Hartenberger, J. L. 1965. Les Cricetidae (Rod.) de Can Llobateres (Néogène d’Espagne). *Bulletin De La Societe Geologique De France* 7:487–498.
- Harzhauser, M., G. Daxner-Höck, U. Göhlich, and D. Nagel. 2011. Complex faunal mixing in the early Pannonian palaeo-Danube Delta (Late Miocene, Gaweinstal, Lower Austria). *Annalen des Naturhistorischen Museums in Wien* 113:167–208.
- Heißig, K. 2006. Biostratigraphy of the “main bentonite horizon” of the Upper Freshwater Molas-se in Bavaria. *Palaeontographica Abteilung a-Palaeozoologie-Stratigraphie* 277:93–102.
- Hernández Fernández, M., J. A. Cárdena, J. Cuevas-González, O. Fesharaki, M. J. Salesa, B. Corrales, L. Domingo, J. Elez, P. López-Guerrero, N. Sala-Burgos, J. Morales, and N. López Martínez. 2006. Los yacimientos de vertebrados del Mioceno medio de Somosaguas (Pozuelo de Alarcón, Madrid): implicaciones paleoambientales y paleoclimáticas. *Estudios Geológicos* 62:263–294.
- Hír, J. 2007. *Cricetodon klariankae* n. sp. (Cricetodontini, Rodentia) from Felsőtárkány-Felnémet (Northern Hungary). *Fragmenta Paleontologica Hungarica* 24–25:16–24.
- Hír, J. 2013. Early and Middle Miocene (MN5–MN6) transitional rodent fauna from Litke (North Hungary, Nógrád County). *Fragmenta Paleontologica Hungarica* 30:101–137.
- Hír, J., and J. Kókay. 2010. A systematic study of the middle-late Miocene rodents and lagomorphs (Mammalia) of Felsőtárkány 3/8 and 3/10 (Northern Hungary) with stratigraphical relations. *Geodiversitas* 32:307–329.
- Kälin, D., and O. Kempf. 2009. High-resolution stratigraphy from the continental record of the Middle Miocene Northern Alpine Foreland Basin of Switzerland. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 254:177–235.

- Kostopoulos, D. S. 2009. The Pikermian event: Temporal and spatial resolution of the Turolian large mammal fauna in SE Europe. *Palaeogeography, Palaeoclimatology, Palaeoecology* 274:82–95.
- Koufos, G. D. 2013. Neogene Mammal Biostratigraphy and Chronology of Greece; pp. 595–628 in X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- Klein Hofmeijer, G., and H. De Bruijn. 1988. The mammals from the lower Miocene of Aliveri (Island of Evia, Greece) Part 8: The Cricetidae. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen Series B* 91:185–204.
- López-Antoñanzas, R., and P. Mein. 2009. First detailed description of *Hispanomys bijugatus* Mein & Freudenthal, 1971 (Rodentia, Cricetodontinae) from the Upper Aragonian of La Grive-Saint Alban (France): Biostratigraphical implications. *Geobios* 42:783–796.
- López-Antoñanzas, R., and P. Mein. 2011. First detailed description of *Hispanomys decedens* (Rodentia) from the Middle Miocene of La Grive-Saint Alban (France). *Swiss Journal of Geosciences* 104:345–353.
- López-Antoñanzas, R., P. Peláez-Campomanes, M. A. Álvarez-Sierra, and I. García-Paredes. 2010. New species of *Hispanomys* (Rodentia, Cricetodontinae) from the Upper Miocene of Batallones (Madrid, Spain). *Zoological Journal of the Linnean Society* 160:725–747.
- López-Guerrero, P., I. García-Paredes, and M. A. Álvarez-Sierra. 2013. Revision of *Cricetodon soriae* (Rodentia Mammalia), new data from the middle Aragonian (Middle Miocene) of the Calatayud-Daroca Basin (Zaragoza, Spain). *Journal of Vertebrate Paleontology* 33:169–184.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. 2014. New Cricetodontini from the middle Miocene of Europe: an example of mosaic evolution. *Bulletin of Geosciences* 89: 573–592.
- López-Guerrero, P., M. A. Álvarez-Sierra, I. García-Paredes, and P. Peláez-Campomanes. In press. Cricetodontini from the Calatayud-Daroca Basin (Spain): a taxonomical description and up-to-date of their stratigraphical distributions. *Comptes Rendus Palevol*. DOI: <http://dx.doi.org/10.1016/j.crpv.2014.07.002>.
- López-Guerrero, P., M. A. Álvarez Sierra, R. López-Antoñanzas, A. Oliver Pérez, and P. Peláez-Campomanes. 2008. Cricetodontini (Cricetidae, Rodentia) del Aragoniense superior (Mioceno medio) de Nombrevilla 2, (Zaragoza, España). *Palaeontologica Nova* 8:259–271.

- López-Guerrero, P., M. Álvarez-Sierra, I. García-Paredes, R. López-Antonanzas, y A. Oliver. 2009. Cricetodontini (Rodentia, Mammalia) from the Upper Aragonian and Lower Vallesian of the Toril-Nombrevilla Section (Middle and Upper Miocene, Calatayud-Daroca Basin, Zaragoza, Spain). *Journal of Vertebrate Paleontology* 29:161A–161A.
- López-Guerrero, P., I. García-Paredes, L. W. Van den Hoek Ostende, J. A. Van Dam, M. A. Álvarez-Sierra, V. Hernández-Ballarín, A. J. Van der Meulen, A. Oliver, y P. Peláez-Campomanes. 2011b. Cañada: a new micromammal succession from the lower Vallesian and Turolian of the Daroca area (Calatayud-Montalbán basin, Spain). *Estudios Geológicos* 67:443–453.
- López-Martínez, N., C. Sesé Benito, and J. L. Sanz García. 1977. La microfauna (Rodentia, Insectivora, Lagomorpha y Reptilia) de las fisuras del Mioceno Medio de Escobosa de Calatañazor (Soria, España). *Trabajos del Neógeno/Cuaternario* 8:47–73.
- Markovic, Z. 2003. The Miocene small mammals of Serbia, a review; pp. 393–398 in J.W.F. Reumer, and W. Wessels, W. (eds.), *Distribution and migration of tertiary mammals in Eurasia. A volume in honour of Hans de Bruijn-Deinsea*.
- Markovic, Z. 2008. Rodents of middle Miocene localities of Lazarevac village and Bele vode (central Serbia). *Bulletin of the Natural History Museum* 1:79–98.
- Markovic, Z., and M. Milivojevic. 2010. The Neogene Small Mammals from Serbia-Collection Methods and Results. *Bulletin of the Natural History Museum* 3:105–114
- Mein, P., and M. Freudenthal. 1971a. Une nouvelle classification des Cricetidae (Mam. Rod.) du Tertiaire d'Europe. *Scripta Geologica* 2:1–37.
- Mein, P., and M. Freudenthal. 1971b. Les Cricetidae (Mammalia, Rodentia) du Néogène Moyen de Vieux-Collonges. Part 1. Le genre *Cricetodon* Lartet, 1851. *Scripta Geologica* 5:1–51.
- Peláez-Campomanes, P. and R. Daams. 2002. Middle Miocene rodents from Pasalar, Anatolia, Turkey. *Acta Paleontologica Polonica* 47:125–132.
- Pickford, M. L. Gabuna, P. Mein, J. Morales, and B. Azanza. 2000. The Middle Miocene mammalian site of Belometchetskaya, North Caucasus: an important biostratigraphic link between Europe and China. *Geobios* 33:257–267.
- Prieto, J. 2007. Kleinsäuger-Biostratigraphie und Paläoökologie des höheren Mittelmiozäns (MN 8) Bayerns: Spaltenfüllungen der Fränkischen Alb und Lokalitäten der Oberen Süßwassermolasse im Vergleich. Ph.D. Dissertation, Ludwig Maximilians Universität Munich, 217 pp.

- Prieto, J., M. Bohme, and M. Gross. 2010. The cricetid rodents from Gratkorn (Austria, Styria): a benchmark locality for the continental Sarmatian *sensu stricto* (late Middle Miocene) in the Central Paratethys. *Geologica Carpathica* 61:419–436.
- Prieto, J., C. Angelone, I. Casanovas-Vilar, M. Gross, J. Hír, L. W. Van den Hoek Ostende, L. C. Maul, and D. Vasilyan. 2014. The small mammals from Gratkorn: an overview. *Palaeobiology and Palaeoenvironments* 94:135–162.
- Qiu, Z. D. 2010. Cricetids rodents from the early Miocene Xiacaowan formation, Sihong, Jiangsu. *Vertebrata Palasiatica* 48:27–47.
- Qiu, Z-D., and Z-X. Qiu. 2013. Early Miocene Xiejiahe and Sihong Fossil Localities and Their Faunas, Eastern China; pp. 595–628 in X. Wang, L. Flynn and M. Fortelius (eds.), *Fossil Mammals of Asia. Neogene Biostratigraphy and Chronology*. Columbia University Press, New York.
- Rummel, M. 1998. Die Cricetiden aus dem Mittel-und Obermiozän der Türkei. *Documenta naturae* 123:1–300.
- Rummel, M. 2000. Miocene Cricetodontini from Petersbuch near Eichstaett. The genus *Cricetodon* Lartet 1851. *Senckenbergiana lethaea* 80:149–171.
- Rummel, M. 2001. Ein neuer *Cricetodon* aus dem Miozän von Petersbuch bei Eichstätt. *Stuttgarter Beiträge zur Naturkunde B* 311:1–6.
- Rummel, M., and D. Kálin. 2003. Die Gattung *Cricetodon* (Mammalia, Rodentia) aus den Mitelmiozän der Schweizer Molasse. *Zitteliana* 43:123–141.
- Seehuber, U. 2008. Litho- und biostratigraphische Untersuchungen in der Oberen Süßwassermolasse in der Umgebung von Kirchheim in Schwaben. P.h.D. Dissertation, Ludwig-Maximilians-Universität, Munich, 359 pp.
- Sen, S., and E. Ünay. 1979. Sur quelques Cricetodontini (Rod.) du Miocène moyen d'Anatolie. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen* 82:293–301.
- Sen, S., P. O. Antoine, B. Varol, T. Ayyildiz, and K. Sözeri. 2011. Giant rhinoceros *Paraceratherium* and other vertebrates from Oligocene and middle Miocene deposits of the Kağızman-Tuzluca Basin, Eastern Turkey. *Naturwissenschaften* 98:407–423.
- Tobien, H. 1978. New species of Cricetodontini (Rod. Mam.) from the Miocene of Turkey. *Mainzer Geowissenschaftliche Mitteilungen Geologisches Landesamt Rheinland-Pfalz* 6:209–219.
- Ünay, E. 1980. The Cricetodontini (Rod. Mam.) from the Bayraktepe section (Çanakkale, Tur-

- key). Proceedings of the Koninklijke Nederlandse Akademie Van Wetenschappen, Series B 83:399–418.
- Van Dam, J., W. Krijgsman, H. A. Abels, M. A. Álvarez-Sierra, I. García-Paredes, P. López-Guerrero, P. Peláez-Campomanes y D. Ventrà. En prensa. Updated chronology for middle to late Miocene mammal sites of the Daroca area (Calatayud-Montalbán Basin), Spain. DOI: <http://dx.doi.org/10.1016/j.geobios.2014.07.002>
- Van Dam, J. A., H. Abdul Aziz, M. A. Álvarez-Sierra, F. J. Hilgen, L. W. Van den Hoek Ostende, L. J. Lourens, P. Mein, A. J. van der Meulen, and P. Peláez-Campomanes. 2006. Long-period astronomical forcing of mammal turnover. *Nature* 443:687–691.
- Van de Weerd, A. 1976. Rodent faunas of the Mio-Pliocene continental sediments of the Teruel-Alfambra region, Spain; pp 1–218 in C. W. Drooger (ed.), *Utrecht micropaleontological bulletins. Special issue 2*. University of Utrecht, Utrecht.
- Vasileiadou, K., and G. Koufos. 2005. The micromammals from the Early/Middle Miocene locality of Antonios, Chalkidiki, Greece. *Annales de Paléontologie* 91:197–225.
- Wessels, W. 2009. Miocene rodent evolution and migration. Muroidea from Pakistan, Turkey and Northern Africa. *Geologica Ultraiectina* 307:1–290.
- Wu, W. Y., J. Meng, J. Ye, X. J., Ni, S. D. Bi, and Y. P. Wei. 2009. The Miocene mammals from Dinshanyanchi Formation of North Junggar Basin, Xinjiang. *Vertebrata Palasiatica* 47:208–233.

IV. Resumen y conclusiones

Capítulo 11

Resumen y conclusiones



RESUMEN Y CONCLUSIONES

"I am turned into a sort of machine for observing facts and grinding out conclusions."

Charles Darwin

El trabajo realizado ha permitido determinar el material estudiado a nivel específico, conocer la distribución estratigráfica de todas las especies reconocidas y evaluar su utilidad como marcadores bioestratigráficos, así como estudiar sus relaciones filogenéticas y sus distribuciones paleobiogeográficas.

En esta tesis doctoral se han estudiado morfométricamente 3.181 piezas dentales pertenecientes a 54 yacimientos datados desde el Aragoniense medio hasta el principio del Vallesiense (Mioceno medio y superior).

El estudio morfológico está basado en la observación y el análisis de 82 caracteres cuya información se ha recogido en una base de datos. Han sido calculadas las frecuencias de cada carácter en todos los yacimientos, permitiendo la comparación entre distintas localidades. Hemos tomado las medidas de longitud y anchura en los casos en los que la preservación del fósil lo permitía. Las medidas se han incluido en la base de datos y se han calculado los estadísticos descriptivos que han permitido comparar las poblaciones de cada yacimiento.

Se ha estudiado si las muestras de cada localidad presentaban uno o varios grupos morfológicos o métricos. Seguidamente, se ha caracterizado la variabilidad morfológica y métrica de cada uno de los grupos y utilizando toda esta información obtenida, se ha asignado taxonómicamente cada grupo morfológico.

Esta parte se ha acompañado de una revisión de las especies encontradas, enmendando las diagnósticos originales y diferenciales de cuatro de ellas (*C. soriae*, *C. jotae*, *H. nombrevillae* e *H. aragonensis*). En un caso, los fósiles encontrados reunían las suficientes características taxonómicas únicas como para describir con ellos una nueva especie (*C. nievei*).

Los patrones morfológicos de los Cricetodontini de la Cuenca de Calatayud-Daroca se ajustan al siguiente modelo: morfología basal en las primeras especies registradas durante el Aragoniense medio (*C. soriae*, *C. sansaniensis*), configuración en mosaico en la mayor parte de las especies del Aragoniense superior (*C. jotae*, *C. nievei*, *H. aguirrei* e *H. lavocati*) y finalmente, se han observado formas más derivadas en la parte final del Aragoniense superior y en el Vallesiense inferior (*H. nombrevillae*, *H. aragonensis*). A continuación se detallan y explican las conclusiones obtenidas.

CONCLUSIONES TAXONÓMICAS

El estudio taxonómico concluye que los fósiles se reparten entre ocho especies pertenecientes a dos géneros de la tribu. Los resultados se han presentado en los capítulos 5-8. En las siguientes líneas se resumen las cuestiones más importantes de cada uno de ellos.

Capítulo 5. Revisión de *Cricetodon soriae* (Rodentia, Mammalia), nuevos datos del Aragoniense medio (Mioceno medio) de la cuenca de Calatayud-Daroca (Zaragoza, Spain)

Cricetodon soriae es la especie más antigua de la tribu Cricetodontini en la cuenca y, además, es el miembro más antiguo de la tribu en la Península Ibérica. Está presente en la biozona E, entre los 14,06 Ma y los 13,80 Ma. Fue descrita en la localidad de Somosaguas, en la Cuenca de Madrid sobre un total de 8 piezas dentales. Después de este trabajo *Cricetodon soriae* se encuentra en ocho yacimientos más y el número de especímenes que se conocen ha aumentado en 52. Se ha ampliado la distribución geográfica de la especie y ahora se encuentra, además de en la Cuenca de Madrid, también en Calatayud-Daroca.

El estudio de *C. soriae* de la Cuenca de Calatayud-Daroca permite completar la información morfológica y métrica de la especie, así como enmendar su diagnosis original y proponer una nueva diagnosis diferencial más detallada. Los rasgos morfológicos más importantes son: ausencia de estilos y estilidos, bajo desarrollo de los mesolofos y mesolódifos, presencia de protolofulos anteriores y posteriores en el M1 y la ausencia del ectolofa anterior en el M1 y M3; posterolofa bien desarrollado en el M3; presencia de metalofúlidos I y II juntos, así como metalofúlido II sólo; presencia de metalofúlido I incompleto; m3 más largo que el m1; gran tamaño del m3 y los mesolódifos largos y dobles en el m3.

Por otra parte, las comparaciones con el resto de especies de *Cricetodon*, incluyendo los

primeros representantes de Europa Central y Oriental, indican una semejanza morfométrica entre *C. soriae* y las especies de *Cricetodon* de Anatolia. Estas semejanzas son: presencia de protolófulo anterior en el M1 y el hecho de que el m3 sea más largo que el m1, así como la presencia de un metalofúlido I incompleto en el m1.

Capítulo 6. *Cricetodon* del Aragoniense medio y superior de la Cuenca de Calatayud-Daroca: una descripción taxonómica.

En este capítulo se han descrito y discutido dos especies de *Cricetodon*: *C. sansaniensis* y *C. jotae*.

Cricetodon sansaniensis está representado por 89 ejemplares provenientes de cinco localidades pertenecientes a la biozona local F y que abarcan un intervalo temporal de 13,76 a 13,68 Ma. Este estudio permite caracterizar el material, que había sido previamente publicado en nomenclatura abierta. *Cricetodon sansaniensis* de la Cuenca de Calatayud-Daroca se describe como muy homogéneo morfológicamente con ectolofos débiles y ambos metalofúlidos, I y II, en el mismo m1, además de tener mesolofos y mesolófidos. Todos estos rasgos se consideran basales dentro de la tribu.

Se trata de una especie de gran tamaño y no se han observado grandes diferencias en talla entre los yacimientos estudiados.

Se han estudiado 414 molares que corresponden a *C. jotae*. Estos fósiles pertenecen a 13 yacimientos situados en las biozonas locales, G1 y G2 (MN6 *pro parte*), así como en la parte inferior de la G3 (MN7/8 pp.). La primera presencia de *C. jotae* se produce en Las Planas 5B (biozona local G1), datada en 13,56 Ma. La última presencia datada numéricamente de *C. jotae* es en Las Planas 5K (13,08 Ma; biozona local G2). Sin embargo, *Cricetodon jotae* está presente en otros yacimientos más recientes que pertenecen a la biozona local G3, de los cuales no se dispone de edad numérica.

C. jotae presenta una combinación de caracteres basales, como es el metalofúlido II, y derivados—ausencia de mesolofos, presencia de ectolofos dobles; mesolófidos poco desarrollados o ausentes; ausencia de ectomesolófido y anterolófido labial del m1 débil o ausente—que se ha reconocido como un patrón morfológico en mosaico.

A pesar de su amplia distribución estratigráfica, no se han observado diferencias de talla entre las asociaciones a lo largo del tiempo.

Cuando se ha comparado *C. jotae* con otras especies de Cricetodontini se ha observado que

guarda similitudes con *C. meini* y con *H. aguirrei*, ambas presentes en Europa occidental.

Capítulo 7. Nuevo *Cricetodontini* del Mioceno medio de Europa: un ejemplo de evolución en mosaico.

La nueva especie *Cricetodon nievei* se ha encontrado en los yacimientos de Toril2, 3A, 3B y Las Planas 5H, (12,65-12,60 Ma; biozona G3). Esta especie se caracteriza por la combinación de caracteres primitivos y derivados. Los molares superiores tienen las siguientes características: coronas bajas, contorno sinuoso, ectolofos medianamente desarrollados que no forman una pared continua, mesolofos y entomesolofos, estilos y formaciones cingulares en los valles, M1 con cuatro raíces, y M3 apenas reducido. Estas características se presentan con frecuencia en *Cricetodon* y son consideradas como primitivas en la tribu. Por otro lado, los molares inferiores tienen mesolofidos cortos o ausentes, metalofúlidos II ausentes en el m1, y la raíz posterior del m2 parcialmente dividida. Estos caracteres se describen en muchas especies de *Hispanomys* y se consideran como derivados dentro de la tribu *Cricetodontini*. Sin embargo, otros caracteres, tales como la presencia de metalofúlido I y la ausencia de metalofúlido II en el m1, así como tres raíces en los m2 también se encuentran en algunas especies de *Cricetodon*. Por lo tanto hemos decidido incluir esta especie en el género *Cricetodon*. Aun así, se ha considerado necesario incluir a las especies de *Hispanomys* en la diagnosis diferencial.

En cuanto al tamaño, los resultados del análisis métrico muestran que no hay diferencias significativas entre las tallas de las muestras estudiadas.

Después de comparar con el resto de las especies europeas y asiáticas de *Cricetodon* e *Hispanomys* se concluye que *Cricetodon nievei* es más parecido a las especies de *Cricetodon* e *Hispanomys* de la Cuenca de Vallès-Penedès, Francia y Europa Central que a las especies registradas en Calatayud-Daroca.

Capítulo 8. Especies de *Hispanomys* del Aragoniense superior y Vallesiense inferior (Mioceno medio y superior) de la Cuenca de Calatayud-Daroca.

Los 2.359 fósiles estudiados en este capítulo han sido asignados a cuatro especies diferentes de *Hispanomys*. Dos de ellas, *H. aguirrei* e *H. lavocati* se han encontrado en el Aragoniense superior; una, *H. aragonensis*, en el Vallesiense inferior, mientras que *H. nombrevillae* está presente tanto en el Aragoniense superior como en el Vallesiense inferior.

El análisis morfológico de *H. aguirrei* muestra que esta especie también posee un patrón

morfológico en mosaico: no tiene el anterocono totalmente dividido, que es un carácter basal, pero presenta caracteres derivados como puede ser el metalofúlido I en el m1.

Se ha comprobado que las tallas de las localidades más modernas son menores que las de las más antiguas. Esto se ha interpretado como una tendencia hacia una disminución de la talla dentro de *H. aguirrei*.

La comparación con otras especies de *Hispanomys* ha concluido que *H. aguirrei* tiene grandes similitudes con otras especies de la cuenca como *C. jotae* e *H. lavocati*.

Hispanomys lavocati se ha incluido en el género *Hispanomys* a pesar de las propuestas de otros autores, que lo incluyen en *Cricetodon*. *Hispanomys lavocati*, posee caracteres basales como son la presencia de metalofúlido II en el m1, mientras que también presenta otros derivados como la alta frecuencia de ectolofos dobles y conectados.

Hispanomys lavocati es una de las especies más grandes de *Hispanomys* descritas y la más grande de la Cuenca de Calatayud-Daroca para este rango temporal.

La comparación con otras especies de Cricetodontini manifiesta que posee algunas características compartidas con los representantes de *Cricetodon* y *Byzantinia* de Anatolia, como son el gran tamaño del anterocono del M1 y el pequeño cíngulo situado entre los lóbulos del mismo en el M1. Estas similitudes apuntan hacia una posible relación de *H. lavocati* con los miembros de Asia Menor.

Las otras dos especies estudiadas en este capítulo son *H. nombrevillae* e *H. aragonensis*, que tienen todos los rasgos característicos de *Hispanomys*. Su estudio ha permitido enmendar las diagnósis original y diferencial de ambas especies.

Hispanomys nombrevillae presenta características claramente derivadas para la tribu, como son: hipsodoncia moderada; pequeño tamaño; M3 muy reducido; ausencia de metalofúlido II en el m1 y tres raíces en el m2. Sin embargo, estas mismas características denotan una condición basal dentro del género *Hispanomys*.

No hay grandes variaciones de talla entre los fósiles de las localidades estudiadas, por lo que no se han distinguido tendencias al aumento o reducción del tamaño.

La comparación entre *Hispanomys nombrevillae* y otros miembros de la tribu indica varias similitudes con las otras especies de la cuenca, así como con *H. bijugatus* (descrita en el sur de Francia) con las que podría estar relacionada.

Hispanomys aragonensis presenta una gran variabilidad intraespecífica. La morfología parece

seguir una tendencia hacia un patrón más simple. Así, el material de las localidades más antiguas tiene los valles más frecuentemente cerrados y, además, por estructuras más grandes que en el material de las localidades más modernas. Igualmente, la muestra de las localidades más antiguas presenta mesolofos y mesolófidos con más frecuencia que los fósiles de los yacimientos más modernos. Además, los mesolófidos con también más largos en los especímenes de los yacimientos más antiguos.

Hay diferencias de talla entre los especímenes de las localidades más antiguas (que son más grandes) y los de las más modernas (que son más pequeñas).

Los resultados de la comparación con otras especies indican que *H. decedens* e *H. peralensis* son las especies más parecidas a *H. aragonensis* y con las que ésta podría guardar algún tipo de relación filogenética.

Comparando *H. nombrevillae* e *H. aragonensis*, es reconocible que *H. nombrevillae* posee rasgos basales (como la presencia de estructuras que cierran los valles o la raíz posterior del m2 parcialmente dividida en algún ejemplar), mientras que *H. aragonensis* es más derivada (contorno trilobulado, ausencia de posterolofos en el M3). En cuanto al tamaño, las medidas muestran que *H. aragonensis* es más pequeña que *H. nombrevillae*. Esta diferencia en la talla se ha comprobado estadísticamente al encontrar diferencias significativas en la talla de todos los elementos dentales, ya sea en longitud o en anchura.

CONCLUSIONES BIOESTRATIGRÁFICAS

Capítulo 9. Cricetodontini de la Cuenca de Calatayud-Daroca (España): una puesta al día de sus distribuciones estratigráficas.

La tribu Cricetodontini está presente en las biozonas locales desde la E hasta la I, con un rango de edad estimada de 14,06 a 9,98 Ma. Las distribuciones estratigráficas de las especies estudiadas, en general, se correlacionan con la biozonación local, por lo que se puede indicar que se trata de buenos indicadores bioestratigráficos. Las distribuciones de *C. soriae*, *C. sansaniensis*, *H. nombrevillae* e *H. aragonensis* coinciden con las biozonas locales E, F, I y H, respectivamente. Sin embargo, la distribución de *Cricetodon jotae* es muy amplia y oscila alrededor de un 1 Ma cubriendo varias biozonas locales (G1, G2 y G3). Por su parte, *C. nievei* se encuentra únicamente en la parte media de la G3.

Cuatro especies de Cricetodontini están presentes en la biozona local G3. Llama la atención

que esta biozona es muy larga; las biozonas locales aquí estudiadas cubren, cada una, alrededor de 200 o 300 Ka, mientras que la G3 abarca más de 1 Ma. Teniendo en cuenta que las especies de Cricetodontini son buenos indicadores bioestratigráficos, el número de especies encontradas y la gran duración de la biozona local G3, podría resultar conveniente hacer una revisión de esta biozona o una subdivisión en subzonas. No obstante, es recomendable realizar un estudio completo de la fauna de roedores de la G3 antes de hacer una nueva propuesta.

CONCLUSIONES FILOGENÉTICAS

Capítulo 10. Primera filogenia de Cricetodontini durante el Mioceno y Plioceno en Europa y Asia Menor.

Las especies estudiadas se dividen en tres clados principales que corresponden a *Deperetomys*, *Byzantinia* e *Hispanomys*+*Ruscinomys*. El cladograma indica que *Deperetomys* y *Byzantinia* son monofiléticos, *Cricetodon* es parafilético y que las especies de *Hispanomys* forman un clado junto con las de *Ruscinomys*.

A partir del cladograma datado se pueden sacar una serie de conclusiones paleobiogeográficas, como la confirmación de que el centro de origen de Cricetodontini es claramente Asia Menor. También que *Byzantinia* e *Hispanomys* están estrechamente vinculados con un área geográfica específica: Asia Menor y el suroeste de Europa respectivamente. Por otra parte, *Deperetomys* se extiende por Asia Menor y Europa Central, y *Cricetodon* está ampliamente extendido por toda Europa y Asia Menor. También se han detectado tres momentos migratorios (hace 18 Ma, 15 Ma y 14 Ma respectivamente) que tienen su origen en Anatolia y en los que participan diferentes especies que dan lugar a las encontradas en la Cuenca de Calatayud-Daroca.

Asimismo, se han podido describir los principales cambios morfológicos que presentan, en general, los géneros de Cricetodontini. El patrón morfológico deriva hacia la simplicidad de la superficie oclusal: a partir de formas braquiodontas y bunodontas con alta densidad de crestas transversales se observan formas hipsodontas y parecidas a molares selenodontos con prevalencia de crestas longitudinales.

Una vez realizado el análisis cladístico en el que se han analizado 103 caracteres de 51 especies, se han podido proponer una serie de relaciones filogenéticas. Entre éstas, se han discutido con más detalle las relaciones filogenéticas de cada una de las especies de Cuenca de Calatayud-Daroca, estudiadas en la presente tesis doctoral.

- *Cricetodon soriae* está estrechamente relacionado con otras especies de Europa occidental, como son *C. sansaniensis*, *C. jotae* y *C. meini*. Su relación con las especies de Anatolia no parece ser tan evidente como se había propuesto en el artículo que sirve de base para el capítulo 5.
- *Cricetodon sansaniensis* no parece tener relación con ninguna de las especies con las que se había relacionado anteriormente y se rechaza su relación directa con *C. aureus*. En cambio, sí parece tener una relación estrecha con *C. soriae*.
- *Cricetodon jotae* es grupo hermano de *C. meini*, lo que confirma la relación propuesta por los autores previamente y discutida en el capítulo 6.
- *Cricetodon nievei* está agrupada con las especies procedentes de Europa central, claramente separada de los otros Cricetodontini de la Cuenca de Calatayud-Daroca, lo que confirma la propuesta realizada tras el estudio sistemático efectuado en el capítulo 7.

Las especies de *Hispanomys* se dividen entre dos clados monofiléticos: *H. aguirrei*-*H. lavocati* se encuentran en un clado e *H. nombrevillae*-*H. aragonensis* en otro. El primer clado está situado en una posición más basal que el segundo.

- *Hispanomys aguirrei* se asigna a *Hispanomys*, ya que se encuentra entre las otras especies de este género. Parece confirmarse que *H. aguirrei* no está relacionada con *C. jotae*, como se explica en los capítulos 6 y 8, sino con *H. lavocati*, su especie hermana.
- *Hispanomys lavocati* está también asignado a *Hispanomys* y no parece tener una relación directa con las especies de *Byzantinia* (que aparecen en otro clado separado), ni con las especies grandes de *Cricetodon*.
- *Hispanomys nombrevillae* aparece como una especie basal al clado que contiene todas las especies derivadas de *Hispanomys*. Esto apoya la discusión realizada en el capítulo 8 en la que se explica que *H. nombrevillae* posee caracteres considerados basales dentro del género.
- *Hispanomys aragonensis* se encuentra entre las especies de *Hispanomys* más derivadas, el cladograma rechaza su relación directa con *H. decedens*. Se relaciona directamente con *H. peralensis*, apoyando la propuesta descrita en el capítulo 8.

Con este estudio se ha aumentado considerablemente la información disponible de los miembros de esta tribu. Sin embargo, sobre todo en los aspectos bioestratigráficos y filogenéticos, todavía hay cuestiones que posiblemente se solucionen estudiando en profundidad otros grupos de roedores.

De esta forma, las futuras líneas de investigación en el campo de la bioestratigrafía local deben enfocarse hacia el resto de géneros de roedores del Aragoniense superior y del Vallesiense inferior de la Cuenca de Calatayud-Daroca. Por otro lado, sería interesante ampliar el estudio filogenético con información morfológica del resto de géneros de Cricetodontini y otros géneros de Asia Menor. De este modo se podrían resolver las cuestiones en géneros tales como *Deperetomys*, *Cricetodon* y *Ruscinomys*.

Durante el desarrollo de esta tesis doctoral la identificación de las especies de la tribu ha sido difícil debido que no se disponía de criterios suficientemente claros en las diagnósicos de algunas especies descritas anteriormente. Para encontrar estos criterios, se ha tenido que profundizar en el estudio de la morfología dental.

Por otro lado, en ocasiones, a pesar de tener una gran cantidad de fósiles, se ha contado con pocas piezas. 3.150 fósiles repartidos entre 8 especies es un número muy bajo si se compara con el material de otros géneros que estos mismos yacimientos presentan. La solución para poder resolver la taxonomía ha pasado por ampliar el número de variables morfológicas observadas y por completar la observación con material de otros yacimientos de fuera de la cuenca. Para eso ha sido fundamental la recogida de datos de Austria, Alemania y Francia. De este modo, se ha podido distinguir qué caracteres presentaban una mayor variabilidad geográfica y cuáles eran más homogéneos y por lo tanto, más útiles taxonómicamente.

Esta observación de las localidades y especies de fuera de la cuenca ha abierto las puertas a futuros trabajos. Con todos los datos recopilados en una base de datos morfológica de poblaciones de toda Europa, se pueden realizar estudios paleofilogeográficos que precisen cómo se relacionan filogenéticamente las poblaciones de Cricetodontini entre sí, si se agrupan de alguna forma y si esa agrupación se corresponde con la especie a la que esas poblaciones están asignadas. Igualmente, un estudio paleofilogeográfico nos permitiría caracterizar la variabilidad intraespecífica y determinar de una manera más efectiva qué caracteres son más diagnósticos dentro de una especie.

Las metodologías aplicadas, en especial la cladística, pueden ser el motor para llevar a cabo estas investigaciones en paleofilogeografía, aunque un trabajo de este tipo necesita nuevas técnicas y metodologías. Además de ayudar en las cuestiones anteriormente mencionadas (describir las relaciones filogenéticas y la variabilidad intraespecífica), la paleofilogeografía podría utilizarse para apoyar la asignación taxonómica de poblaciones con poco material fósil.

Pese a la amplitud e interés de la paleontología descriptiva, cada vez son más los trabajos de paleontología analítica y esta tesis doctoral es un punto de partida para éstos. Se ha recopilado una inmensa cantidad de datos y ahora hay que preguntarse qué más se puede hacer con ellos y cómo hacerlo. Estas preguntas no hacen más que abrir nuevos retos que van hacia el concepto más amplio e integrador de la paleontología.



Las ilustraciones utilizadas en esta memoria son creaciones de: Melissa van de Paardt, Wayne Anderson, Andreas Deja, Brian Wildsmith, Sebas Roa, Norbet Martin y Beatrix Potter. También se han utilizado ilustraciones del CD-ROM *Full color animals illustrations*, Editado por Dover publications, ISBN: 9780486995461.